Jordana Georgin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7991132/publications.pdf

Version: 2024-02-01

218662 254170 2,098 87 26 43 citations g-index h-index papers 88 88 88 1137 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Application of fly ash modified by alkaline fusion as an effective adsorbent to remove methyl violet 10B in water. Chemical Engineering Communications, 2022, 209, 184-195.	2.6	4
2	Effective adsorptive removal of textile pollutant using coal bottom ash with high surface area obtained by alkaline fusion route. Environmental Technology (United Kingdom), 2022, 43, 2418-2429.	2.2	4
3	Optimization of ketoprofen adsorption from aqueous solutions and simulated effluents using H2SO4 activated Campomanesia guazumifolia bark. Environmental Science and Pollution Research, 2022, 29, 2122-2135.	5.3	6
4	Effective adsorptive removal of atrazine herbicide in river waters by a novel hydrochar derived from Prunus serrulata bark. Environmental Science and Pollution Research, 2022, 29, 3672-3685.	5.3	22
5	Woody residues of the grape production chain as an alternative precursor of high porous activated carbon with remarkable performance for naproxen uptake from water. Environmental Science and Pollution Research, 2022, 29, 16988-17000.	5.3	4
6	Applicability of amethyst mining rejects as a novel photo-fenton catalyst for the abatement of an emerging pollutant in water. Applied Geochemistry, 2022, 136, 105136.	3.0	8
7	A study of single and quaternary adsorption of Cu2+, Co2+, Ni2+ and Ag+ on sludge modified by alkaline fusion. Chemical Engineering Journal, 2022, 433, 133674.	12.7	7
8	Adsorption of atrazine and 2,4-D pesticides on alternative biochars from cedar bark sawdust (Cedrella) Tj ETQq0	0 <u>0 ;</u> gBT /	Overlock 10 T
9	Development of activated carbon from Schizolobium parahyba (guapuruvu) residues employed for the removal of ketoprofen. Environmental Science and Pollution Research, 2022, 29, 21860-21875.	5.3	3
10	Adsorption of atrazine herbicide from water by diospyros kaki fruit waste activated carbon. Journal of Molecular Liquids, 2022, 347, 117990.	4.9	27
11	Residual peel of pitaya fruit (Hylocereus undatus) as a precursor to obtaining an efficient carbon-based adsorbent for the removal of metanil yellow dye from water. Journal of Environmental Chemical Engineering, 2022, 10, 107006.	6.7	19
12	Effective removal of non-steroidal anti-inflammatory drug from wastewater by adsorption process using acid-treated Fagopyrum esculentum husk. Environmental Science and Pollution Research, 2022, 29, 31085-31098.	5 . 3	4
13	Application of biowaste generated by the production chain of pitaya fruit (Hylocereus undatus) as an efficient adsorbent for removal of naproxen in water. Environmental Science and Pollution Research, 2022, 29, 39754-39767.	5.3	5
14	Application of ara \tilde{A} § \tilde{A}_i fruit husks (Psidium cattleianum) in the preparation of activated carbon with FeCl3 for atrazine herbicide adsorption. Chemical Engineering Research and Design, 2022, 180, 67-78.	5.6	24
15	Investigation of biochar from Cedrella fissilis applied to the adsorption of atrazine herbicide from an aqueous medium. Journal of Environmental Chemical Engineering, 2022, 10, 107408.	6.7	36
16	Preparation of activated carbons from fruit residues for the removal of naproxen (NPX): Analytical interpretation via statistical physical model. Journal of Molecular Liquids, 2022, 356, 119021.	4.9	22
17	A review of the toxicology presence and removal of ketoprofen through adsorption technology. Journal of Environmental Chemical Engineering, 2022, 10, 107798.	6.7	29
18	Pore volume and surface diffusion model (PVSDM) applied for single and binary dye adsorption systems. Chemical Engineering Research and Design, 2022, 182, 645-658.	5.6	5

#	Article	IF	CITATIONS
19	Enhanced adsorption of ketoprofen and 2,4-dichlorophenoxyactic acid on Physalis peruviana fruit residue functionalized with H2SO4: Adsorption properties and statistical physics modeling. Chemical Engineering Journal, 2022, 445, 136773.	12.7	22
20	Conversion of Erythrina speciosa pods to porous adsorbent for Ibuprofen removal. Journal of Environmental Chemical Engineering, 2022, 10, 108070.	6.7	13
21	An advanced combination of density functional theory simulations and statistical physics modeling in the unveiling and prediction of adsorption mechanisms of 2,4-D pesticide to activated carbon. Journal of Molecular Liquids, 2022, 361, 119639.	4.9	21
22	Advances made in removing paraquat herbicide by adsorption technology: A review. Journal of Water Process Engineering, 2022, 49, 102988.	5 . 6	26
23	Application of Cordia trichotoma sawdust as an effective biosorbent for removal of crystal violet from aqueous solution in batch system and fixed-bed column. Environmental Science and Pollution Research, 2021, 28, 6771-6783.	5. 3	26
24	Transforming pods of the species Capparis flexuosa into effective biosorbent to remove blue methylene and bright blue in discontinuous and continuous systems. Environmental Science and Pollution Research, 2021, 28, 8036-8049.	5. 3	5
25	Successful adsorption of bright blue and methylene blue on modified pods of Caesalpinia echinata in discontinuous system. Environmental Science and Pollution Research, 2021, 28, 8407-8420.	5. 3	12
26	High-performance removal of 2,4-dichlorophenoxyacetic acid herbicide in water using activated carbon derived from Queen palm fruit endocarp (Syagrus romanzoffiana). Journal of Environmental Chemical Engineering, 2021, 9, 104911.	6.7	79
27	Transforming shrub waste into a high-efficiency adsorbent: Application of Physalis peruvian chalice treated with strong acid to remove the 2,4-dichlorophenoxyacetic acid herbicide. Journal of Environmental Chemical Engineering, 2021, 9, 104574.	6.7	56
28	Macro-fungal (Agaricus bisporus) wastes as an adsorbent in the removal of the acid red 97 and crystal violet dyes from ideal colored effluents. Environmental Science and Pollution Research, 2021, 28, 405-415.	5. 3	24
29	Application of seed residues from Anadenanthera macrocarpa and Cedrela fissilis as alternative adsorbents for remarkable removal of methylene blue dye in aqueous solutions. Environmental Science and Pollution Research, 2021, 28, 2342-2354.	5.3	23
30	Adsorption investigation of 2,4-D herbicide on acid-treated peanut (Arachis hypogaea) skins. Environmental Science and Pollution Research, 2021, 28, 36453-36463.	5. 3	14
31	Conversion of the forest species Inga marginata and Tipuana tipu wastes into biosorbents: Dye biosorption study from isotherm to mass transfer. Environmental Technology and Innovation, 2021, 22, 101521.	6.1	10
32	An overview of forest residues as promising low-cost adsorbents. Gondwana Research, 2021, , .	6.0	14
33	Development of highly porous activated carbon from Jacaranda mimosifolia seed pods for remarkable removal of aqueous-phase ketoprofen. Journal of Environmental Chemical Engineering, 2021, 9, 105676.	6.7	54
34	Adsorption of ketoprofen and paracetamol and treatment of a synthetic mixture by novel porous carbon derived from Butia capitata endocarp. Journal of Molecular Liquids, 2021, 339, 117184.	4.9	73
35	Highly effective adsorption of synthetic phenol effluent by a novel activated carbon prepared from fruit wastes of the Ceiba speciosa forest species. Journal of Environmental Chemical Engineering, 2021, 9, 105927.	6.7	51
36	Adsorption and mass transfer studies of methylene blue onto comminuted seedpods from Luehea divaricata and Inga laurina. Environmental Science and Pollution Research, 2021, 28, 20854-20868.	5. 3	8

#	Article	IF	CITATIONS
37	One step acid modification of the residual bark from <i>Campomanesia guazumifolia</i> i> using H ₂ SO ₄ and application in the removal of 2,4-dichlorophenoxyacetic from aqueous solution. Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes, 2021, 56, 995-1006.	1.5	2
38	Transforming agricultural waste into adsorbent: application of Fagopyrum esculentum wheat husks treated with H2SO4 to adsorption of the 2,4-D herbicide. Journal of Environmental Chemical Engineering, 2021, 9, 106872.	6.7	22
39	Preparation of activated carbon from the residues of the mushroom (Agaricus bisporus) production chain for the adsorption of the 2,4-dichlorophenoxyacetic herbicide. Journal of Environmental Chemical Engineering, 2021, 9, 106843.	6.7	47
40	Efficient removal of naproxen from aqueous solution by highly porous activated carbon produced from Grapetree (Plinia cauliflora) fruit peels. Journal of Environmental Chemical Engineering, 2021, 9, 106820.	6.7	24
41	Paddle cactus (Tacinga palmadora) as potential low-cost adsorbent to treat textile effluents containing crystal violet. Chemical Engineering Communications, 2020, 207, 1368-1379.	2.6	16
42	Water treatment plant sludge as iron source to catalyze a heterogeneous photo-Fenton reaction. Environmental Technology and Innovation, 2020, 17, 100544.	6.1	38
43	Preparation and characterization of a novel mountain soursop seeds powder adsorbent and its application for the removal of crystal violet and methylene blue from aqueous solutions. Chemical Engineering Journal, 2020, 391, 123617.	12.7	70
44	A mass transfer study considering intraparticle diffusion and axial dispersion for fixed-bed adsorption of crystal violet on pecan pericarp (Carya illinoensis). Chemical Engineering Journal, 2020, 397, 125423.	12.7	52
45	Utilization of Pacara Earpod tree (Enterolobium contortisilquum) and Ironwood (Caesalpinia) Tj ETQq1 1 0.784 Pollution Research, 2020, 27, 33307-33320.	1314 rgBT /(5.3	Overlock 10 T 59
46	Adsorption of hazardous dyes on functionalized multiwalled carbon nanotubes in single and binary systems: Experimental study and physicochemical interpretation of the adsorption mechanism. Chemical Engineering Journal, 2020, 389, 124467.	12.7	125
47	Araticum (Annona crassiflora) seed powder (ASP) for the treatment of colored effluents by biosorption. Environmental Science and Pollution Research, 2020, 27, 11184-11194.	5.3	28
48	Powdered biosorbent from the mandacaru cactus (cereus jamacaru) for discontinuous and continuous removal of Basic Fuchsin from aqueous solutions. Powder Technology, 2020, 364, 584-592.	4.2	47
49	Solid wastes from the enzyme production as a potential biosorbent to treat colored effluents containing crystal violet dye. Environmental Science and Pollution Research, 2020, 27, 10484-10494.	5 . 3	15
50	Evaluation of Ocotea puberula bark powder (OPBP) as an effective adsorbent to uptake crystal violet from colored effluents: alternative kinetic approaches. Environmental Science and Pollution Research, 2020, 27, 25727-25739.	5. 3	27
51	Treatment of water containing methylene by biosorption using Brazilian berry seeds (Eugenia) Tj ETQq $1\ 1\ 0.78$	34314.rgBT /	Overlock 10
52	Insights of the adsorption mechanism of methylene blue on brazilian berries seeds: Experiments, phenomenological modelling and DFT calculations. Chemical Engineering Journal, 2020, 394, 125011.	12.7	60
53	Powdered biosorbent from pecan pericarp (Carya illinoensis) as an efficient material to uptake methyl violet 2B from effluents in batch and column operations. Advanced Powder Technology, 2020, 31, 2843-2852.	4.1	40
54	Adsorption of crystal violet on biomasses from pecan nutshell, para chestnut husk, araucaria bark and palm cactus: Experimental study and theoretical modeling via monolayer and double layer statistical physics models. Chemical Engineering Journal, 2019, 378, 122101.	12.7	148

#	Article	IF	Citations
55	Potentiality of the Phoma sp. inactive fungal biomass, a waste from the bioherbicide production, for the treatment of colored effluents. Chemosphere, 2019, 235, 596-605.	8.2	22
56	Adsorptive decontamination of wastewater containing methylene blue dye using golden trumpet tree bark (Handroanthus albus). Environmental Science and Pollution Research, 2019, 26, 31924-31933.	5.3	34
57	Potential of Cedrella fissilis bark as an adsorbent for the removal of red 97 dye from aqueous effluents. Environmental Science and Pollution Research, 2019, 26, 19207-19219.	5.3	50
58	Biosorption of crystal violet dye using inactive biomass of the fungus Diaporthe schini. Water Science and Technology, 2019, 79, 709-717.	2.5	26
59	Residual biomass of Nigrospora sp. from process of the microbial oil extraction for the biosorption of procion red H–E7B dye. Journal of Water Process Engineering, 2019, 31, 100818.	5.6	7
60	Applicability of Coal Bottom Ash from Thermoelectric Power Plant as an Alternative Heterogeneous Catalyst in Photo-Fenton Reaction. Water, Air, and Soil Pollution, 2019, 230, 1.	2.4	13
61	Application of Beauveria bassiana spore waste as adsorbent to uptake acid red 97 dye from aqueous medium. Environmental Science and Pollution Research, 2019, 26, 36967-36977.	5.3	7
62	Preparação, caracterização e avaliação catalÃŧica do compósito Fe2O3/grafite em reação foto-Fentor Revista Materia, 2019, 24, .	٦.	2
63	Biosorption of cationic dyes by Par $ ilde{A}_i$ chestnut husk (Bertholletia excelsa). Water Science and Technology, 2018, 77, 1612-1621.	2.5	48
64	Removal of Procion Red dye from colored effluents using H2SO4-/HNO3-treated avocado shells (Persea americana) as adsorbent. Environmental Science and Pollution Research, 2018, 25, 6429-6442.	5.3	44
65	Potential of Araucaria angustifolia bark as adsorbent to remove Gentian Violet dye from aqueous effluents. Water Science and Technology, 2018, 78, 1693-1703.	2.5	43
66	Valorização dos resÃduos do processamento da Nóz pecã (Carya illinoensis) como adsorvente dos corantes azul de metileno e violeta cristal. Engevista, 2017, 19, 1449.	0.1	0
67	Preparation of activated carbon from peanut shell by conventional pyrolysis and microwave irradiation-pyrolysis to remove organic dyes from aqueous solutions. Journal of Environmental Chemical Engineering, 2016, 4, 266-275.	6.7	158
68	ESTIMATIVA DO CONSUMO PER CAPITA DE ÃGUA TRATADA PARA USO DOMÉSTICO POR MEIO DE FERRAMENTAS ESTATÃSTICAS: ESTUDO DE CASO DA CIDADE DE ARIQUEMES-RO. , 2016, 64, 32-38.	0.2	1
69	Avaliação dos Componentes de Rendimento do Trigo quando Submetido a Diferentes Fontes de Nitrogênio. Revista Eletrônica Em Gestão EducaçÁ£o E Tecnologia Ambiental, 2016, 20, 524.	0.0	4
70	Avaliação da qualidade da água subterrânea: estudo de caso de Vilhena – RO. Revista Ãguas Subterrâneas, 2015, 29, 213.	0.1	4
71	As modalidades mais frequentes de licenciamento realizadas em municÃpios da região norte do Rio Grande do Sul. Revista Monografias Ambientais, 2015, 14, 136-143.	0.1	O
72	Festas rurais: mÃdia, dimensão festiva e impacto social em duas pequenas comunidades do Rio Grande do Sul. Revista Monografias Ambientais, 2015, 14, 115-122.	0.1	0

#	Article	IF	CITATIONS
73	A participação feminina na agricultura agroecológica: um estudo do caso na região norte do Rio Grande do Sul. Revista Monografias Ambientais, 2015, 14, 01-09.	0.1	2
74	PRÃTICAS DE CONSCIENTIZAÇÃ f O AMBIENTAL EM ESCOLAS PÊBLICAS DE RONDA ALTA/RS. Revista Monografias Ambientais, 2014, 13, .	0.1	1
75	PLANTIO DE PINUS ELLIOTTII EM PEQUENAS PROPRIEDADES RURAIS NO NORTE DO RIO GRANDE DO SUL. Revista Monografias Ambientais, 2014, 13, .	0.1	1
76	ANÃŁISE DO PROGRAMA DE QUALIFICAÇÃO PROFISSIONAL DE AGRICULTORES – EMATER, SOB A PERSPECTI MIDIÃTICA. Revista Monografias Ambientais, 2014, 13, .	VA O.1	0
77	BRASIL: O ACESSO UNIVERSAL AO SANEAMENTO BÃSICO. Revista Monografias Ambientais, 2014, 13, .	0.1	О
78	ANÃŁISE ESTRUTURAL E DISTRIBUIÇÕES DE FREQUÊNCIA EM UM FRAGMENTO FLORESTAL NO MUNICÃPIO DE FREDERICO WESTPHALEN – RS, BRASIL. Revista EletrÃ′nica Em Gestão Educação E Tecnologia Ambiental, 2014, 18, .	0.0	0
79	INDUÇÃO AO ENRAIZAMENTO DE ESTACAS DE TECA (TECTONA GRANDIS L.F). Revista Eletrônica Em Gestão Educação E Tecnologia Ambiental, 2014, 18, .	0.0	2
80	A AGRICULTURA FAMILIAR NO CENÃRIO DA PRODUÇÃ f O DE FLORESTAS ENERGÃ%TICAS. Revista Monografias Ambientais, 2014, 13, .	0.1	0
81	Influência do Teor de Umidade na Germinação de Sementes de Parapiptadenia rigida (Benth.) Brenan. Nativa, 2014, 2, 124-128.	0.4	0
82	ADSORÇÃO DE CORANTES TÊXTEIS UTILIZANDO A CASCA DA CASTANHA DO PARÕ, 0, , .		0
83	USO DE RESÃDUOS DA AGROINDà STRIA NA REMOà ‡Ã JO DO CORANTE VIOLETA CRISTAL., 0, , .		0
84	ADSORÇÃO DE CORANTES TÊXTEIS UTILIZANDO A CASCA DA CASTANHA DO PARÕ, 0, , 9-15.		0
85	USO DE RESÃDUOS DA AGROINDÃĕSTRIA NA REMOÇÃO DO CORANTE VIOLETA. , 0, , 151-157.		O
86	Adsorption of the Paracetamol First-Line Covid Treatment Drug Onto Activated Carbon from Residual Pods of Erythrina Speciosa. SSRN Electronic Journal, 0, , .	0.4	0
87	O ATUAL CONTEXTO DA PRODUÇÃO DE TRIGO NO RIO GRANDE DO SUL. Revista Eletrônica Em Gestão Educação E Tecnologia Ambiental, 0, , 246-257.	0.0	0