

Vasiliki Koliaraki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7992171/publications.pdf>

Version: 2024-02-01

27
papers

1,443
citations

471061

17
h-index

552369

26
g-index

28
all docs

28
docs citations

28
times ranked

2633
citing authors

#	ARTICLE	IF	CITATIONS
1	Col6a1+/CD201+ mesenchymal cells regulate intestinal morphogenesis and homeostasis. Cellular and Molecular Life Sciences, 2022, 79, 1.	2.4	12
2	Fibroblastic reticular cell lineage convergence in Peyer's patches governs intestinal immunity. Nature Immunology, 2021, 22, 510-519.	7.0	35
3	An intrinsic role of IL-33 in Treg cell-mediated tumor immunoevasion. Nature Immunology, 2020, 21, 75-85.	7.0	82
4	The mesenchymal context in inflammation, immunity and cancer. Nature Immunology, 2020, 21, 974-982.	7.0	168
5	Unfolding innate mechanisms in the cancer microenvironment: The emerging role of the mesenchyme. Journal of Experimental Medicine, 2020, 217, .	4.2	11
6	Fibroblast Reprogramming in Gastrointestinal Cancer. Frontiers in Cell and Developmental Biology, 2020, 8, 630.	1.8	19
7	Paracrine orchestration of intestinal tumorigenesis by a mesenchymal niche. Nature, 2020, 580, 524-529.	13.7	183
8	The BACH1-HMOX1 Regulatory Axis Is Indispensable for Proper Macrophage Subtype Specification and Skeletal Muscle Regeneration. Journal of Immunology, 2019, 203, 1532-1547.	0.4	22
9	Innate Sensing through Mesenchymal TLR4/MyD88 Signals Promotes Spontaneous Intestinal Tumorigenesis. Cell Reports, 2019, 26, 536-545.e4.	2.9	38
10	Mesenchymal MAPKAPK2/HSP27 drives intestinal carcinogenesis. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, E5546-E5555.	3.3	29
11	Mesenchymal Cells in Colon Cancer. Gastroenterology, 2017, 152, 964-979.	0.6	158
12	Targeted deletion of RANKL in M cell inducer cells by the Col6a1-Cre driver. Biochemical and Biophysical Research Communications, 2017, 493, 437-443.	1.0	14
13	CollagenVI-Cre mice: A new tool to target stromal cells in secondary lymphoid organs. Scientific Reports, 2016, 6, 33027.	1.6	17
14	Isolation of Intestinal Mesenchymal Cells from Adult Mice. Bio-protocol, 2016, 6, .	0.2	13
15	IKK β in intestinal mesenchymal cells promotes initiation of colitis-associated cancer. Journal of Experimental Medicine, 2015, 212, 2235-2251.	4.2	109
16	IKK β in intestinal mesenchymal cells promotes initiation of colitis-associated cancer. Journal of Cell Biology, 2015, 211, 2115-2127.	2.3	1
17	Intestinal myofibroblast-specific Tpl2-Cox-2-PGE ₂ pathway links innate sensing to epithelial homeostasis. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, E4658-67.	3.3	83
18	Tpl2 regulates intestinal myofibroblast HGF release to suppress colitis-associated tumorigenesis. Journal of Clinical Investigation, 2012, 122, 4231-4242.	3.9	64

#	ARTICLE	IF	CITATIONS
19	A New Role for Myeloid HO-1 in the Innate to Adaptive Crosstalk and Immune Homeostasis. <i>Advances in Experimental Medicine and Biology</i> , 2011, 780, 101-111.	0.8	25
20	Association of haptoglobin genotype and common cardiovascular risk factors with the amount of iron in atherosclerotic carotid plaques. <i>Atherosclerosis</i> , 2011, 216, 131-138.	0.4	29
21	Serum hepcidin levels are related to the severity of liver histological lesions in chronic hepatitis C. <i>Journal of Viral Hepatitis</i> , 2010, 17, 800-806.	1.0	53
22	A Novel Immunological Assay for Heparin Quantification in Human Serum. <i>PLoS ONE</i> , 2009, 4, e4581.	1.1	72
23	Results of the first international round robin for the quantification of urinary and plasma hepcidin assays: need for standardization. <i>Haematologica</i> , 2009, 94, 1748-1752.	1.7	161
24	Iron regulatory and bactericidal properties of human recombinant hepcidin expressed in <i>Pichia pastoris</i> . <i>Biochimie</i> , 2008, 90, 726-735.	1.3	30
25	Results of the First International Round Robin for the Quantification of Urinary and Plasma Heparin: Need for Standardization. <i>Blood</i> , 2008, 112, 120-120.	0.6	7
26	Protein BmP0 from the silkworm <i>Bombyx mori</i> can be assembled and is functional in the <i>Saccharomyces cerevisiae</i> ribosomal stalk in the absence of the acidic P1 and P2 proteins. <i>Gene</i> , 2003, 314, 173-179.	1.0	8
27	Innate Sensing by Mesenchymal TLR4/MyD88 Signals Promotes Spontaneous Intestinal Tumorigenesis. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0