

Leonid E Popov

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7999625/publications.pdf>

Version: 2024-02-01

127
papers

2,428
citations

186209

28
h-index

302012

39
g-index

128
all docs

128
docs citations

128
times ranked

1144
citing authors

#	ARTICLE	IF	CITATIONS
1	Stratigraphic and volcanic signatures of Miaolingian-Late Ordovician rift pulses in the Alborz Mountains, northern Iran. <i>Journal of Asian Earth Sciences</i> , 2022, 233, 105240.	1.0	10
2	Submarine metalliferous carbonate mounds in the Cambrian of the Baltoscandian Basin induced by vent networks and water column stratification. <i>Scientific Reports</i> , 2022, 12, 8475.	1.6	1
3	The oldest Cambrian trilobite “ brachiopod association in South China. <i>Gondwana Research</i> , 2021, 89, 147-167.	3.0	16
4	Cambrian brachiopods from the area of the former Semipalatinsk nuclear-testing site, Chingiz Ranges, Kazakhstan. <i>Palaontologische Zeitschrift</i> , 2021, 95, 275-290.	0.8	2
5	Silurian (Aeronian) rhynchonelliform brachiopods of Shabdjereh, south-west Central Iran and their significance for early spiriferide evolution. <i>Journal of Systematic Palaeontology</i> , 2021, 19, 191-219.	0.6	2
6	The first Early Ordovician graptolites and marine incursions in eastern Alborz, Iran. <i>Geological Magazine</i> , 2021, 158, 1719-1732.	0.9	4
7	The identity and significance of the high-latitude Early Ordovician Mediterranean brachiopod Province. <i>Geological Magazine</i> , 2021, 158, 2187-2208.	0.9	8
8	Linguliform brachiopods from the Cambrian (Guzhangian) Karpinsk Formation of Novaya Zemlya. <i>Papers in Palaeontology</i> , 2020, 6, 571-592.	0.7	2
9	Early Devonian (Lochkovian) trilobites and brachiopods from the west Balkhash Region, Kazakhstan and their significance for regional terrane dynamics. <i>Geobios</i> , 2019, 56, 49-64.	0.7	2
10	Glendonite occurrences in the Tremadocian of Baltica: first Early Palaeozoic evidence of massive ikaite precipitation at temperate latitudes. <i>Scientific Reports</i> , 2019, 9, 7205.	1.6	19
11	Cambrian rhynchonelliform nisusoid brachiopods: phylogeny and distribution. <i>Papers in Palaeontology</i> , 2019, 5, 559-575.	0.7	7
12	Early Ordovician brachiopods from south-west Wales. <i>Proceedings of the Geologists Association</i> , 2019, 130, 677-690.	0.6	8
13	Ordovician cold water brachiopods from the Ougarta Mountain Range, Algerian Sahara. <i>Bulletin of Geosciences</i> , 2019, , 41-70.	0.5	6
14	Evolutionary significance of a middle Cambrian (Series 3) in-situ occurrence of the pedunculate rhynchonelliform brachiopod <i>Nisusia sulcata</i> . <i>Lethaia</i> , 2018, 51, 424-432.	0.6	6
15	Earliest ontogeny of early Cambrian acrotretoid brachiopods “ first evidence for metamorphosis and its implications. <i>BMC Evolutionary Biology</i> , 2018, 18, 42.	3.2	24
16	The attachment strategies of Cambrian kutorginate brachiopods: the curious case of two pedicle openings and their phylogenetic significance. <i>Journal of Paleontology</i> , 2018, 92, 33-39.	0.5	8
17	A paired apatite and calcite clumped isotope thermometry approach to estimating Cambro-Ordovician seawater temperatures and isotopic composition. <i>Geochimica Et Cosmochimica Acta</i> , 2018, 224, 18-41.	1.6	60
18	First report on the Early Devonian (Lochkovian) brachiopods from eastern Central Pamirs, Tajikistan. <i>Journal of Asian Earth Sciences</i> , 2017, 138, 427-438.	1.0	1

#	ARTICLE	IF	CITATIONS
19	Ecology, biofacies, biogeography and systematics of micromorphic lingulate brachiopods from the Ordovician (Darriwilian–Sandbian) of south-central China. <i>Papers in Palaeontology</i> , 2017, 3, 317-361.	0.7	4
20	Unusual pitted Ordovician brachiopods from the East Baltic: the significance of coarsely pitted ornamentations in linguliforms. <i>Papers in Palaeontology</i> , 2017, 3, 387-399.	0.7	5
21	The World's second oldest strophomenoid-dominated benthic assemblage in the first Dapingian (Middle Ordovician) brachiopod fauna identified from Iran. <i>Journal of Asian Earth Sciences</i> , 2017, 140, 1-12.	1.0	10
22	Earliest ontogeny of the Silurian orthotetide brachiopod <i>Coolinia</i> and its significance for interpreting strophomenate phylogeny. <i>Lethaia</i> , 2017, 50, 504-510.	0.6	7
23	Late Ordovician palaeogeography and the positions of the Kazakh terranes through analysis of their brachiopod faunas. <i>Acta Geologica Polonica</i> , 2017, 67, 323-380.	0.9	31
24	A new orthide brachiopod genus from the Middle Ordovician of the Alborz Mountains, Iran. <i>Gff</i> , 2017, 139, 327-332.	0.4	0
25	Silurian stratigraphy of Central Iran – an update. <i>Acta Geologica Polonica</i> , 2017, 67, 201-233.	0.9	8
26	Brachiopods: origin and early history. <i>Palaeontology</i> , 2017, 60, 609-631.	1.0	39
27	First occurrence of <i>Lesueurilla prima</i> (Gastropoda) from the Middle Ordovician (Darriwilian) of Iran. <i>Gff</i> , 2016, 138, 510-518.	0.4	6
28	Reassessment of the early Triassic lingulid brachiopod <i>Lingula borealis</i> Bittner, 1899 and related problems of lingulid taxonomy. <i>Gff</i> , 2016, 138, 519-525.	0.4	5
29	Review of the Ordovician stratigraphy and fauna of the Anarak Region in Central Iran. <i>Acta Geologica Polonica</i> , 2015, 65, 403-435.	0.9	9
30	Himalayan Cambrian brachiopods. <i>Papers in Palaeontology</i> , 2015, 1, 345-399.	0.7	52
31	First record of the Ordovician fauna in Mila-Kuh, eastern Alborz, northern Iran. <i>Estonian Journal of Earth Sciences</i> , 2015, 64, 121.	0.4	9
32	Taxonomy, morphology, shell structure and early ontogeny of <i>Pelmanotreta</i> nom. nov. from the lower Cambrian of Siberia. <i>Gff</i> , 2015, 137, 1-8.	0.4	15
33	Late Ordovician cephalopods, tentaculitides, machaeridians and echinoderms from Kuh-e Faraghan, High Zagros, Iran. <i>Alcheringa</i> , 2015, 39, 530-549.	0.5	12
34	First reported Late Ordovician trilobites from the High Zagros Ranges, Iran: A biogeographic link between Gondwanan Chinese and Mediterranean faunas. <i>Geobios</i> , 2015, 48, 351-369.	0.7	13
35	An Early Silurian (Aeronian) cephalopod fauna from Kopet-Dagh, north-eastern Iran: including the earliest records of non-orthocerid cephalopods from the Silurian of Northern Gondwana. <i>Bulletin of Geosciences</i> , 2015, , 479-507.	0.5	4
36	Late Ordovician brachiopods from the Chingiz Terrane, Kazakhstan, and their palaeogeography. <i>Journal of Systematic Palaeontology</i> , 2014, 12, 687-758.	0.6	20

#	ARTICLE	IF	CITATIONS
37	A diverse Upper Darriwilian radiolarian assemblage from the Shundy Formation of Kazakhstan: insights into late Middle Ordovician radiolarian biodiversity. <i>Journal of Micropalaeontology</i> , 2014, 33, 149-163.	1.3	11
38	Ordovician "Silurian Chileida" First Post-Cambrian Records of an Enigmatic Group of Brachiopoda. <i>Journal of Paleontology</i> , 2014, 88, 488-496.	0.5	6
39	Early Silurian (Rhuddanian) rugose corals and sponges from the Ak-Kerme Peninsula, Kazakhstan. <i>Palaontologische Zeitschrift</i> , 2014, 88, 279-295.	0.8	1
40	Late Ordovician and early Silurian brachiopods from the Zagros Ranges, Iran. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2014, 105, 159-187.	0.3	6
41	Dapingian-lower Darriwilian (Ordovician) stratigraphic gap in the Faraghan Mountains, Zagros Ranges, south-eastern Iran. <i>Bulletin of Geosciences</i> , 2014, , 679-706.	0.5	31
42	New and revised inarticulate Radiolaria and associated trilobites from the upper Darriwilian (Ordovician) Shundy Formation of Kazakhstan. <i>Journal of Paleontology</i> , 2013, 87, 1143-1159.	0.5	14
43	First report of craniid brachiopods in the Palaeozoic of Iran (<i>Pseudocrania</i> , Ordovician), and Early to Mid-Ordovician biogeography of the Craniida. <i>Palaeontology</i> , 2013, 56, 209-216.	1.0	6
44	Chapter 10 Biogeography of Ordovician linguliform and craniiform brachiopods. <i>Geological Society Memoir</i> , 2013, 38, 117-126.	0.9	18
45	Chapter 19 Global Cambrian trilobite palaeobiogeography assessed using parsimony analysis of endemism. <i>Geological Society Memoir</i> , 2013, 38, 273-296.	0.9	39
46	The radiation of early Silurian spiriferid brachiopods, with new taxa from the Llandovery of Iran. <i>Alcheringa</i> , 2013, 37, 558-564.	0.5	8
47	Long-period orbital climate forcing in the early Palaeozoic?. <i>Journal of the Geological Society</i> , 2013, 170, 707-710.	0.9	17
48	Silurian craniid brachiopods from Gotland. <i>Palaeontology</i> , 2013, 56, 1029-1044.	1.0	7
49	Morphology, ontogeny and affinities of the Hirnantian triplisid brachiopod <i>Schreptis undifera</i> from Baltoscandia. <i>Palaeontology</i> , 2013, 56, 961-970.	1.0	3
50	Cambrian (Furongian) rhynchonelliform brachiopods from the Eastern Alborz Mountains, Iran. <i>Bulletin of Geosciences</i> , 2013, , 525-538.	0.5	10
51	Earliest ontogeny of Early Palaeozoic Craniiformea: compelling evidence for lecithotrophy. <i>Lethaia</i> , 2012, 45, 566-573.	0.6	10
52	An obolellate brachiopod with soft-part preservation from the Early Cambrian Chengjiang fauna of China. <i>Journal of Paleontology</i> , 2011, 85, 460-463.	0.5	18
53	Stratigraphic evidence for the Hirnantian (latest Ordovician) glaciation in the Zagros Mountains, Iran. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2011, 307, 1-16.	1.0	61
54	New, early ostracods from the Ordovician (Tremadocian) of Iran: systematic, biogeographical and palaeoecological significance. <i>Alcheringa</i> , 2011, 35, 517-529.	0.5	17

#	ARTICLE	IF	CITATIONS
55	Ordovician radiolaria and conodonts from the peri-Gondwanan Karatau-Naryn microplate (Sarydzhas, Tajikistan). <i>Tectonophysics</i> , 2011, 511, 1-10.	0.7	14
56	Relic aragonite from Ordovician to Silurian brachiopods: Implications for the evolution of calcification. <i>Geology</i> , 2011, 39, 967-970.	2.0	43
57	Early Ordovician (Tremadocian) brachiopods from the Eastern Alborz Mountains, Iran. <i>Estonian Journal of Earth Sciences</i> , 2011, 60, 65.	0.4	30
58	Earliest ontogeny of Early Palaeozoic Craniiformea: implications for brachiopod phylogeny. <i>Lethaia</i> , 2010, 43, 323-333.	0.6	16
59	Late Ordovician trilobites from the Karagach Formation of the western Tarbagatai Range, Kazakhstan. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2010, 101, 161-187.	0.3	13
60	Late Silurian trilobites from the Nuratau and Turkestan ranges, Uzbekistan and Tajikistan. <i>Geobios</i> , 2009, 42, 715-737.	0.7	5
61	Early ontogeny and soft tissue preservation in siphonotretide brachiopods: New data from the Cambrian to Ordovician of Iran. <i>Gondwana Research</i> , 2009, 16, 151-161.	3.0	30
62	Autoradiographic study on the distribution of ²⁴¹ Am in the shell of the freshwater zebra mussel <i>Dreissena polymorpha</i> . <i>Mikrochimica Acta</i> , 2009, 167, 173-178.	2.5	10
63	THE ENIGMATIC EARLY CAMBRIAN <i>SALANYGOLINA</i> : A STEM GROUP OF RHYNCHONELLIFORM CHILEATE BRACHIOPODS?. <i>Palaeontology</i> , 2009, 52, 1-10.	1.0	45
64	BILLENGSELLIDE AND ORTHIDE BRACHIOPODS: NEW INSIGHTS INTO EARLIEST ORDOVICIAN EVOLUTION AND BIOGEOGRAPHY FROM NORTHERN IRAN. <i>Palaeontology</i> , 2009, 52, 35-52.	1.0	27
65	GENERIC HOMES FOR BRITISH SILURIAN LINGULOID BRACHIOPODS. <i>Palaeontology</i> , 2009, 52, 349-367.	1.0	7
66	Silicified Middle Cambrian trilobites from Kyrgyzstan. <i>Palaeontology</i> , 2009, 52, 1039-1056.	1.0	2
67	Gondwanan faunal signatures from Early Palaeozoic terranes of Kazakhstan and Central Asia: evidence and tectonic implications. <i>Geological Society Special Publication</i> , 2009, 325, 23-64.	0.8	40
68	Brachiopoda from the Soom Shale Lagerstätte (Upper Ordovician, South Africa). <i>Journal of Paleontology</i> , 2009, 83, 614-623.	0.5	23
69	First Report on the Occurrence of <i>Neseuretina</i> and <i>Ovalocephalus</i> Trilobites in the Middle Ordovician of Iran. <i>Acta Palaeontologica Polonica</i> , 2009, 54, 125-133.	0.4	21
70	The earliest ostracods: the geological evidence. <i>Senckenbergiana Lethaea</i> , 2008, 88, 11-21.	0.3	71
71	Ordovician (Caradoc) Gastropoda of the Katkoyeh Formation, Kerman Province, Iran. <i>Geobios</i> , 2008, 41, 605-624.	0.7	9
72	Short-term episodes of carbonate productivity in a Cambrian uplifted rift shoulder of the Coastal Meseta, Morocco. <i>Gondwana Research</i> , 2008, 14, 410-428.	3.0	28

#	ARTICLE	IF	CITATIONS
73	Early to Middle Ordovician lingulate brachiopods from the Lashkarak Formation, Eastern Alborz Mountains, Iran. <i>Alcheringa</i> , 2008, 32, 1-35.	0.5	29
74	A New Genus of Late Ordovician–Early Silurian Pentameride Brachiopods and Its Phylogenetic Relationships. <i>Acta Palaeontologica Polonica</i> , 2008, 53, 221-236.	0.4	8
75	Columnar shell structures in early linguloid brachiopods – new data from the Middle Cambrian of Sweden. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2007, 98, 221-232.	0.3	16
76	Pedicle preservation in a Silurian rhynchonelliformean brachiopod from Herefordshire, England: soft-tissue or an artefact of interpretation?. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2007, 98, 303-308.	0.3	8
77	Biogeography and affinities of the bradoriid arthropods: Cosmopolitan microbenthos of the Cambrian seas. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2007, 248, 202-232.	1.0	60
78	A new Middle Ordovician arthropod fauna (Trilobita, Ostracoda, Bradoriida) from the Lashkarak Formation, Eastern Alborz Mountains, northern Iran. <i>Gff</i> , 2007, 129, 245-254.	0.4	18
79	Earliest ontogeny of Middle Ordovician rhynchonelliform brachiopods (<i>Clitambonitoidea</i> and) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10</i>	0.8	37
80	LATE ORDOVICIAN BRACHIOPODS FROM THE DULANKARA FORMATION OF THE CHU-ILI RANGE, KAZAKHSTAN: THEIR SYSTEMATICS, PALAEOECOLOGY AND PALAEOBIOGEOGRAPHY. <i>Palaeontology</i> , 2006, 49, 247-283.	1.0	23
81	Proposed stratotype for the base of the highest Cambrian stage at the first appearance datum of <i>Cordylodus andresi</i> , Lawson Cove section, Utah, USA. <i>Palaeoworld</i> , 2006, 15, 384-405.	0.5	31
82	ORDOVICIAN (ARENIG-CARADOC) SYNTROPHIIDINE BRACHIOPODS FROM THE EAST BALTIC REGION. <i>Palaeontology</i> , 2005, 48, 739-761.	1.0	14
83	TWO ENIGMATIC BRYOZOANS FROM THE MIDDLE ORDOVICIAN OF THE EAST BALTIC. <i>Palaeontology</i> , 2005, 48, 1065-1074.	1.0	6
84	Neodymium isotopic composition of Cambrian–Ordovician biogenic apatite in the Baltoscandian Basin: implications for palaeogeographical evolution and patterns of biodiversity. <i>Geological Magazine</i> , 2005, 142, 419-439.	0.9	40
85	LOWER ORDOVICIAN (TREMADOCIAN) LINGULATE BRACHIOPODS FROM THE HOUSE AND FILLMORE FORMATIONS, IBEX AREA, WESTERN UTAH, USA. <i>Journal of Paleontology</i> , 2005, 79, 884-906.	0.5	33
86	The oldest-known metazoan parasite?. <i>Journal of Paleontology</i> , 2004, 78, 1214-1216.	0.5	15
87	Conodont biostratigraphy and faunal assemblages in radiolarian ribbon-banded cherts of the Burubaital Formation, West Balkhash Region, Kazakhstan. <i>Geological Magazine</i> , 2004, 141, 699-715.	0.9	34
88	Late Cambrian relative age constraints by acritarchs on the post-Timanian deposition on Kolguev Island, Arctic Russia. <i>Geological Society Memoir</i> , 2004, 30, 159-168.	0.9	0
89	Upper Cambrian-Ordovician successions overlying Timanian complexes: new evidence of acritarchs and brachiopods from Kolguev Island, Arctic Russia. <i>Geobios</i> , 2004, 37, 239-251.	0.7	3
90	THE OLDEST-KNOWN METAZOAN PARASITE?. <i>Journal of Paleontology</i> , 2004, 78, 1214-1216.	0.5	27

#	ARTICLE	IF	CITATIONS
91	A new Ordovician rostroconch mollusc from Kazakhstan. <i>Alcheringa</i> , 2003, 27, 173-179.	0.5	6
92	Ordovician radiolarian biodiversity : insights based on new and revised data from Kazakhstan. <i>Bulletin - Societe Geologique De France</i> , 2003, 174, 325-335.	0.9	33
93	Understanding linguloid brachiopods: <i>Obolus</i> and <i>Ungula</i> as examples. <i>Carnets De Geologie</i> , 2003, , .	0.4	3
94	LINGULATE BRACHIOPODS FROM THE CAMBRIAN-ORDOVICIAN BOUNDARY BEDS OF UTAH. <i>Journal of Paleontology</i> , 2002, 76, 211-228.	0.5	17
95	Brachiopods: Cambrian-Tremadoc precursors to Ordovician radiation events. <i>Geological Society Special Publication</i> , 2002, 194, 13-23.	0.8	37
96	Lingulate brachiopods from the Cambrian-Ordovician boundary beds of Utah. <i>Journal of Paleontology</i> , 2002, 76, 211-228.	0.5	24
97	The Hunneberg Stage (Ordovician) in the area east of St. Petersburg, north-western Russia. <i>Palaontologische Zeitschrift</i> , 2001, 74, 543-561.	0.8	28
98	Brachiopods of the redefined family <i>Tritoechiidae</i> from the Ordovician of Kazakhstan and South Urals. <i>Geobios</i> , 2001, 34, 131-155.	0.7	31
99	Evidence for 15 m.y. of continuous deep-sea biogenic siliceous sedimentation in early Paleozoic oceans. <i>Geology</i> , 2001, 29, 755.	2.0	52
100	Late Ordovician Brachiopods From The Otar Member Of The Chu-Ili Range, South Kazakhstan. <i>Palaeontology</i> , 2000, 43, 833-870.	1.0	24
101	Late Ordovician (Ashgill)linguliformean microbrachiopods from the Bestyube Formation, north-central Kazakhstan. <i>Geobios</i> , 2000, 33, 419-435.	0.7	9
102	Relative sea-level changes in Baltoscandia in the Cambrian and early Ordovician: the predominance of tectonic factors and the absence of large scale eustatic fluctuations. <i>Tectonophysics</i> , 2000, 320, 375-407.	0.9	48
103	Early Ordovician organophosphatic brachiopods with Baltoscandian affinities from the Alay Range, southern Kyrgyzstan. <i>Gff</i> , 2000, 122, 367-375.	0.4	13
104	Late Ordovician linguliformean microbrachiopods from north-central Kazakhstan. <i>Alcheringa</i> , 2000, 24, 257-275.	0.5	12
105	Early Ordovician bryozoans from north-western Russia. <i>Palaeontology</i> , 1999, 42, 171-189.	1.0	21
106	The earliest atrypides and athyridides (Brachiopoda) from the Ordovician of Kazakhstan. <i>Palaeontology</i> , 1999, 42, 625-661.	1.0	43
107	Organophosphatic brachiopods:Patterns of biodiversification and extinction in the Early Palaeozoic. <i>Geobios</i> , 1999, 32, 145-163.	0.7	54
108	Cambrian phosphatic brachiopods from the Precordillera of western Argentina. <i>Gff</i> , 1999, 121, 227-242.	0.4	13

#	ARTICLE	IF	CITATIONS
109	Nd isotope composition and rare earth element distribution in early Paleozoic biogenic apatite from Baltoscandia: A signature of Iapetus ocean water. <i>Geology</i> , 1998, 26, 1083.	2.0	38
110	Early Cambrian Lingulellotreta (Lingulata, Brachiopoda) from South Kazakhstan (Malyi Karatau Range) and South China (Eastern Yunnan). <i>Journal of Paleontology</i> , 1997, 71, 577-584.	0.5	35
111	Late Ordovician and early Silurian Trimerellide brachiopods from Kazakhstan. <i>Journal of Paleontology</i> , 1997, 71, 584-598.	0.5	25
112	Strophomenid and triplesiid brachiopods from an Upper Ordovician carbonate mound in central Kazakhstan. <i>Alcheringa</i> , 1996, 20, 1-20.	0.5	18
113	Middle Ordovician (Llanvirn) unguulate brachiopods and conodonts from the Malyi Karatau Range, Kazakhstan. <i>Palaontologische Zeitschrift</i> , 1996, 70, 481-495.	0.8	6
114	Late Ordovician brachiopod assemblage of Hiberno-Salairian type from Central Kazakhstan. <i>Gff</i> , 1996, 118, 83-96.	0.4	17
115	The elkaniid brachiopod <i>Volborthia</i> from the lower Ordovician of Baltoscandia. <i>Palaontologische Zeitschrift</i> , 1995, 69, 213-221.	0.8	3
116	Review of the Cambrian acrotretid brachiopod <i>Neotreta</i> . <i>Alcheringa</i> , 1994, 18, 345-357.	0.5	7
117	First record of upper Cambrian from the eastern White Sea coast: New evidence from obolids (Brachiopoda). <i>Gff</i> , 1994, 116, 31-35.	0.4	10
118	Cambrian-Ordovician lingulate brachiopods from Scandinavia, Kazakhstan, and South Ural Mountains. <i>Lethaia</i> , 1994, 27, 1-156.	0.6	13
119	Revision of the type species of <i>Acrotreta</i> and related lingulate brachiopods. <i>Journal of Paleontology</i> , 1994, 68, 433-450.	0.5	18
120	First occurrence of the genus <i>Bicarinata</i> (Brachiopoda, Inarticulata) from the Middle Devonian in North America (Alaska). <i>Journal of Paleontology</i> , 1994, 68, 1214-1218.	0.5	11
121	Phylogenetic analysis of higher taxa of Brachiopoda. <i>Lethaia</i> , 1993, 26, 1-5.	0.6	75
122	Phylogenetic analysis and classification of the Brachiopoda - reply and comments. <i>Lethaia</i> , 1993, 26, 385-386.	0.6	8
123	The acrotretacean brachiopod <i>Ceratreta tanneri</i> (Metzger) from the Upper Cambrian of Baltoscandia. <i>Gff</i> , 1990, 112, 249-263.	0.4	13
124	On the origin systematic position of the calcareous-shelled inarticulate brachiopods. <i>Lethaia</i> , 1986, 19, 233-240.	0.6	47
125	Lower Paleozoic radiolarians of Kazakhstan. <i>International Geology Review</i> , 1977, 19, 913-920.	1.1	8
126	Were transgressive black shales a negative feedback modulating glacioeustasy in the Early Palaeozoic Icehouse? , 0, , 123-156.		49

#	ARTICLE	IF	CITATIONS
127	Cambrian (Stage 4 to Wuliuan) brachiopods from Sonora, Mexico. Journal of Paleontology, 0, , 1-21.	0.5	2