## **Christophe Fuerer**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8003174/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Wnt Signaling Mediates Self-Organization and Axis Formation in Embryoid Bodies. Cell Stem Cell, 2008, 3, 508-518.	5.2	406
2	Lentiviral Vectors to Probe and Manipulate the Wnt Signaling Pathway. PLoS ONE, 2010, 5, e9370.	1.1	241
3	Wnt Proteins Promote Bone Regeneration. Science Translational Medicine, 2010, 2, 29ra30.	5.8	235
4	Wnt Signaling and Stem Cell Control. Cold Spring Harbor Symposia on Quantitative Biology, 2008, 73, 59-66.	2.0	203
5	Wnt signaling mediates regional specification in the vertebrate face. Development (Cambridge), 2007, 134, 3283-3295.	1.2	188
6	Wnt5a can both activate and repress Wnt/β-catenin signaling during mouse embryonic development. Developmental Biology, 2012, 369, 101-114.	0.9	185
7	Secreted Wingless-interacting molecule (Swim) promotes long-range signaling by maintaining Wingless solubility. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 370-377.	3.3	157
8	Wnt/β-Catenin Signaling in Murine Hepatic Transit Amplifying Progenitor Cells. Gastroenterology, 2007, 133, 1579-1591.e1.	0.6	154
9	A study on the interactions between heparan sulfate proteoglycans and Wnt proteins. Developmental Dynamics, 2010, 239, 184-190.	0.8	93
10	The microenvironment patterns the pluripotent mouse epiblast through paracrine Furin and Pace4 proteolytic activities. Genes and Development, 2011, 25, 1871-1880.	2.7	42
11	AOAC SMPR® 2016.002. Journal of AOAC INTERNATIONAL, 2016, 99, 1122-1124.	0.7	39
12	Late Expression of Nitroreductase in an Oncolytic Adenovirus Sensitizes Colon Cancer Cells to the Prodrug CB1954. Human Gene Therapy, 2005, 16, 1473-1483.	1.4	38
13	Nodal·Gdf1 Heterodimers with Bound Prodomains Enable Serum-independent Nodal Signaling and Endoderm Differentiation. Journal of Biological Chemistry, 2014, 289, 17854-17871.	1.6	36
14	Amyloidâ€beta oligomerization is associated with the generation of a typical peptide fragment fingerprint. Alzheimer's and Dementia, 2016, 12, 996-1013.	0.4	17
15	Total Amino Acids by UHPLC–UV in Infant Formulas and Adult Nutritionals, First Action 2018.06. Journal of AOAC INTERNATIONAL, 2019, 102, 1574-1588.	0.7	15
16	Quantification of Whey Protein Content in Infant Formulas by Sodium Dodecyl Sulfate-Capillary Gel Electrophoresis (SDS-CGE): Single-Laboratory Validation, First Action 2016.15. Journal of AOAC INTERNATIONAL, 2017, 100, 510-521.	0.7	11
17	Food Fraud Vulnerabilities in the Supply Chain: An Industry Perspective. , 2019, , 670-678.		9
18	Quantification of Whey Protein Content in Milk-Based Infant Formula Powders by Sodium Dodecyl Sulfate–Capillary Gel Electrophoresis (SDS-CGE): Multilaboratory Testing Study, Final Action 2016.15. Journal of AOAC INTERNATIONAL, 2018, 101, 1566-1577.	0.7	7

#	Article	IF	CITATIONS
19	Fusion of the BCL9 HD2 domain to E1A increases the cytopathic effect of an oncolytic adenovirus that targets colon cancer cells. BMC Cancer, 2006, 6, 236.	1.1	3
20	Quantification of Whey Protein Content in Infant Formulas by Sodium Dodecyl Sulfate-Capillary Gel Electrophoresis (SDS-CGE): Single-Laboratory Validation, First Action 2016.15. Journal of AOAC INTERNATIONAL, 2017, 100, 1177-1180.	0.7	2
21	Determination of Total Amino Acids in Infant Formulas, Adult Nutritionals, Dairy, and Cereal Matrixes by UHPLC–UV: Interlaboratory Validation Study, Final Action 2018.06. Journal of AOAC INTERNATIONAL, 2022, 105, 1625-1639.	0.7	2