

Ana J Gonzalez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8011305/publications.pdf>

Version: 2024-02-01

28
papers

401
citations

759233

12
h-index

794594

19
g-index

28
all docs

28
docs citations

28
times ranked

404
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Clavibacter michiganensis</i> subsp. <i>phaseoli</i> subsp. nov., pathogenic in bean. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1752-1755.	1.7	39
2	Identification of an Emergent and Atypical <i>Pseudomonas viridiflava</i> Lineage Causing Bacteriosis in Plants of Agronomic Importance in a Spanish Region. Applied and Environmental Microbiology, 2003, 69, 2936-2941.	3.1	36
3	Pathovars of <i>Pseudomonas syringae</i> Causing Bacterial Brown Spot and Halo Blight in <i>Phaseolus vulgaris</i> L. Are Distinguishable by Ribotyping. Applied and Environmental Microbiology, 2000, 66, 850-854.	3.1	30
4	<i>Pseudomonas asturiensis</i> sp. nov., isolated from soybean and weeds. Systematic and Applied Microbiology, 2013, 36, 320-324.	2.8	25
5	<i>Erwinia persicina</i> Causing Chlorosis and Necrotic Spots in Leaves and Tendrils of <i>Pisum sativum</i> in Southeastern Spain. Plant Disease, 2007, 91, 460-460.	1.4	25
6	Plasmid typing of <i>Shigella sonnei</i> epidemic strains and molecular relationship of their R-plasmids. European Journal of Epidemiology, 1988, 4, 158-163.	5.7	24
7	Clonal population structure and introductions of the chestnut blight fungus, <i>Cryphonectria parasitica</i> , in Asturias, northern Spain. European Journal of Plant Pathology, 2011, 131, 67-79.	1.7	20
8	Identification, incidence and control of bacterial blotch disease in mushroom crops by management of environmental conditions. Scientia Horticulturae, 2018, 229, 10-18.	3.6	20
9	Bacterial Wilt of Beans (<i>Phaseolus vulgaris</i>) Caused by <i>Curtobacterium flaccumfaciens</i> in Southeastern Spain. Plant Disease, 2005, 89, 1361-1361.	1.4	20
10	First Report of <i>Erwinia persicina</i> from <i>Phaseolus vulgaris</i> in Spain. Plant Disease, 2005, 89, 109-109.	1.4	17
11	Brown Blotch Caused by <i>Pseudomonas tolaasii</i> on Cultivated <i>Pleurotus eryngii</i> in Spain. Plant Disease, 2009, 93, 667-667.	1.4	17
12	<i>Pseudomonas syringae</i> pv. <i>phaseolicola</i> isolated from weeds in bean crop fields. Letters in Applied Microbiology, 2016, 62, 344-348.	2.2	16
13	First Report of Cobweb on White Button Mushroom (<i>Agaricus bisporus</i>) in Spain Caused by <i>Cladobotryum mycophilum</i> . Plant Disease, 2012, 96, 1067-1067.	1.4	14
14	Phylogenetic and phenotypic characterisation of <i>Sirococcus castaneae</i> comb. nov. (synonym <i>Diplodina</i>) Tj ETQq0 0,0,rgBT /Overlock 10	2.5	13
15	<i>Brenneria goodwinii</i> and <i>Gibbsiella quercinecans</i> isolated from weeping cankers on <i>Quercus robur</i> L. in Spain. European Journal of Plant Pathology, 2020, 156, 965-969.	1.7	13
16	Identification and RAPD-typing of <i>Ewingella americana</i> on cultivated mushrooms in Castilla-La Mancha, Spain. European Journal of Plant Pathology, 2012, 133, 517-522.	1.7	11
17	Occurrence of hypovirus-infected <i>Cryphonectria parasitica</i> isolates in northern Spain: an encouraging situation for biological control of chestnut blight in Asturian forests. European Journal of Plant Pathology, 2017, 149, 503-514.	1.7	10
18	First Report of <i>Gnomoniopsis castaneae</i> Causing Canker on Hybrid Plants of <i>Castanea sativa</i> – <i>C. crenata</i> in Spain. Plant Disease, 2018, 102, 1040-1040.	1.4	9

#	ARTICLE	IF	CITATIONS
19	Regional variations in the population structure of <i>Pseudomonas syringae</i> pathovar <i>phaseolicola</i> from Spain are revealed by typing with Pmel pulsed-field gel electrophoresis, plasmid profiling and virulence gene complement. <i>Microbiology (United Kingdom)</i> , 2010, 156, 1795-1804.	1.8	7
20	A <i>Pseudomonas viridiflava</i> -Related Bacterium Causes a Dark-Reddish Spot Disease in <i>Glycine max.</i> <i>Applied and Environmental Microbiology</i> , 2012, 78, 3756-3758.	3.1	6
21	Diversity of culturable bacteria and occurrence of phytopathogenic species in bean seeds (<i>Phaseolus</i>) Tj ETQq1 1 0,784314 rgBT /Ove	1.6	8
22	Disease of Floral Buds of Kiwifruit in Spain Caused by <i>Pseudomonas syringae</i> . <i>Plant Disease</i> , 2001, 85, 1287-1287.	1.4	5
23	Short communication: Occurrence of angular leaf spot caused by <i>Pseudocercospora griseola</i> in <i>Phaseolus vulgaris</i> in Asturias, Spain. <i>Spanish Journal of Agricultural Research</i> , 2017, 15, e10SC03.	0.6	5
24	<i>Pseudomonas viridiflava</i> Causing Necrotic Leaf Spots and Defoliation on <i>Hebe</i> spp. in Northern Spain. <i>Plant Disease</i> , 2006, 90, 830-830.	1.4	4
25	Chestnut Blight Control by Agrochemicals in <i>Castanea sativa</i> under Managed Conditions. <i>Journal of Plant Diseases and Protection</i> , 2015, 122, 120-124.	2.9	3
26	Strains of <i>Neopestalotiopsis</i> sp. are in vitro antagonists of <i>Cryphonectria parasitica</i> . <i>Biological Control</i> , 2020, 143, 104187.	3.0	3
27	First Report of <i>Neofusicoccum parvum</i> Causing Canker on <i>Castanea sativa</i> in Spain. <i>Plant Disease</i> , 2022, 106, 1299.	1.4	2
28	Phenotypic and genotypic analysis of <i>Pseudomonas syringae</i> recovered from symptomatic beans and associated weeds in Northern Spain. <i>European Journal of Plant Pathology</i> , 2020, 157, 377-387.	1.7	1