
Claire L Nollett

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/802303/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Training results in increased practitioner confidence and identification of depression in people with low vision: a mixed methods study. Ophthalmic and Physiological Optics, 2021, 41, 316-330.	2.0	6
2	Normalisation process theory and the implementation of a new glaucoma clinical pathway in hospital eye services: Perspectives of doctors, nurses and optometrists. PLoS ONE, 2021, 16, e0255564.	2.5	0
3	Barriers to integrating routine depression screening into community low vision rehabilitation services: a mixed methods study. BMC Psychiatry, 2020, 20, 419.	2.6	9
4	<p>Chronic Conditions and Behavioural Change Approaches to Medication Adherence: Rethinking Clinical Guidance and Recommendations</p> . Patient Preference and Adherence, 2020, Volume 14, 581-586.	1.8	4
5	Acceptability, adherence and economic analyses of a new clinical pathway for the identification of non-responders to glaucoma eye drops: a prospective observational study. British Journal of Ophthalmology, 2020, 104, 1704-1709.	3.9	1
6	Depressive symptoms in people with vision impairment: a cross-sectional study to identify who is most at risk. BMJ Open, 2019, 9, e026163.	1.9	27
7	How do community-based eye care practitioners approach depression in patients with low vision? A mixed methods study. BMC Psychiatry, 2019, 19, 426.	2.6	12
8	Feasibility of weekly participant-reported data collection in a pragmatic randomised controlled trial in primary care: experiences from the BATHE trial (Bath Additives for the Treatment of cHildhood) Tj ETQq0 0 C) rgB T./ Over	loc k 10 Tf 50
9	Emollient bath additives for the treatment of childhood eczema (BATHE): multicentre pragmatic parallel group randomised controlled trial of clinical and cost effectiveness. BMJ: British Medical Journal, 2018, 361, k1332.	2.3	50
10	Pragmatic RAndomised controlled trial of a trauma-focused guided self-help Programme versus InDividual trauma-focused cognitive Behavioural therapy for post-traumatic stress disorder (RAPID): trial protocol. BMC Psychiatry, 2018, 18, 77.	2.6	10
11	Adding emollient bath additives to standard eczema management for children with eczema: the BATHE RCT. Health Technology Assessment, 2018, 22, 1-116.	2.8	14
12	Depression in Visual Impairment Trial (DEPVIT): A Randomized Clinical Trial of Depression Treatments in People With Low Vision. , 2016, 57, 4247.		36
13	Site visits to initiate recruitment in a clinical trial: does it matter who conducts the visit? Protocol for implementation in trials. Journal of Evidence-Based Medicine, 2016, 9, 225-227.	1.8	0
14	High Prevalence of Untreated Depression in Patients Accessing Low-Vision Services. Ophthalmology, 2016, 123, 440-441.	5.2	45
15	The effectiveness of bariatric surgery on long term psychosocial quality of life – A systematic review. Obesity Research and Clinical Practice, 2016, 10, 225-242.	1.8	61
16	The Depression in Visual Impairment Trial (DEPVIT): trial design and protocol. BMC Psychiatry, 2012, 12, 57.	2.6	35
17	Don't forget EDNOS (eating disorder not otherwise specified): patterns of service use in an eating disorders service. Psychiatric Bulletin, 2005, 29, 134-136.	0.3	25

18Questionnaire measures of psychopathology in eating disorders: comparisons between clinical
groups. European Eating Disorders Review, 2005, 13, 211-215.4.113

#	Article	IF	CITATIONS
19	Guided, internet based, cognitive behavioural therapy for post-traumatic stress disorder: pragmatic, multicentre, randomised controlled non-inferiority trial (RAPID). BMJ, The, 0, , e069405.	6.0	10