## **Insung Kang**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8039257/publications.pdf

Version: 2024-02-01

567281 610901 25 691 15 24 citations h-index g-index papers 25 25 25 1173 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Upregulation of SNAP25 by HDAC inhibition ameliorates Niemannâ€Pick Type C disease phenotypes via autophagy induction. Clinical and Translational Medicine, 2022, 12, e776.	4.0	O
2	Oral administration of microbiome-friendly graphene quantum dots as therapy for colitis. 2D Materials, 2021, 8, 025036.	4.4	7
3	Graphene Quantum Dots from Carbonized Coffee Bean Wastes for Biomedical Applications. Nanomaterials, 2021, 11, 1423.	4.1	27
4	Induced neural stem cells from human patient-derived fibroblasts attenuate neurodegeneration in Niemann-Pick type C mice. Journal of Veterinary Science, 2021, 22, e7.	1.3	2
5	Graphene Quantum Dots Alleviate Impaired Functions in Niemann-Pick Disease Type C in Vivo. Nano Letters, 2021, 21, 2339-2346.	9.1	17
6	cAMP/EPAC Signaling Enables ETV2 to Induce Endothelial Cells with High Angiogenesis Potential. Molecular Therapy, 2020, 28, 466-478.	8.2	13
7	Graphene quantum dots as anti-inflammatory therapy for colitis. Science Advances, 2020, 6, eaaz2630.	10.3	88
8	Interferon- $\hat{i}^3$ -mediated secretion of tryptophanyl-tRNA synthetases has a role in protection of human umbilical cord blood-derived mesenchymal stem cells against experimental colitis. BMB Reports, 2019, 52, 318-323.	2.4	11
9	Disease-specific primed human adult stem cells effectively ameliorate experimental atopic dermatitis in mice. Theranostics, 2019, 9, 3608-3621.	10.0	26
10	Human umbilical cord blood plasma alleviates age-related olfactory dysfunction by attenuating peripheral TNF-α expression. BMB Reports, 2019, 52, 259-264.	2.4	5
11	Donor-dependent variation of human umbilical cord blood mesenchymal stem cells in response to hypoxic preconditioning and amelioration of limb ischemia. Experimental and Molecular Medicine, 2018, 50, 1-15.	7.7	56
12	Stem cell-secreted 14,15- epoxyeicosatrienoic acid rescues cholesterol homeostasis and autophagic flux in Niemann–Pick-type C disease. Experimental and Molecular Medicine, 2018, 50, 1-14.	7.7	13
13	GATA4-dependent regulation of the secretory phenotype via MCP-1 underlies lamin A-mediated human mesenchymal stem cell aging. Experimental and Molecular Medicine, 2018, 50, 1-12.	7.7	24
14	MIS416 Enhances Therapeutic Functions of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Against Experimental Colitis by Modulating Systemic Immune Milieu. Frontiers in Immunology, 2018, 9, 1078.	4.8	18
15	Single-Factor SOX2 Mediates Direct Neural Reprogramming of Human Mesenchymal Stem Cells via Transfection of <i>In Vitro</i> Transcribed mRNA. Cell Transplantation, 2018, 27, 1154-1167.	2.5	23
16	Cell Surface Nanoâ€modulation for Nonâ€invasive inâ€vivo Nearâ€iR Stem Cell Monitoring. ChemMedChem, 2017, 12, 28-32.	3.2	2
17	Human adipose tissue-derived mesenchymal stem cells alleviate atopic dermatitis via regulation of B lymphocyte maturation. Oncotarget, 2017, 8, 512-522.	1.8	61
18	Direct Conversion of Human Umbilical Cord Blood into Induced Neural Stem Cells with SOX2 and HMGA2. International Journal of Stem Cells, 2017, 10, 227-234.	1.8	13

#	Article	IF	CITATION
19	Generation of patient specific human neural stem cells from Niemann-Pick disease type C patient-derived fibroblasts. Oncotarget, 2017, 8, 85428-85441.	1.8	22
20	PGE2 maintains self-renewal of human adult stem cells via EP2-mediated autocrine signaling and its production is regulated by cell-to-cell contact. Scientific Reports, 2016, 6, 26298.	3.3	69
21	Cathepsin S contributes to microglia-mediated olfactory dysfunction through the regulation of Cx3cl1-Cx3cr1 axis in a Niemann-Pick disease type C1 model. Glia, 2016, 64, 2291-2305.	4.9	36
22	Inducible HGF-secreting Human Umbilical Cord Blood-derived MSCs Produced via TALEN-mediated Genome Editing Promoted Angiogenesis. Molecular Therapy, 2016, 24, 1644-1654.	8.2	45
23	BMI1 inhibits senescence and enhances the immunomodulatory properties of human mesenchymal stem cells via the direct suppression of MKP-1/DUSP1. Aging, 2016, 8, 1670-1689.	3.1	24
24	A p38 MAPK-Mediated Alteration of COX-2/PGE2 Regulates Immunomodulatory Properties in Human Mesenchymal Stem Cell Aging. PLoS ONE, 2014, 9, e102426.	2.5	58
25	Excessive microglial activation aggravates olfactory dysfunction by impeding the survival of newborn neurons in the olfactory bulb of Niemann–Pick disease type C1 mice. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2014, 1842, 2193-2203.	3.8	31