

Michele A Kelly

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8042358/publications.pdf>

Version: 2024-02-01

15
papers

2,447
citations

567281

15
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

2221
citing authors

#	ARTICLE	IF	CITATIONS
1	Exocrine Gland Dysfunction in MC5-R-Deficient Mice: Evidence for Coordinated Regulation of Exocrine Gland Function by Melanocortin Peptides. <i>Cell</i> , 1997, 91, 789-798.	28.9	466
2	Pituitary Lactotroph Hyperplasia and Chronic Hyperprolactinemia in Dopamine D2 Receptor-Deficient Mice. <i>Neuron</i> , 1997, 19, 103-113.	8.1	398
3	Locomotor Activity in D2 Dopamine Receptor-Deficient Mice Is Determined by Gene Dosage, Genetic Background, and Developmental Adaptations. <i>Journal of Neuroscience</i> , 1998, 18, 3470-3479.	3.6	395
4	Alcohol preference and sensitivity are markedly reduced in mice lacking dopamine D2 receptors. <i>Nature Neuroscience</i> , 1998, 1, 610-615.	14.8	236
5	Dopamine D ₂ Receptor-Deficient Mice Exhibit Decreased Dopamine Transporter Function but No Changes in Dopamine Release in Dorsal Striatum. <i>Journal of Neurochemistry</i> , 1999, 72, 148-156.	3.9	206
6	The Dopamine D2, but not D3 or D4, Receptor Subtype is Essential for the Disruption of Prepulse Inhibition Produced by Amphetamine in Mice. <i>Journal of Neuroscience</i> , 1999, 19, 4627-4633.	3.6	169
7	Pituitary Lactotroph Adenomas Develop after Prolonged Lactotroph Hyperplasia in Dopamine D2 Receptor-Deficient Mice. <i>Endocrinology</i> , 1999, 140, 5348-5355.	2.8	159
8	Lack of prolactin receptor signaling in mice results in lactotroph proliferation and prolactinomas by dopamine-dependent and -independent mechanisms. <i>Journal of Clinical Investigation</i> , 2002, 110, 973-981.	8.2	95
9	Functional Uncoupling of Adenosine A _{2A} Receptors and Reduced Response to Caffeine in Mice Lacking Dopamine D ₂ Receptors. <i>Journal of Neuroscience</i> , 2000, 20, 5949-5957.	3.6	89
10	Lack of prolactin receptor signaling in mice results in lactotroph proliferation and prolactinomas by dopamine-dependent and -independent mechanisms. <i>Journal of Clinical Investigation</i> , 2002, 110, 973-981.	8.2	56
11	Neuropeptide B-deficient mice demonstrate hyperalgesia in response to inflammatory pain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 9942-9947.	7.1	55
12	Pituitary Lactotroph Adenomas Develop after Prolonged Lactotroph Hyperplasia in Dopamine D2 Receptor-Deficient Mice. <i>Endocrinology</i> , 1999, 140, 5348-5355.	2.8	45
13	The mapping of quantitative trait loci underlying strain differences in locomotor activity between 129S6 and C57BL/6J mice. <i>Mammalian Genome</i> , 2003, 14, 692-702.	2.2	36
14	Cloning of the mouse gonadotropin β -subunit-encoding genes, I. Structure of the follicle-stimulating hormone β -subunit-encoding gene. <i>Gene</i> , 1995, 166, 333-334.	2.2	24
15	Selective increase of Nurr1 mRNA expression in mesencephalic dopaminergic neurons of D2 dopamine receptor-deficient mice. <i>Molecular Brain Research</i> , 2000, 80, 1-6.	2.3	18