

Kazuo Eguchi

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8062938/kazuo-eguchi-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

202
papers

5,941
citations

40
h-index

69
g-index

206
ext. papers

6,711
ext. citations

3.2
avg, IF

5.34
L-index

#	Paper	IF	Citations
202	Prognostic Effect of the Nocturnal Blood Pressure Fall in Hypertensive Patients: The Ambulatory Blood Pressure Collaboration in Patients With Hypertension (ABC-H) Meta-Analysis. <i>Hypertension</i> , 2016 , 67, 693-700	8.5	282
201	U-curve relationship between orthostatic blood pressure change and silent cerebrovascular disease in elderly hypertensives: orthostatic hypertension as a new cardiovascular risk factor. <i>Journal of the American College of Cardiology</i> , 2002 , 40, 133-41	15.1	187
200	Prognostic impact from clinic, daytime, and night-time systolic blood pressure in nine cohorts of 13,844 patients with hypertension. <i>Journal of Hypertension</i> , 2014 , 32, 2332-40; discussion 2340	1.9	175
199	Masked hypertension: a review. <i>Hypertension Research</i> , 2007 , 30, 479-88	4.7	162
198	Ambulatory blood pressure is a better marker than clinic blood pressure in predicting cardiovascular events in patients with/without type 2 diabetes. <i>American Journal of Hypertension</i> , 2008 , 21, 443-50	2.3	155
197	Differential effects between a calcium channel blocker and a diuretic when used in combination with angiotensin II receptor blocker on central aortic pressure in hypertensive patients. <i>Hypertension</i> , 2009 , 54, 716-23	8.5	154
196	Morning hypertension: the strongest independent risk factor for stroke in elderly hypertensive patients. <i>Hypertension Research</i> , 2006 , 29, 581-7	4.7	145
195	Night time blood pressure variability is a strong predictor for cardiovascular events in patients with type 2 diabetes. <i>American Journal of Hypertension</i> , 2009 , 22, 46-51	2.3	128
194	Reproducibility of arterial stiffness indices (pulse wave velocity and augmentation index) simultaneously assessed by automated pulse wave analysis and their associated risk factors in essential hypertensive patients. <i>Hypertension Research</i> , 2004 , 27, 851-7	4.7	128
193	Morning blood pressure surge and hypertensive cerebrovascular disease: role of the alpha adrenergic sympathetic nervous system. <i>American Journal of Hypertension</i> , 2004 , 17, 668-75	2.3	127
192	Association between diabetes mellitus and left ventricular hypertrophy in a multiethnic population. <i>American Journal of Cardiology</i> , 2008 , 101, 1787-91	3	124
191	Added predictive value of night-time blood pressure variability for cardiovascular events and mortality: the Ambulatory Blood Pressure-International Study. <i>Hypertension</i> , 2014 , 64, 487-93	8.5	121
190	Morning and Evening Home Blood Pressure and Risks of Incident Stroke and Coronary Artery Disease in the Japanese General Practice Population: The Japan Morning Surge-Home Blood Pressure Study. <i>Hypertension</i> , 2016 , 68, 54-61	8.5	117
189	Visit-to-visit and ambulatory blood pressure variability as predictors of incident cardiovascular events in patients with hypertension. <i>American Journal of Hypertension</i> , 2012 , 25, 962-8	2.3	105
188	Maximum value of home blood pressure: a novel indicator of target organ damage in hypertension. <i>Hypertension</i> , 2011 , 57, 1087-93	8.5	98
187	Relationship between extreme dippers and orthostatic hypertension in elderly hypertensive patients. <i>Hypertension</i> , 1998 , 31, 77-82	8.5	97
186	Nighttime home blood pressure and the risk of hypertensive target organ damage. <i>Hypertension</i> , 2012 , 60, 921-8	8.5	96

185	Short sleep duration as an independent predictor of cardiovascular events in Japanese patients with hypertension. <i>Archives of Internal Medicine</i> , 2008 , 168, 2225-31		93
184	Effect of dosing time of angiotensin II receptor blockade titrated by self-measured blood pressure recordings on cardiorenal protection in hypertensives: the Japan Morning Surge-Target Organ Protection (J-TOP) study. <i>Journal of Hypertension</i> , 2010 , 28, 1574-83	1.9	90
183	The misdiagnosis of hypertension: the role of patient anxiety. <i>Archives of Internal Medicine</i> , 2008 , 168, 2459-65		84
182	Consistency of blood pressure differences between the left and right arms. <i>Archives of Internal Medicine</i> , 2007 , 167, 388-93		78
181	Greater impact of coexistence of hypertension and diabetes on silent cerebral infarcts. <i>Stroke</i> , 2003 , 34, 2471-4	6.7	78
180	Greater change of orthostatic blood pressure is related to silent cerebral infarct and cardiac overload in hypertensive subjects. <i>Hypertension Research</i> , 2004 , 27, 235-41	4.7	78
179	Association of morning and evening blood pressure at home with asymptomatic organ damage in the J-HOP Study. <i>American Journal of Hypertension</i> , 2014 , 27, 939-47	2.3	64
178	An alpha-adrenergic blocker titrated by self-measured blood pressure recordings lowered blood pressure and microalbuminuria in patients with morning hypertension: the Japan Morning Surge-1 Study. <i>Journal of Hypertension</i> , 2008 , 26, 1257-65	1.9	62
177	Determinants of exaggerated difference in morning and evening blood pressure measured by self-measured blood pressure monitoring in medicated hypertensive patients: Jichi Morning Hypertension Research (J-MORE) Study. <i>American Journal of Hypertension</i> , 2005 , 18, 958-65	2.3	61
176	Predictive value of night-time heart rate for cardiovascular events in hypertension. The ABP-International study. <i>International Journal of Cardiology</i> , 2013 , 168, 1490-5	3.2	60
175	Nocturnal nondipping of heart rate predicts cardiovascular events in hypertensive patients. <i>Journal of Hypertension</i> , 2009 , 27, 2265-70	1.9	60
174	Comparison of valsartan and amlodipine on ambulatory and morning blood pressure in hypertensive patients. <i>American Journal of Hypertension</i> , 2004 , 17, 112-7	2.3	60
173	Low-grade inflammation is a risk factor for clinical stroke events in addition to silent cerebral infarcts in Japanese older hypertensives: the Jichi Medical School ABPM Study, wave 1. <i>Stroke</i> , 2007 , 38, 911-7	6.7	58
172	Sleep Blood Pressure Self-Measured at Home as a Novel Determinant of Organ Damage: Japan Morning Surge Home Blood Pressure (J-HOP) Study. <i>Journal of Clinical Hypertension</i> , 2015 , 17, 340-8	2.3	57
171	Masked nocturnal hypertension and target organ damage in hypertensives with well-controlled self-measured home blood pressure. <i>Hypertension Research</i> , 2007 , 30, 143-9	4.7	57
170	Nocturnal blood pressure elevation predicts progression of albuminuria in elderly people with type 2 diabetes. <i>Journal of Clinical Hypertension</i> , 2008 , 10, 12-20	2.3	56
169	Subclinical arterial damage in untreated masked hypertensive subjects detected by home blood pressure measurement. <i>American Journal of Hypertension</i> , 2007 , 20, 385-91	2.3	56
168	Short sleep duration is an independent predictor of stroke events in elderly hypertensive patients. <i>Journal of the American Society of Hypertension</i> , 2010 , 4, 255-62		52

167	Diabetic brain damage in hypertension: role of renin-angiotensin system. <i>Hypertension</i> , 2005 , 45, 887-938.5		52
166	Effect of Standard vs Intensive Blood Pressure Control on the Risk of Recurrent Stroke: A Randomized Clinical Trial and Meta-analysis. <i>JAMA Neurology</i> , 2019 , 76, 1309-1318	17.2	50
165	Differential impacts of adiponectin on low-grade albuminuria between obese and nonobese persons without diabetes. <i>Journal of Clinical Hypertension</i> , 2007 , 9, 775-82	2.3	50
164	Reproducibility of ambulatory blood pressure in treated and untreated hypertensive patients. <i>Journal of Hypertension</i> , 2010 , 28, 918-24	1.9	42
163	Masked hypertension in diabetes mellitus: a potential risk. <i>Journal of Clinical Hypertension</i> , 2007 , 9, 601-7.3		41
162	Glomerular hyperfiltration is a predictor of adverse cardiovascular outcomes. <i>Kidney International</i> , 2018 , 93, 195-203	9.9	40
161	Regular alcohol drinking is a determinant of masked morning hypertension detected by home blood pressure monitoring in medicated hypertensive patients with well-controlled clinic blood pressure: the Jichi Morning Hypertension Research (J-MORE) study. <i>Hypertension Research</i> , 2006 , 29, 679-86	4.7	39
160	Collagen metabolism in extracellular matrix may be involved in arterial stiffness in older hypertensive patients with left ventricular hypertrophy. <i>Hypertension Research</i> , 2005 , 28, 995-1001	4.7	38
159	Early morning surge in blood pressure. <i>Blood Pressure Monitoring</i> , 2001 , 6, 349-53	1.3	37
158	Comparison of the effects of cilnidipine and amlodipine on ambulatory blood pressure. <i>Hypertension Research</i> , 2005 , 28, 1003-8	4.7	36
157	Additional impact of morning haemostatic risk factors and morning blood pressure surge on stroke risk in older Japanese hypertensive patients. <i>European Heart Journal</i> , 2011 , 32, 574-80	9.5	35
156	Association between the morning-evening difference in home blood pressure and cardiac damage in untreated hypertensive patients. <i>Journal of Hypertension</i> , 2009 , 27, 712-20	1.9	35
155	Nocturnal hypoxia is associated with silent cerebrovascular disease in a high-risk Japanese community-dwelling population. <i>American Journal of Hypertension</i> , 2005 , 18, 1489-95	2.3	35
154	Comparison of the effects of pioglitazone and metformin on insulin resistance and hormonal markers in patients with impaired glucose tolerance and early diabetes. <i>Hypertension Research</i> , 2007 , 30, 23-30	4.7	34
153	Validation of an oscillometric home blood pressure monitor in an end-stage renal disease population and the effect of arterial stiffness on its accuracy. <i>Blood Pressure Monitoring</i> , 2007 , 12, 227-32.3		34
152	Increased low-grade inflammation and plasminogen-activator inhibitor-1 level in nondippers with sleep apnea syndrome. <i>Journal of Hypertension</i> , 2008 , 26, 1181-7	1.9	33
151	Smoking and antihypertensive medication: interaction between blood pressure reduction and arterial stiffness. <i>Hypertension Research</i> , 2005 , 28, 631-8	4.7	32
150	Riser Pattern Is a Novel Predictor of Adverse Events in Heart Failure Patients With Preserved Ejection Fraction. <i>Circulation Journal</i> , 2017 , 81, 220-226	2.9	31

149	Comparison of wrist-type and arm-type 24-h blood pressure monitoring devices for ambulatory use. <i>Blood Pressure Monitoring</i> , 2013 , 18, 57-62	1.3	30
148	Smoking is associated with silent cerebrovascular disease in a high-risk Japanese community-dwelling population. <i>Hypertension Research</i> , 2004 , 27, 747-54	4.7	29
147	The first multicenter, randomized, controlled trial of home telemonitoring for Japanese patients with heart failure: home telemonitoring study for patients with heart failure (HOMES-HF). <i>Heart and Vessels</i> , 2018 , 33, 866-876	2.1	28
146	What is the optimal interval between successive home blood pressure readings using an automated oscillometric device?. <i>Journal of Hypertension</i> , 2009 , 27, 1172-7	1.9	28
145	Orthostatic hypertension detected by self-measured home blood pressure monitoring: a new cardiovascular risk factor for elderly hypertensives. <i>Hypertension Research</i> , 2008 , 31, 1509-16	4.7	28
144	Effects of new calcium channel blocker, azelnidipine, and amlodipine on baroreflex sensitivity and ambulatory blood pressure. <i>Journal of Cardiovascular Pharmacology</i> , 2007 , 49, 394-400	3.1	28
143	Comparison of candesartan with lisinopril on ambulatory blood pressure and morning surge in patients with systemic hypertension. <i>American Journal of Cardiology</i> , 2003 , 92, 621-4	3	27
142	The effect of pulse rate and blood pressure dipping status on the risk of stroke and cardiovascular disease in Japanese hypertensive patients. <i>American Journal of Hypertension</i> , 2010 , 23, 749-55	2.3	26
141	Gestational hypertension as a subclinical preeclampsia in view of serum levels of angiogenesis-related factors. <i>Hypertension Research</i> , 2011 , 34, 212-7	4.7	26
140	Relationship between morning hypertension identified by home blood pressure monitoring and brain natriuretic peptide and estimated glomerular filtration rate: the Japan Morning Surge 1 (JMS-1) Study. <i>Journal of Clinical Hypertension</i> , 2008 , 10, 34-42	2.3	26
139	Differential effects of a long-acting angiotensin converting enzyme inhibitor (temocapril) and a long-acting calcium antagonist (amlodipine) on ventricular ectopic beats in older hypertensive patients. <i>Hypertension Research</i> , 2002 , 25, 329-33	4.7	26
138	Clarithromycin associated with torsades de pointes. <i>Japanese Circulation Journal</i> , 1999 , 63, 421-2		26
137	Increased heart rate variability during sleep is a predictor for future cardiovascular events in patients with type 2 diabetes. <i>Hypertension Research</i> , 2010 , 33, 737-42	4.7	25
136	Differential impact of left ventricular mass and relative wall thickness on cardiovascular prognosis in diabetic and nondiabetic hypertensive subjects. <i>American Heart Journal</i> , 2007 , 154, 79.e9-15	4.9	25
135	Association between aldosterone induced by antihypertensive medication and arterial stiffness reduction: the J-CORE study. <i>Atherosclerosis</i> , 2011 , 215, 184-8	3.1	24
134	Type 2 diabetes is associated with left ventricular concentric remodeling in hypertensive patients. <i>American Journal of Hypertension</i> , 2005 , 18, 23-9	2.3	24
133	Sleep pulse pressure and awake mean pressure as independent predictors for stroke in older hypertensive patients. <i>American Journal of Hypertension</i> , 2004 , 17, 439-45	2.3	24
132	Rationale and design of a randomized trial to test the safety and non-inferiority of canagliflozin in patients with diabetes with chronic heart failure: the CANDLE trial. <i>Cardiovascular Diabetology</i> , 2016 , 15, 57	8.7	24

131	Is home blood pressure variability itself an interventional target beyond lowering mean home blood pressure during anti-hypertensive treatment?. <i>Hypertension Research</i> , 2012 , 35, 862-6	4.7	23
130	Prediction of strokes versus cardiac events by ambulatory monitoring of blood pressure: results from an international database. <i>Blood Pressure Monitoring</i> , 2007 , 12, 397-9	1.3	23
129	Altered aortic properties in elderly orthostatic hypertension. <i>Hypertension Research</i> , 2005 , 28, 15-9	4.7	23
128	Aggressive blood pressure-lowering therapy guided by home blood pressure monitoring improves target organ damage in hypertensive patients with type 2 diabetes/prediabetes. <i>Journal of Clinical Hypertension</i> , 2012 , 14, 422-8	2.3	21
127	Soluble tumor necrosis factor receptor 1 level is associated with left ventricular hypertrophy: the northern Manhattan study. <i>American Journal of Hypertension</i> , 2009 , 22, 763-9	2.3	21
126	The effects of the L/N-type calcium channel blocker (cilnidipine) on sympathetic hyperactive morning hypertension: results from ACHIEVE-ONE. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 133-42	2.3	20
125	Effects of Celiprolol and Bisoprolol on Blood Pressure, Vascular Stiffness, and Baroreflex Sensitivity. <i>American Journal of Hypertension</i> , 2015 , 28, 858-67	2.3	20
124	Orthostatic hypertension: home blood pressure monitoring for detection and assessment of treatment with doxazosin. <i>Hypertension Research</i> , 2012 , 35, 100-6	4.7	20
123	Association of an abnormal blood glucose level and morning blood pressure surge in elderly subjects with hypertension. <i>American Journal of Hypertension</i> , 2009 , 22, 611-6	2.3	20
122	Urinary albumin excretion during angiotensin II receptor blockade: comparison of combination treatment with a diuretic or a calcium-channel blocker. <i>American Journal of Hypertension</i> , 2011 , 24, 466-73	2.3	20
121	Morning hypertension assessed by home monitoring is a strong predictor of concentric left ventricular hypertrophy in patients with untreated hypertension. <i>Journal of Clinical Hypertension</i> , 2010 , 12, 776-83	2.3	20
120	Cardiovascular risks of dipping status and chronic kidney disease in elderly Japanese hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2008 , 10, 787-94	2.3	20
119	The Japan Morning Surge-1 (JMS-1) study: protocol description. <i>Hypertension Research</i> , 2006 , 29, 153-9	4.7	20
118	Riser Blood Pressure Pattern Is Associated With Mild Cognitive Impairment in Heart Failure Patients. <i>American Journal of Hypertension</i> , 2016 , 29, 194-201	2.3	19
117	The time course of flow-mediated vasodilation and endothelial dysfunction in patients with a cardiovascular risk factor. <i>Journal of the American Society of Hypertension</i> , 2012 , 6, 109-16		19
116	Association of home and ambulatory blood pressure changes with changes in cardiovascular biomarkers during antihypertensive treatment. <i>American Journal of Hypertension</i> , 2012 , 25, 306-12	2.3	19
115	Effect of Intensive Salt-Restriction Education on Clinic, Home, and Ambulatory Blood Pressure Levels in Treated Hypertensive Patients During a 3-Month Education Period. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 385-92	2.3	19
114	Effect of a novel calcium channel blocker on abnormal nocturnal blood pressure in hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 465-72	2.3	18

113	A bedtime dose of ARB was better than a morning dose in improving baroreflex sensitivity and urinary albumin excretion--the J-TOP study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 488-92	2.2	18
112	Masked tachycardia. A predictor of adverse outcome in hypertension. <i>Journal of Hypertension</i> , 2017 , 35, 487-492	1.9	17
111	Febuxostat does not delay progression of carotid atherosclerosis in patients with asymptomatic hyperuricemia: A randomized, controlled trial. <i>PLoS Medicine</i> , 2020 , 17, e1003095	11.6	17
110	Ambulatory blood pressure monitoring in diabetes and obesity-a review. <i>International Journal of Hypertension</i> , 2011 , 2011, 954757	2.4	17
109	Add-On Use of Eplerenone Is Effective for Lowering Home and Ambulatory Blood Pressure in Drug-Resistant Hypertension. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 1250-1257	2.3	17
108	Cardiovascular prognosis of sustained and white-coat hypertension in patients with type 2 diabetes mellitus. <i>Blood Pressure Monitoring</i> , 2008 , 13, 15-20	1.3	16
107	Factors associated with baroreflex sensitivity: association with morning blood pressure. <i>Hypertension Research</i> , 2007 , 30, 723-8	4.7	16
106	Rationale and design of a multicenter randomized study for evaluating vascular function under uric acid control using the xanthine oxidase inhibitor, febuxostat: the PRIZE study. <i>Cardiovascular Diabetology</i> , 2016 , 15, 87	8.7	16
105	Riser Pattern: Another Determinant of Heart Failure With Preserved Ejection Fraction. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 994-999	2.3	16
104	Impact of arterial stiffness reduction on urinary albumin excretion during antihypertensive treatment: the Japan morning Surge-1 study. <i>Journal of Hypertension</i> , 2010 , 28, 1752-60	1.9	15
103	Effect of doxazosin on the left ventricular structure and function in morning hypertensive patients: the Japan Morning Surge 1 study. <i>Journal of Hypertension</i> , 2008 , 26, 1463-71	1.9	15
102	Two cases of malignant hypertension with reversible diffuse leukoencephalopathy exhibiting a reversible nocturnal blood pressure "riser" pattern. <i>Hypertension Research</i> , 2002 , 25, 467-73	4.7	15
101	Rationale and design of a multicenter randomized controlled study to evaluate the preventive effect of ipragliflozin on carotid atherosclerosis: the PROTECT study. <i>Cardiovascular Diabetology</i> , 2016 , 15, 133	8.7	15
100	Effects of Antihypertensive Therapy on Blood Pressure Variability. <i>Current Hypertension Reports</i> , 2016 , 18, 75	4.7	15
99	High Salt Intake Is Independently Associated With Hypertensive Target Organ Damage. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 315-21	2.3	15
98	Prognostic impact of sex-ambulatory blood pressure interactions in 10 cohorts of 17 312 patients diagnosed with hypertension: systematic review and meta-analysis. <i>Journal of Hypertension</i> , 2015 , 33, 212-20	1.9	14
97	Correlations between different measures of clinic, home, and ambulatory blood pressure in hypertensive patients. <i>Blood Pressure Monitoring</i> , 2011 , 16, 142-8	1.3	14
96	An increased visceral-subcutaneous adipose tissue ratio is associated with difficult-to-treat hypertension in men. <i>Journal of Hypertension</i> , 2010 , 28, 1340-6	1.9	14

95	Why is blood pressure so hard to control in patients with type 2 diabetes?. <i>Journal of the Cardiometabolic Syndrome</i> , 2007 , 2, 114-8		14
94	Home telemonitoring study for Japanese patients with heart failure (HOMES-HF): protocol for a multicentre randomised controlled trial. <i>BMJ Open</i> , 2013 , 3,	3	13
93	Impaired flow-mediated vasodilatation is associated with increased left ventricular mass in a multiethnic population. The Northern Manhattan Study. <i>American Journal of Hypertension</i> , 2010 , 23, 413-9 ³		13
92	Association between asleep blood pressure and brain natriuretic peptide during antihypertensive treatment: the Japan Morning Surge-Target Organ Protection (J-TOP) study. <i>Journal of Hypertension</i> , 2012 , 30, 1015-21	1.9	13
91	Masked hypertension defined by ambulatory blood pressure monitoring is associated with an increased serum glucose level and urinary albumin-creatinine ratio. <i>Journal of Clinical Hypertension</i> , 2010 , 12, 578-87	2.3	13
90	The influence of wave reflection on left ventricular hypertrophy in hypertensive patients is modified by age and gender. <i>Hypertension Research</i> , 2008 , 31, 649-56	4.7	13
89	Coexistence of PM2.5 and low temperature is associated with morning hypertension in hypertensives. <i>Clinical and Experimental Hypertension</i> , 2015 , 37, 468-72	2.2	12
88	New-onset Takayasu's Arteritis as Acute Myocardial Infarction. <i>Internal Medicine</i> , 2018 , 57, 1415-1420	1.1	12
87	An adverse pregnancy-associated outcome due to overlooked primary aldosteronism. <i>Internal Medicine</i> , 2014 , 53, 2499-504	1.1	12
86	Correlation of Central Blood Pressure to Hypertensive Target Organ Damages During Antihypertensive Treatment: The J-TOP Study. <i>American Journal of Hypertension</i> , 2015 , 28, 980-6	2.3	12
85	Is very low dose hydrochlorothiazide combined with candesartan effective in uncontrolled hypertensive patients?. <i>Blood Pressure Monitoring</i> , 2010 , 15, 308-11	1.3	12
84	Factors associated with incident ischemic stroke in hospitalized heart failure patients: a pilot study. <i>Hypertension Research</i> , 2008 , 31, 289-94	4.7	12
83	Circadian variation of blood pressure and neurohumoral factors during the acute phase of stroke. <i>Clinical and Experimental Hypertension</i> , 2002 , 24, 109-14	2.2	12
82	Association of Extreme Nocturnal Dipping With Cardiovascular Events Strongly Depends on Age. <i>Hypertension</i> , 2020 , 75, 324-330	8.5	12
81	Association between nondipper pulse rate and measures of cardiac overload: The J-HOP Study. <i>Journal of Clinical Hypertension</i> , 2017 , 19, 402-409	2.3	11
80	Home BP monitoring using a telemonitoring system is effective for controlling BP in a remote island in Japan. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 814-9	2.3	11
79	Controlling evening BP as well as morning BP is important in hypertensive patients with prediabetes/diabetes: the JMS-1 study. <i>American Journal of Hypertension</i> , 2010 , 23, 522-7	2.3	11
78	Increase Trend in Home Blood Pressure on a Single Occasion Is Associated With B-Type Natriuretic Peptide and the Estimated Glomerular Filtration Rate. <i>American Journal of Hypertension</i> , 2015 , 28, 1098-105	2.3	10

77	Exaggerated blood pressure variability is associated with memory impairment in very elderly patients. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 637-644	2.3	10
76	High central blood pressure is associated with incident cardiovascular events in treated hypertensives: the ABC-J II Study. <i>Hypertension Research</i> , 2018 , 41, 947-956	4.7	10
75	Differential effects of strict blood pressure lowering by losartan/hydrochlorothiazide combination therapy and high-dose amlodipine monotherapy on microalbuminuria: the ALPHABET study. <i>Journal of the American Society of Hypertension</i> , 2012 , 6, 73-82		10
74	Masked hypertension defined by home blood pressure monitoring is associated with impaired flow-mediated vasodilatation in patients with cardiovascular risk factors. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 630-6	2.3	10
73	A home blood pressure monitor equipped with a graphic function facilitates faster blood pressure control than the conventional home blood pressure monitor. <i>Journal of Clinical Hypertension</i> , 2009 , 11, 422-5	2.3	10
72	Adrenergic blockade improved insulin resistance in patients with morning hypertension: the Japan Morning Surge-1 Study. <i>Journal of Hypertension</i> , 2009 , 27, 1252-7	1.9	10
71	Changes in self-monitored pulse pressure correlate with improvements in B-type natriuretic Peptide and urinary albumin in treated hypertensive patients. <i>American Journal of Hypertension</i> , 2007 , 20, 1268-75	2.3	10
70	Usefulness of fasting blood glucose to predict vascular outcomes among individuals without diabetes mellitus (from the Northern Manhattan Study). <i>American Journal of Cardiology</i> , 2007 , 100, 1404-9	3.9	10
69	Short telomere length is associated with renal impairment in Japanese subjects with cardiovascular risk. <i>PLoS ONE</i> , 2017 , 12, e0176138	3.7	10
68	Plasma B-type natriuretic peptide is a useful tool for assessing coronary heart disease risk in a Japanese general population. <i>Hypertension Research</i> , 2015 , 38, 74-9	4.7	9
67	Why the radial augmentation index is low in patients with diabetes: The J-HOP study. <i>Atherosclerosis</i> , 2016 , 246, 338-43	3.1	9
66	Exaggerated blood pressure variability in patients with pneumoconiosis: a pilot study. <i>American Journal of Hypertension</i> , 2014 , 27, 1456-63	2.3	9
65	Change in high-sensitive cardiac troponin T on hypertensive treatment. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 40-4	2.2	9
64	Short sleep duration and type 2 diabetes enhance the risk of cardiovascular events in hypertensive patients. <i>Diabetes Research and Clinical Practice</i> , 2012 , 98, 518-23	7.4	9
63	A case of vertebral artery dissection associated with morning blood pressure surge. <i>Hypertension Research</i> , 2005 , 28, 847-51	4.7	9
62	Comparison of valsartan and amlodipine on ambulatory blood pressure variability in hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2016 , 38, 721-724	2.2	9
61	Lung Disease and Hypertension. <i>Pulse</i> , 2014 , 2, 103-12	1.6	8
60	Determinants of negative white-coat effect in treated hypertensive patients: the Jichi Morning Hypertension Research (J-MORE) study. <i>American Journal of Hypertension</i> , 2009 , 22, 35-40	2.3	8

59	Cognitive dysfunction and physical disability are associated with mortality in extremely elderly patients. <i>Hypertension Research</i> , 2008 , 31, 1331-8	4.7	8
58	Impact of blood pressure vs. glycemic factors on target organ damage in patients with type 2 diabetes mellitus. <i>Journal of Clinical Hypertension</i> , 2006 , 8, 404-10	2.3	8
57	A novel and simple protocol for the validation of home blood pressure monitors in clinical practice. <i>Blood Pressure Monitoring</i> , 2012 , 17, 210-3	1.3	7
56	Metabolic syndrome less strongly associated with target organ damage than syndrome components in a healthy, working population. <i>Journal of Clinical Hypertension</i> , 2007 , 9, 337-44	2.3	7
55	Added predictive value of high uric acid for cardiovascular events in the Ambulatory Blood Pressure International Study. <i>Journal of Clinical Hypertension</i> , 2019 , 21, 966-974	2.3	6
54	Tom Pickering as a clinical scientist: masked hypertension. <i>Blood Pressure Monitoring</i> , 2010 , 15, 85-9	1.3	6
53	Plasma tissue inhibitor of matrix metalloproteinase-1 level is increased in normotensive non-dippers in association with impaired glucose metabolism. <i>Hypertension Research</i> , 2008 , 31, 2045-51	4.7	6
52	Association of high-sensitivity cardiac troponin T and N-terminal pro-brain-type natriuretic peptide with left ventricular structure: J-HOP study. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 354-61	2.3	5
51	Prognostic Value of Ambulatory Blood Pressure in the Obese: The Ambulatory Blood Pressure-International Study. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 111-8	2.3	5
50	Left ventricular diastolic function evaluated by the E/eRatio is impaired in patients with masked uncontrolled hypertension. <i>Clinical and Experimental Hypertension</i> , 2014 , 36, 538-44	2.2	4
49	Low-grade inflammation and ambulatory blood pressure response to antihypertensive treatment: the ALPHABET study. <i>American Journal of Hypertension</i> , 2013 , 26, 784-92	2.3	4
48	Locomotive syndrome is associated with large blood pressure variability in elderly hypertensives: the Japan Ambulatory Blood Pressure Prospective (JAMP) substudy. <i>Journal of Clinical Hypertension</i> , 2017 , 19, 388-394	2.3	3
47	A New Proposal for the Target Value for Home BP in Type 2 Diabetes Patients: The J-HOP Study. <i>American Journal of Hypertension</i> , 2015 , 28, 971-9	2.3	3
46	Changes in Central Hemodynamics in Women With Hypertensive Pregnancy Between Before and After Delivery. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 329-36	2.3	3
45	Cardiac sarcoidosis, the complete atrioventricular block of which was completely recovered by intravenous steroid pulse therapy. <i>Journal of Cardiology Cases</i> , 2016 , 13, 21-24	0.6	3
44	Blood Pressure Management in Patients with Type 2 Diabetes. <i>Internal Medicine</i> , 2015 , 54, 2285-9	1.1	3
43	Evening heart rate measured at home is associated with visceral obesity and abnormal fat distribution in patients with hypertension. <i>American Journal of Hypertension</i> , 2011 , 24, 783-8	2.3	3
42	Monitoring of the central pulse pressure is useful for detecting cardiac overload during antiadrenergic treatment: the Japan Morning Surge 1 study. <i>Journal of Hypertension</i> , 2008 , 26, 1928-34	1.9	3

41	Age-specific impact of self-monitored pulse pressure on hypertensive target organ damage in treated hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2007 , 9, 522-9	2.3	3
40	Life- and limb-saving endovascular therapy in a patient with acute abdominal aortic occlusion. <i>Cardiovascular Intervention and Therapeutics</i> , 2017 , 32, 190-195	2.5	2
39	Target home morning SBP be below 125 mmHg in type 2 diabetes patients. <i>Journal of Hypertension</i> , 2018 , 36, 1284-1290	1.9	2
38	Elevated pulse amplification in hypertensive patients with advanced kidney disease. <i>Hypertension Research</i> , 2018 , 41, 299-307	4.7	2
37	Association between decreased respiratory function and increased blood pressure variability. <i>Blood Pressure Monitoring</i> , 2018 , 23, 79-84	1.3	2
36	Maximum home blood pressure readings are associated with left atrial diameter in essential hypertensives. <i>Journal of Human Hypertension</i> , 2018 , 32, 432-439	2.6	2
35	The measurement of orthostatic blood pressure as a screening tool for masked hypertension with abnormal circadian blood pressure rhythm. <i>Hypertension Research</i> , 2016 , 39, 631-2	4.7	2
34	Relationship between endothelial dysfunction and kidney disease. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 156	2.3	2
33	Contribution of the ABP-International study to the definition of night-time tachycardia. <i>Journal of Hypertension</i> , 2014 , 32, 2099-100	1.9	2
32	Differences in effects of age and blood pressure on augmentation index. <i>American Journal of Hypertension</i> , 2014 , 27, 1479-85	2.3	2
31	Cardiac positron emission tomography as a prognostic indicator of cardiac sarcoidosis. <i>Journal of the American College of Cardiology</i> , 2014 , 63, 2589-2590	15.1	2
30	A call for wider use of ambulatory blood pressure monitoring in patients with diabetes. <i>Hypertension Research</i> , 2011 , 34, 1171-2	4.7	2
29	Poor blood pressure and urinary albumin excretion responses to home blood pressure-based antihypertensive therapy in depressive hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2010 , 12, 345-9	2.3	2
28	Can ischemic stroke be caused by acute reduction of blood pressure in the acute phase of cardiovascular disease?. <i>Journal of Clinical Hypertension</i> , 2008 , 10, 195-200	2.3	2
27	Determinants of self-measured pulse rate profile in medicated hypertensives: the Jichi Morning Surge-1 (JMS-1) study. <i>Clinical and Experimental Hypertension</i> , 2008 , 30, 255-65	2.2	2
26	Daytime blood pressure surges following hypoxic episodes in a case of pneumoconiosis with lacunar stroke recurrences. <i>Blood Pressure Monitoring</i> , 2017 , 22, 175-177	1.3	1
25	An α -glucosidase inhibitor could reduce T-wave alternans in type 2 diabetes patients. <i>Journal of Electrocardiology</i> , 2018 , 51, 21-26	1.4	1
24	Electron Microscopy of Contact Between a Monocyte and a Multinucleated Giant Cell in Cardiac Sarcoidosis. <i>Canadian Journal of Cardiology</i> , 2016 , 32, 1577.e19-1577.e20	3.8	1

23	A case of syncope induced in the supine position. <i>International Journal of Hypertension</i> , 2011 , 2011, 3421-4	1.0	1
22	Relationship between the change in left ventricular hypertrophy and asleep blood pressure after sodium restriction and/or diuretic treatment. <i>Blood Pressure Monitoring</i> , 2011 , 16, 172-9	1.3	1
21	Is nondipping harmful in normotensive, healthy subjects?. <i>American Journal of Hypertension</i> , 2010 , 23, 222-3	2.3	1
20	Impact of white-coat hypertension on microvascular complications in type 2 diabetes: response to Kramer et al. <i>Diabetes Care</i> , 2009 , 32, e24; author reply e25	14.6	1
19	Ambulatory versus home versus clinic blood pressure. <i>Hypertension</i> , 2012 , 59, e25; author reply e26	8.5	1
18	Carcinomatous lymphangitis mimicking pulmonary thromboembolism. <i>Japanese Circulation Journal</i> , 2001 , 65, 683-4		1
17	Myofibroblasts in fibrous tumors and fibrosis in various organs. <i>Pathology International</i> , 1981 , 31, 423-37	1.8	1
16	Night Time Blood Pressure Variability Is a Strong Predictor for Cardiovascular Events in Patients With Type 2 Diabetes		1
15	Evaluation of day-by-day variability of home blood pressure using a home blood pressure telemonitoring system. <i>Blood Pressure Monitoring</i> , 2016 , 21, 184-8	1.3	1
14	Long-Term Effect of Febuxostat on Endothelial Function in Patients With Asymptomatic Hyperuricemia: A Sub-Analysis of the PRIZE Study.. <i>Frontiers in Cardiovascular Medicine</i> , 2022 , 9, 882821	5.4	0
13	Regarding the article "Clinical implication of visit-to-visit blood pressure variability". <i>Hypertension Research</i> , 2019 , 42, 1089	4.7	
12	Recent clinical trial of central hemodynamics. <i>Vascular Failure</i> , 2017 , 1, 9-14	0.3	
11	Clinical implications of the change in glomerular filtration rate with adrenergic blockers in patients with morning hypertension: the Japan morning surge-1 study. <i>International Journal of Hypertension</i> , 2013 , 2013, 413469	2.4	
10	Masked Hypertension and Diabetes. <i>Current Hypertension Reviews</i> , 2011 , 7, 1-4	2.3	
9	Primary Aldosteronism Complicated by Severe Hypokalemic Myopathy 2010 , 20, 298-300		
8	Doxazosin and heart failure: to be or not to be. <i>Journal of Hypertension</i> , 2009 , 27, 434-435	1.9	
7	Trial finds valsartan and amlodipine equally effective for preventing cardiac disease among people with hypertension at high risk. <i>Evidence-based Cardiovascular Medicine</i> , 2004 , 8, 306-7; discussion 308-9		
6	P-194: Morning BP reduction: which drug can lower most?. <i>American Journal of Hypertension</i> , 2004 , 17, S104	2.3	

- 5 Effect of angiotensin-receptor blockade on reduced cerebrovascular reserve and neuronal mass in diabetic hypertensives. *American Journal of Hypertension*, **2004**, 17, S155-S156 2.3
- 4 Peripheral and central pulse pressure are not useful in predicting cardiovascular prognosis. *Vascular Failure*, **2020**, 3, 37-42 0.3
- 3 Is High Central BP but Normal Office Brachial BP a risk?-The ABC-J II Study-. *Vascular Failure*, **2018**, 2, 66-73 0.3
- 2 Ambulatory blood pressure variability increases over a 19-year follow-up in a clinic on a solitary island. *Blood Pressure Monitoring*, **2018**, 23, 283-287 1.3
- 1 New Insight into Effects of β Blockers on Arterial Functions. *Pulse*, **2016**, 3, 190-4 1.6