

# Maja Thiele

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8068951/maja-thiele-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

74  
papers

2,302  
citations

25  
h-index

47  
g-index

97  
ext. papers

3,434  
ext. citations

6.7  
avg, IF

4.95  
L-index

#	Paper	IF	Citations
74	Risk of infections and their role on subsequent mortality in biopsy-proven alcohol-related liver disease.. <i>United European Gastroenterology Journal</i> , <b>2022</b> ,	5.3	1
73	Stage-dependent expression of fibrogenic markers in alcohol-related liver disease.. <i>Pathology Research and Practice</i> , <b>2022</b> , 231, 153798	3.4	2
72	Treatment of alcohol use disorder in patients with liver disease.. <i>Current Opinion in Pharmacology</i> , <b>2022</b> , 62, 145-151	5.1	0
71	Artificial intelligence outperforms standard blood-based scores in identifying liver fibrosis patients in primary care.. <i>Scientific Reports</i> , <b>2022</b> , 12, 2914	4.9	0
70	Reply to: Correspondence on "EASL Clinical Practice Guidelines on non-invasive tests for evaluation of liver disease severity and prognosis - 2021 update". <i>Journal of Hepatology</i> , <b>2021</b> ,	13.4	
69	Binge drinking induces an acute burst of markers of hepatic fibrogenesis (PRO-C3). <i>Liver International</i> , <b>2021</b> ,	7.9	1
68	2D-shear wave elastography: number of acquisitions can be reduced according to clinical setting. <i>Insights Into Imaging</i> , <b>2021</b> , 12, 145	5.6	0
67	The Role of Diagnostic Biomarkers, Omics Strategies, and Single-Cell Sequencing for Nonalcoholic Fatty Liver Disease in Severely Obese Patients. <i>Journal of Clinical Medicine</i> , <b>2021</b> , 10,	5.1	2
66	Assessment of hepatic steatosis by controlled attenuation parameter using the M and XL probes: an individual patient data meta-analysis. <i>The Lancet Gastroenterology and Hepatology</i> , <b>2021</b> , 6, 185-198	18.8	36
65	A Dynamic Aspartate-to-Alanine Aminotransferase Ratio Provides Valid Predictions of Incident Severe Liver Disease. <i>Hepatology Communications</i> , <b>2021</b> , 5, 1021-1035	6	5
64	Level of MFAP4 in ascites independently predicts 1-year transplant-free survival in patients with cirrhosis. <i>JHEP Reports</i> , <b>2021</b> , 3, 100287	10.3	
63	AGO2 localizes to cytokinetic protrusions in a p38-dependent manner and is needed for accurate cell division. <i>Communications Biology</i> , <b>2021</b> , 4, 726	6.7	0
62	Prognostic performance of 7 biomarkers compared to liver biopsy in early alcohol-related liver disease. <i>Journal of Hepatology</i> , <b>2021</b> , 75, 1017-1025	13.4	7
61	PRO-C3 and ADAPT algorithm accurately identify patients with advanced fibrosis due to alcohol-related liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2021</b> , 54, 699-708	6.1	5
60	Mortality in biopsy-proven alcohol-related liver disease: a population-based nationwide cohort study of 3453 patients. <i>Gut</i> , <b>2021</b> , 70, 170-179	19.2	15
59	Refining the Baveno VI elastography criteria for the definition of compensated advanced chronic liver disease. <i>Journal of Hepatology</i> , <b>2021</b> , 74, 1109-1116	13.4	39
58	Risk of Cancer in Biopsy-Proven Alcohol-Related Liver Disease: A Population-Based Cohort Study of 3410 Persons. <i>Clinical Gastroenterology and Hepatology</i> , <b>2021</b> ,	6.9	5

57	Two-dimensional shear wave elastography predicts survival in advanced chronic liver disease. <i>Gut</i> , <b>2021</b> ,	19.2	16
56	2D Shear Wave Elastography Predicts Survival in Advanced Chronic Liver Disease: AIXPLORER PREDICT and VALIDATE <b>2021</b> , 59,		2
55	Effects of Ethanol Feeding in Early-Stage NAFLD Mice Induced by Western Diet. <i>Livers</i> , <b>2021</b> , 1, 27-39		
54	Progressive alcohol-related liver fibrosis is characterised by imbalanced collagen formation and degradation. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2021</b> , 54, 1070-1080	6.1	5
53	Diagnostic accuracy of routine liver function tests to identify patients with significant and advanced alcohol-related liver fibrosis. <i>Scandinavian Journal of Gastroenterology</i> , <b>2021</b> , 56, 1088-1095	2.4	1
52	Population screening for liver fibrosis: Toward early diagnosis and intervention for chronic liver diseases. <i>Hepatology</i> , <b>2021</b> ,	11.2	8
51	We need to address health disparities to combat alcohol-related harm. <i>Lancet Regional Health - Europe, The</i> , <b>2021</b> , 8, 100198		
50	EASL Clinical Practice Guidelines on non-invasive tests for evaluation of liver disease severity and prognosis - 2021 update. <i>Journal of Hepatology</i> , <b>2021</b> , 75, 659-689	13.4	102
49	Comprehensive lipidomics reveals phenotypic differences in hepatic lipid turnover in ALD and NAFLD during alcohol intoxication. <i>JHEP Reports</i> , <b>2021</b> , 3, 100325	10.3	3
48	Development and prognostic relevance of a histologic grading and staging system for alcohol-related liver disease. <i>Journal of Hepatology</i> , <b>2021</b> , 75, 810-819	13.4	7
47	Low accuracy of FIB-4 and NAFLD fibrosis scores for screening for liver fibrosis in the population.. <i>Clinical Gastroenterology and Hepatology</i> , <b>2021</b> ,	6.9	5
46	2D shear wave liver elastography by Aixplorer to detect portal hypertension in cirrhosis: An individual patient data meta-analysis. <i>Liver International</i> , <b>2020</b> , 40, 1435-1446	7.9	22
45	Describing the fecal metabolome in cryogenically collected samples from healthy participants. <i>Scientific Reports</i> , <b>2020</b> , 10, 885	4.9	8
44	Prediction of liver fibrosis severity in alcoholic liver disease by human microfibrillar-associated protein 4. <i>Liver International</i> , <b>2020</b> , 40, 1701-1712	7.9	12
43	Collagen proportionate area predicts clinical outcomes in patients with alcohol-related liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2020</b> , 52, 1728-1739	6.1	4
42	Elastography in Combination with Other Biomarkers: Role of Algorithms <b>2020</b> , 551-561		
41	Steatosis Assessment with Controlled Attenuation Parameter (CAP) in Various Diseases <b>2020</b> , 441-457		
40	Recent advances in alcohol-related liver disease (ALD): summary of a Gut round table meeting. <i>Gut</i> , <b>2020</b> , 69, 764-780	19.2	39

39	Metabolic and Genetic Risk Factors Are the Strongest Predictors of Severity of Alcohol-Related Liver Fibrosis. <i>Clinical Gastroenterology and Hepatology</i> , <b>2020</b> ,	6.9	9
38	Cost-Effectiveness of Noninvasive Screening for Alcohol-Related Liver Fibrosis. <i>Hepatology</i> , <b>2020</b> , 71, 2093-2104	11.2	10
37	Transient elastography for screening of liver fibrosis: Cost-effectiveness analysis from six prospective cohorts in Europe and Asia. <i>Journal of Hepatology</i> , <b>2019</b> , 71, 1141-1151	13.4	51
36	Spleen stiffness to liver stiffness ratio significantly differs between ALD and HCV and predicts disease-specific complications. <i>JHEP Reports</i> , <b>2019</b> , 1, 99-106	10.3	21
35	Combination of CCl with alcoholic and metabolic injuries mimics human liver fibrosis. <i>American Journal of Physiology - Renal Physiology</i> , <b>2019</b> , 317, G182-G194	5.1	18
34	Clinical Aspects of Alcoholic Liver Disease <b>2019</b> , 3-21		
33	Controlled attenuation parameter and alcoholic hepatic steatosis: Diagnostic accuracy and role of alcohol detoxification. <i>Journal of Hepatology</i> , <b>2018</b> , 68, 1025-1032	13.4	55
32	Accuracy of the Enhanced Liver Fibrosis Test vs FibroTest, Elastography, and Indirect Markers in Detection of Advanced Fibrosis in Patients With Alcoholic Liver Disease. <i>Gastroenterology</i> , <b>2018</b> , 154, 1369-1379	13.3	97
31	Assessment of biopsy-proven liver fibrosis by two-dimensional shear wave elastography: An individual patient data-based meta-analysis. <i>Hepatology</i> , <b>2018</b> , 67, 260-272	11.2	213
30	Non-invasive diagnosis of liver fibrosis in patients with alcohol-related liver disease by transient elastography: an individual patient data meta-analysis. <i>The Lancet Gastroenterology and Hepatology</i> , <b>2018</b> , 3, 614-625	18.8	54
29	Antifibrotic and molecular aspects of rifaximin in alcoholic liver disease: study protocol for a randomized controlled trial. <i>Trials</i> , <b>2018</b> , 19, 143	2.8	10
28	Reliability Criteria for Liver Stiffness Measurements with Real-Time 2D Shear Wave Elastography in Different Clinical Scenarios of Chronic Liver Disease. <i>Ultraschall in Der Medizin</i> , <b>2017</b> , 38, 648-654	3.8	35
27	Contemporary use of elastography in liver fibrosis and portal hypertension. <i>Clinical Physiology and Functional Imaging</i> , <b>2017</b> , 37, 235-242	2.4	14
26	EFSUMB Guidelines and Recommendations on the Clinical Use of Liver Ultrasound Elastography, Update 2017 (Long Version). <i>Ultraschall in Der Medizin</i> , <b>2017</b> , 38, e16-e47	3.8	383
25	EFSUMB Guidelines and Recommendations on the Clinical Use of Liver Ultrasound Elastography, Update 2017 (Short Version). <i>Ultraschall in Der Medizin</i> , <b>2017</b> , 38, 377-394	3.8	62
24	Pharmacological interventions for alcoholic liver disease (alcohol-related liver disease): an attempted network meta-analysis. <i>The Cochrane Library</i> , <b>2017</b> , 3, CD011646	5.2	7
23	High risk of misinterpreting liver and spleen stiffness using 2D shear-wave and transient elastography after a moderate or high calorie meal. <i>PLoS ONE</i> , <b>2017</b> , 12, e0173992	3.7	34
22	Prediction of presence of oesophageal varices just by shear-wave elastography of the liver and spleen. <i>Liver International</i> , <b>2017</b> , 37, 1406-1407	7.9	1

21	Shear-wave elastography of the liver and spleen identifies clinically significant portal hypertension: A prospective multicentre study. <i>Liver International</i> , <b>2017</b> , 37, 396-405	7.9	90
20	Is liver stiffness equal to liver fibrosis?. <i>Hepatology</i> , <b>2017</b> , 65, 749	11.2	3
19	Feasibility of transient elastography versus real-time two-dimensional shear wave elastography in difficult-to-scan patients. <i>Scandinavian Journal of Gastroenterology</i> , <b>2016</b> , 51, 1354-9	2.4	17
18	Algorithm to rule out clinically significant portal hypertension combining Shear-wave elastography of liver and spleen: a prospective multicentre study. <i>Gut</i> , <b>2016</b> , 65, 1057-8	19.2	43
17	Transient and 2-Dimensional Shear-Wave Elastography Provide Comparable Assessment of Alcoholic Liver Fibrosis and Cirrhosis. <i>Gastroenterology</i> , <b>2016</b> , 150, 123-33	13.3	131
16	Reply. <i>Gastroenterology</i> , <b>2016</b> , 150, 1252-1253	13.3	
15	Beta-blockers alone or in combination with isosorbide mononitrate for secondary prevention of bleeding from gastro-esophageal varices in adults with cirrhosis and gastro-oesophageal varices. <i>The Cochrane Library</i> , <b>2016</b> ,	5.2	1
14	How to achieve a smooth transition from fellowship to faculty. <i>United European Gastroenterology Journal</i> , <b>2016</b> , 4, 717-718	5.3	1
13	O018 : 2D-shear wave elastography is equivalent or superior to transient elastography for liver fibrosis assessment: Results from an individual patient data based meta-analysis. <i>Journal of Hepatology</i> , <b>2015</b> , 62, S199-S200	13.4	8
12	Non-selective beta-blockers may reduce risk of hepatocellular carcinoma: a meta-analysis of randomized trials. <i>Liver International</i> , <b>2015</b> , 35, 2009-16	7.9	43
11	Large variations in risk of hepatocellular carcinoma and mortality in treatment naïve hepatitis B patients: systematic review with meta-analyses. <i>PLoS ONE</i> , <b>2014</b> , 9, e107177	3.7	26
10	The continuous reaction times method for diagnosing, grading, and monitoring minimal/covert hepatic encephalopathy. <i>Metabolic Brain Disease</i> , <b>2013</b> , 28, 231-4	3.9	46
9	Can non-selective beta-blockers prevent hepatocellular carcinoma in patients with cirrhosis?. <i>Medical Hypotheses</i> , <b>2013</b> , 81, 871-4	3.8	17
8	Antiviral therapy for prevention of hepatocellular carcinoma and mortality in chronic hepatitis B: systematic review and meta-analysis. <i>BMJ Open</i> , <b>2013</b> , 3,	3	27
7	Predictors of health-related quality of life in outpatients with cirrhosis: results from a prospective cohort. <i>Hepatitis Research and Treatment</i> , <b>2013</b> , 2013, 479639		16
6	Meta-analysis: banding ligation and medical interventions for the prevention of rebleeding from oesophageal varices. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2012</b> , 35, 1155-65	6.1	70
5	Common variable immunodeficiency diagnosed by capsule endoscopy. <i>Endoscopy</i> , <b>2011</b> , 43 Suppl 2 UCTN, E402	3.4	1
4	Higher intramuscular triacylglycerol in women does not impair insulin sensitivity and proximal insulin signaling. <i>Journal of Applied Physiology</i> , <b>2009</b> , 107, 824-31	3.7	53

3	Sex differences in hormone-sensitive lipase expression, activity, and phosphorylation in skeletal muscle at rest and during exercise. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2006</b> , 291, E1106-14	6	74
2	Higher skeletal muscle alpha2AMPK activation and lower energy charge and fat oxidation in men than in women during submaximal exercise. <i>Journal of Physiology</i> , <b>2006</b> , 574, 125-38	3-9	125
1	A paired liver biopsy and plasma proteomics study reveals circulating biomarkers for alcohol-related liver disease		5