

Shon Lewis

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8072423/shon-lewis-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

106
papers

7,156
citations

42
h-index

84
g-index

124
ext. papers

8,201
ext. citations

5.6
avg, IF

5.41
L-index

#	Paper	IF	Citations
106	Cognitive function in early-phase schizophrenia-spectrum disorder: IQ subtypes, brain volume and immune markers.. <i>Psychological Medicine</i> , 2022 , 1-10	6.9	0
105	The EMPOWER blended digital intervention for relapse prevention in schizophrenia: a feasibility cluster randomised controlled trial in Scotland and Australia.. <i>Lancet Psychiatry</i> , 2022 , 9, 477-486	23.3	2
104	Digital smartphone intervention to recognise and manage early warning signs in schizophrenia to prevent relapse: the EMPOWER feasibility cluster RCT. <i>Health Technology Assessment</i> , 2022 , 26, 1-174	4.4	0
103	Cross-sectional study comparing cognitive function in treatment responsive versus treatment non-responsive schizophrenia: evidence from the STRATA study. <i>BMJ Open</i> , 2021 , 11, e054160	3	0
102	Co-designing new tools for collecting, analysing and presenting patient experience data in NHS services: working in partnership with patients and carers. <i>Research Involvement and Engagement</i> , 2021 , 7, 85	4.4	0
101	Peripheral immune markers and antipsychotic non-response in psychosis. <i>Schizophrenia Research</i> , 2021 , 230, 1-8	3.6	8
100	Deconstructing depression and negative symptoms of schizophrenia; differential and longitudinal immune correlates, and response to minocycline treatment. <i>Brain, Behavior, and Immunity</i> , 2021 , 91, 498-504	16.6	10
99	Symptom Remission and Brain Cortical Networks at First Clinical Presentation of Psychosis: The OPTiMISE Study. <i>Schizophrenia Bulletin</i> , 2021 , 47, 444-455	1.3	1
98	Adaptive Symptom Monitoring Using Hidden Markov Models - An Application in Ecological Momentary Assessment. <i>IEEE Journal of Biomedical and Health Informatics</i> , 2021 , 25, 1770-1780	7.2	1
97	Dopamine and Glutamate in Antipsychotic-Responsive Compared With Antipsychotic-Nonresponsive Psychosis: A Multicenter Positron Emission Tomography and Magnetic Resonance Spectroscopy Study (STRATA). <i>Schizophrenia Bulletin</i> , 2021 , 47, 505-516	1.3	14
96	Digital mental health in China: a systematic review. <i>Psychological Medicine</i> , 2021 , 1-19	6.9	3
95	Early Signs Monitoring to Prevent Relapse in Psychosis and Promote Well-Being, Engagement, and Recovery: Protocol for a Feasibility Cluster Randomized Controlled Trial Harnessing Mobile Phone Technology Blended With Peer Support. <i>JMIR Research Protocols</i> , 2020 , 9, e15058	2	13
94	Smartphone-Enhanced Symptom Management In Psychosis: Open, Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2020 , 22, e17019	7.6	9
93	Developing a Theory-Informed Smartphone App for Early Psychosis: Learning Points From a Multidisciplinary Collaboration. <i>Frontiers in Psychiatry</i> , 2020 , 11, 602861	5	3
92	Digital methods to enhance the usefulness of patient experience data in services for long-term conditions: the DEPEND mixed-methods study. <i>Health Services and Delivery Research</i> , 2020 , 8, 1-128	1.5	3
91	Evidence-based guidelines for the pharmacological treatment of schizophrenia: Updated recommendations from the British Association for Psychopharmacology. <i>Journal of Psychopharmacology</i> , 2020 , 34, 3-78	4.6	85
90	A longitudinal comparison of two neurocognitive test batteries in patients with schizophrenia and healthy volunteers: Time effects on neuropsychological performance and their relation to functional outcome. <i>Schizophrenia Research</i> , 2020 , 216, 347-356	3.6	4

89	"They Are Not Hard-to-Reach Clients. We Have Just Got Hard-to-Reach Services." Staff Views of Digital Health Tools in Specialist Mental Health Services. <i>Frontiers in Psychiatry</i> , 2019 , 10, 344	5	30
88	Stress and cognitive biases in schizotypy: A two-site study of bias against disconfirmatory evidence and jumping to conclusions. <i>European Psychiatry</i> , 2019 , 62, 20-27	6	7
87	Stratification and prediction of remission in first-episode psychosis patients: the OPTiMiSE cohort study. <i>Translational Psychiatry</i> , 2019 , 9, 20	8.6	30
86	Digital biomarkers from geolocation data in bipolar disorder and schizophrenia: a systematic review. <i>Journal of the American Medical Informatics Association: JAMIA</i> , 2019 , 26, 1412-1420	8.6	23
85	Blending active and passive digital technology methods to improve symptom monitoring in early psychosis. <i>Microbial Biotechnology</i> , 2019 , 13, 1271-1275	3.3	17
84	Remission from antipsychotic treatment in first episode psychosis related to longitudinal changes in brain glutamate. <i>NPJ Schizophrenia</i> , 2019 , 5, 12	5.5	12
83	A Theory-Informed Digital Health Intervention in People with Severe Mental Health Problems. <i>Studies in Health Technology and Informatics</i> , 2019 , 264, 526-530	0.5	3
82	Minocycline for negative symptoms of schizophrenia and possible mechanistic actions: the BeneMin RCT. <i>Efficacy and Mechanism Evaluation</i> , 2019 , 6, 1-66	1.7	1
81	Low Intelligence and Poor Educational Achievement Precede Early Onset Schizophrenic Psychosis 2019 , 131-144		2
80	Towards a consensus around standards for smartphone apps and digital mental health. <i>World Psychiatry</i> , 2019 , 18, 97-98	14.4	142
79	Actissist: Proof-of-Concept Trial of a Theory-Driven Digital Intervention for Psychosis. <i>Schizophrenia Bulletin</i> , 2018 , 44, 1070-1080	1.3	85
78	Digital interventions in severe mental health problems: lessons from the Actissist development and trial. <i>World Psychiatry</i> , 2018 , 17, 230-231	14.4	12
77	Early Psychosis Service User Views on Digital Technology: Qualitative Analysis. <i>JMIR Mental Health</i> , 2018 , 5, e10091	6	28
76	The COMMAND trial of cognitive therapy to prevent harmful compliance with command hallucinations: predictors of outcome and mediators of change. <i>Psychological Medicine</i> , 2018 , 48, 1966-1974	6.74	15
75	The benefit of minocycline on negative symptoms of schizophrenia in patients with recent-onset psychosis (BeneMin): a randomised, double-blind, placebo-controlled trial. <i>Lancet Psychiatry</i> , 2018 , 5, 885-894	23.3	88
74	The COMMAND trial of cognitive therapy for harmful compliance with command hallucinations (CTCH): a qualitative study of acceptability and tolerability in the UK. <i>BMJ Open</i> , 2018 , 8, e021657	3	4
73	Culturally-adapted Family Intervention (CaFI) for African-Caribbeans diagnosed with schizophrenia and their families: a feasibility study protocol of implementation and acceptability. <i>Pilot and Feasibility Studies</i> , 2016 , 2, 39	1.9	20
72	Antidepressant Controlled Trial For Negative Symptoms In Schizophrenia (ACTIONS): a double-blind, placebo-controlled, randomised clinical trial. <i>Health Technology Assessment</i> , 2016 , 20, 1-46	4.4	15

71	The optimization of treatment and management of schizophrenia in Europe (OPTiMiSE) trial: rationale for its methodology and a review of the effectiveness of switching antipsychotics. <i>Schizophrenia Bulletin</i> , 2015 , 41, 549-58	1.3	38
70	The promise of biological markers for treatment response in first-episode psychosis: a systematic review. <i>Schizophrenia Bulletin</i> , 2015 , 41, 559-73	1.3	65
69	Calibration and cross-validation of MCCB and CogState in schizophrenia. <i>Psychopharmacology</i> , 2015 , 232, 3873-82	4.7	27
68	Using mobile technology to deliver a cognitive behaviour therapy-informed intervention in early psychosis (Actissist): study protocol for a randomised controlled trial. <i>Trials</i> , 2015 , 16, 404	2.8	28
67	Stratified medicine for mental disorders. <i>European Neuropsychopharmacology</i> , 2014 , 24, 5-50	1.2	121
66	Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. <i>Lancet Psychiatry</i> , 2014 , 1, 23-33	23.3	100
65	Mood variability predicts the course of suicidal ideation in individuals with first and second episode psychosis. <i>Psychiatry Research</i> , 2013 , 206, 240-5	9.9	12
64	Psychotic experiences and psychotic disorders at age 18 in relation to psychotic experiences at age 12 in a longitudinal population-based cohort study. <i>American Journal of Psychiatry</i> , 2013 , 170, 742-50	11.9	206
63	Pathways to violent behavior during first-episode psychosis: a report from the UK National EDEN Study. <i>JAMA Psychiatry</i> , 2013 , 70, 1287-93	14.5	69
62	Reducing duration of untreated psychosis: care pathways to early intervention in psychosis services. <i>British Journal of Psychiatry</i> , 2013 , 203, 58-64	5.4	108
61	A comparison of two delivery modalities of a mobile phone-based assessment for serious mental illness: native smartphone application vs text-messaging only implementations. <i>Journal of Medical Internet Research</i> , 2013 , 15, e60	7.6	95
60	A treatment approach to the patient with first-episode schizophrenia 2012 , 237-251		
59	A treatment approach to the patient with first-episode schizophrenia 2012 , 237-251		
58	The negative and positive self: a longitudinal study examining self-esteem, paranoia and negative symptoms in individuals with first-episode psychosis. <i>Microbial Biotechnology</i> , 2011 , 5, 150-5	3.3	29
57	A multi-centre, randomised controlled trial of cognitive therapy to prevent harmful compliance with command hallucinations. <i>BMC Psychiatry</i> , 2011 , 11, 155	4.2	14
56	Auditory false perceptions are mediated by psychosis risk factors. <i>Cognitive Neuropsychiatry</i> , 2011 , 16, 289-302	2	26
55	Integrated motivational interviewing and cognitive behavioural therapy for people with psychosis and comorbid substance misuse: randomised controlled trial. <i>BMJ, The</i> , 2010 , 341, c6325	5.9	155
54	Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. <i>Archives of General Psychiatry</i> , 2010 , 67, 241-51		491

53	Distress and metacognition in psychosis prone individuals: comparing high schizotypy to the at-risk mental state. <i>Journal of Nervous and Mental Disease</i> , 2010 , 198, 99-104	1.8	26
52	Treatment of Schizophrenia: Discussion 2010 , 441-445		
51	Second-generation antipsychotics: a therapeutic downturn?. <i>Psychological Medicine</i> , 2009 , 39, 1603-6	6.9	3
50	Evaluating integrated MI and CBT for people with psychosis and substance misuse: recruitment, retention and sample characteristics of the MIDAS trial. <i>Addictive Behaviors</i> , 2009 , 34, 859-66	4.2	39
49	Assessing illicit substance use in schizophrenia: the relationship between self report and detection by hair analysis. <i>Schizophrenia Research</i> , 2009 , 114, 180-1	3.6	7
48	A randomized controlled trial of the cost-utility of second-generation antipsychotics in people with psychosis and eligible for clozapine. <i>Value in Health</i> , 2008 , 11, 549-62	3.3	25
47	Cognitive and neural processes in non-clinical auditory hallucinations. <i>British Journal of Psychiatry</i> , 2007 , 51, s76-81	5.4	78
46	The Psychotic Symptom Rating Scales (PSYRATS): their usefulness and properties in first episode psychosis. <i>Schizophrenia Research</i> , 2007 , 89, 119-22	3.6	109
45	Prospects for a cognitive-developmental account of psychotic experiences. <i>British Journal of Clinical Psychology</i> , 2007 , 46, 155-73	3.6	65
44	Cost-effectiveness of first- v. second-generation antipsychotic drugs: results from a randomised controlled trial in schizophrenia responding poorly to previous therapy. <i>British Journal of Psychiatry</i> , 2007 , 191, 14-22	5.4	82
43	Hallucination proneness, schizotypy and meta-cognition. <i>Behaviour Research and Therapy</i> , 2007 , 45, 1401-8	5.8	25
42	First- vs Second-Generation Antipsychotic Drugs in Schizophrenia Reply. <i>Archives of General Psychiatry</i> , 2007 , 64, 979		5
41	The presence of neurological soft signs along the psychosis proneness continuum. <i>Schizophrenia Bulletin</i> , 2006 , 32, 573-7	1.3	52
40	Suicide behaviour over 18 months in recent onset schizophrenic patients: the effects of CBT. <i>Schizophrenia Research</i> , 2006 , 83, 15-27	3.6	72
39	Influence of age on outcome of psychological treatments in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2006 , 188, 250-4	5.4	43
38	Identification of young people at risk of psychosis. <i>Advances in Psychiatric Treatment</i> , 2006 , 12, 249-255		10
37	Cannabis use prior to first onset psychosis predicts spared neurocognition at 10-year follow-up. <i>Schizophrenia Research</i> , 2005 , 75, 135-7	3.6	105
36	Association between duration of untreated psychosis and outcome in cohorts of first-episode patients: a systematic review. <i>Archives of General Psychiatry</i> , 2005 , 62, 975-83		1062

35	Attitudes to atypical and conventional antipsychotic drug treatment in clinicians participating in the cutlass study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2005 , 9, 35-40	2.4	6
34	Psychological interventions in early psychosis. <i>Schizophrenia Bulletin</i> , 2005 , 31, 697-704	1.3	38
33	Essential elements of an early intervention service for psychosis: the opinions of expert clinicians. <i>BMC Psychiatry</i> , 2004 , 4, 17	4.2	55
32	Cognitive-behavioural therapy in first-episode and early schizophrenia. 18-month follow-up of a randomised controlled trial. <i>British Journal of Psychiatry</i> , 2004 , 184, 231-9	5.4	212
31	Neurocognitive function and outcome in first-episode schizophrenia: a 10-year follow-up of an epidemiological cohort. <i>Schizophrenia Research</i> , 2003 , 65, 75-86	3.6	157
30	The evolution of symptoms in the early course of non-affective psychosis. <i>Schizophrenia Research</i> , 2003 , 63, 171-9	3.6	40
29	Is treatment research in people at high risk of psychosis ethical?. <i>Journal of Mental Health</i> , 2003 , 12, 359-362	3.6	1
28	Cognitive-behavioural therapy and motivational intervention for schizophrenia and substance misuse. 18-month outcomes of a randomised controlled trial. <i>British Journal of Psychiatry</i> , 2003 , 183, 418-26	5.4	164
27	Randomised controlled trial of cognitive-behavioural therapy in early schizophrenia: acute-phase outcomes. <i>British Journal of Psychiatry</i> , 2002 , 43, s91-7	5.4	260
26	Postcode prescribing in psychiatry: Clozapine in an English county. <i>Psychiatric Bulletin</i> , 2000 , 24, 420-422		13
25	Cognitive Behavioral Intervention for Individuals with Severe Mental Illness Who Have a Substance Misuse Problem. <i>Psychiatric Rehabilitation Skills</i> , 2000 , 4, 216-233		10
24	Lack of normal pattern of cerebral asymmetry in familial schizophrenic patients and their relatives--The Maudsley Family Study. <i>Schizophrenia Research</i> , 1999 , 40, 111-20	3.6	80
23	Randomised controlled effectiveness trial of a needs-based psychosocial intervention service for carers of people with schizophrenia. <i>British Journal of Psychiatry</i> , 1999 , 174, 505-11	5.4	114
22	Brain changes in schizophrenia. Volumetric MRI study of families multiply affected with schizophrenia--the Maudsley Family Study 5. <i>British Journal of Psychiatry</i> , 1998 , 173, 132-8	5.4	89
21	Individual cognitive-behavioural interventions in early psychosis. <i>British Journal of Psychiatry</i> , 1998 , 172, 101-106	5.4	42
20	The Maudsley Family Study. I: Structural brain changes on magnetic resonance imaging in familial schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1997 , 21, 1297-315	5.5	15
19	Doctor-patient interactions in oncology. <i>Social Science and Medicine</i> , 1996 , 42, 1511-9	5.1	310
18	No news is not good news: information preferences of patients with cancer. <i>Psycho-Oncology</i> , 1995 , 4, 197-202	3.9	190

17	Increased intracerebral cerebrospinal fluid spaces predict unemployment and negative symptoms in psychotic illness. A prospective study. <i>British Journal of Psychiatry</i> , 1995 , 166, 750-8	5.4	36
16	Correlates of left-handedness in a large sample of schizophrenic patients. <i>Schizophrenia Research</i> , 1995 , 18, 37-41	3.6	26
15	Age at onset, sex, and familial psychiatric morbidity in schizophrenia. Camberwell Collaborative Psychosis Study. <i>British Journal of Psychiatry</i> , 1994 , 165, 466-73	5.4	135
14	Information preferences of patients with cancer. <i>Lancet, The</i> , 1994 , 344, 1576	4.0	52
13	Cannabis and acute psychosis. <i>Schizophrenia Research</i> , 1994 , 13, 161-7	3.6	77
12	Low birth weight and schizophrenia. <i>British Journal of Psychiatry</i> , 1994 , 165, 357-62	5.4	99
11	Life events and psychosis. Initial results from the Camberwell Collaborative Psychosis Study. <i>British Journal of Psychiatry</i> , 1993 , 162, 72-9	5.4	206
10	Sex differences in the familial risk of schizophrenia: data from RĀinB study. <i>British Journal of Psychiatry</i> , 1992 , 161, 131-2	5.4	2
9	Sex and schizophrenia: vive la diffĀence. <i>British Journal of Psychiatry</i> , 1992 , 161, 445-50	5.4	85
8	Pre-morbid adjustment and personality in psychosis. Effects of sex and diagnosis. <i>British Journal of Psychiatry</i> , 1991 , 158, 171-6	5.4	168
7	The right stuff? a prospective controlled trial of traineesPresearch. <i>Psychiatric Bulletin</i> , 1991 , 15, 478-480		13
6	Delusional misidentification, AIDS and the right hemisphere. <i>British Journal of Psychiatry</i> , 1990 , 157, 608-10	5.4	29
5	The Heredity of Behavior Disorders in Adults and Children By S. G. Vandenburg, S. M. Singer and D. L. Pauls New York: Plenum. 1986. Pp 301. \$35.00.. <i>British Journal of Psychiatry</i> , 1987 , 151, 277-277	5.4	
4	Transient Psychosis: Diagnosis, Management and Evaluation. Edited by Joe. P. Tupin, Uriel Halbreich and Jesus J. Pena. New York: Brunner/Mazel. 1984. Pp. 307. \$30.00.. <i>British Journal of Psychiatry</i> , 1985 , 146, 226-226	5.4	
3	Falling Incidence of Schizophrenia?. <i>British Journal of Psychiatry</i> , 1985 , 146, 560-561	5.4	3
2	Duration of untreated psychosis125-146		2
1	Baseline levels of C-reactive protein and proinflammatory cytokines are not associated with early response to amisulpride in patients with First Episode Psychosis: the OPTiMiSE cohort study. <i>Schizophrenia Bulletin Open</i> ,	2.2	2