

# Barbora Å umovÃ;

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8073757/publications.pdf>

Version: 2024-02-01

17  
papers

659  
citations

840776

11  
h-index

1125743

13  
g-index

18  
all docs

18  
docs citations

18  
times ranked

1287  
citing authors

#	ARTICLE	IF	CITATIONS
1	Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. <i>Nature Communications</i> , 2017, 8, 1130.	12.8	245
2	Sirt1 regulates canonical TGF- $\beta$ signalling to control fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 226-233.	0.9	115
3	The transcription factor GLI2 as a downstream mediator of transforming growth factor- $\beta$ -induced fibroblast activation in SSc. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 756-764.	0.9	53
4	S100A4 amplifies TGF- $\beta$ -induced fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1748-1755.	0.9	52
5	Tribbles homologue 3 stimulates canonical TGF- $\beta$ signalling to regulate fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 609-616.	0.9	38
6	Calgizzarin (S100A11): a novel inflammatory mediator associated with disease activity of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2017, 19, 79.	3.5	35
7	Composition of TWIST1 dimers regulates fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 244-251.	0.9	28
8	Interleukin 35 Synovial Fluid Levels Are Associated with Disease Activity of Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015, 10, e0132674.	2.5	28
9	Circulating S100 proteins effectively discriminate SLE patients from healthy controls: a cross-sectional study. <i>Rheumatology International</i> , 2019, 39, 469-478.	3.0	23
10	Pro-inflammatory S100A11 is elevated in inflammatory myopathies and reflects disease activity and extramuscular manifestations in myositis. <i>Cytokine</i> , 2019, 116, 13-20.	3.2	17
11	High levels of metastasis-inducing S100A4 protein and treatment outcome in early rheumatoid arthritis: data from the PERAC cohort. <i>Biomarkers</i> , 2015, 20, 47-51.	1.9	12
12	Interleukin-20 is triggered by TLR ligands and associates with disease activity in patients with rheumatoid arthritis. <i>Cytokine</i> , 2017, 97, 187-192.	3.2	10
13	S100A4 is elevated in axial spondyloarthritis: a potential link to disease severity. <i>BMC Rheumatology</i> , 2020, 4, 13.	1.6	1
14	S100A11 (calgizzarin) is released by circulating mononuclear cells and its elevated plasma levels distinguish systemic lupus erythematosus patients from healthy individuals. <i>Clinical and Experimental Rheumatology</i> , 2019, 37, 338-339.	0.8	1
15	Metastasis-inducing S100A4 protein is associated with treatment response in patients with recent-onset rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, A15.2-A15.	0.9	0
16	OP0240 Signal Transducer and Activator of Transcription 3 Regulates Fibroblast Activation as A Downstream Mediator of Transforming Growth Factor-Beta. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 153.1-153.	0.9	0
17	02.37 IL-37 is associated with increased atherogenesis rather than disease activity in patients with rheumatoid arthritis. , 2017, , .		0