
## Barbora Å umovÃ;

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8073757/publications.pdf Version: 2024-02-01



RAPBORA ÅUMOVÃ:

#	Article	IF	CITATIONS
1	S100A4 is elevated in axial spondyloarthritis: a potential link to disease severity. BMC Rheumatology, 2020, 4, 13.	1.6	1
2	Pro-inflammatory S100A11 is elevated in inflammatory myopathies and reflects disease activity and extramuscular manifestations in myositis. Cytokine, 2019, 116, 13-20.	3.2	17
3	Circulating S100 proteins effectively discriminate SLE patients from healthy controls: a cross-sectional study. Rheumatology International, 2019, 39, 469-478.	3.0	23
4	S100A11 (calgizzarin) is released by circulating mononuclear cells and its elevated plasma levels distinguish systemic lupus erythematosus patients from healthy individuals. Clinical and Experimental Rheumatology, 2019, 37, 338-339.	0.8	1
5	The transcription factor GLI2 as a downstream mediator of transforming growth factor-β-induced fibroblast activation in SSc. Annals of the Rheumatic Diseases, 2017, 76, 756-764.	0.9	53
6	Composition of TWIST1 dimers regulates fibroblast activation and tissue fibrosis. Annals of the Rheumatic Diseases, 2017, 76, 244-251.	0.9	28
7	Calgizzarin (S100A11): a novel inflammatory mediator associated with disease activity of rheumatoid arthritis. Arthritis Research and Therapy, 2017, 19, 79.	3.5	35
8	Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. Nature Communications, 2017, 8, 1130.	12.8	245
9	02.37â€Il-37 is associated with increased atherogenesis rather than disease activity in patients with rheumatoid arthritis. , 2017, , .		0
10	Interleukin-20 is triggered by TLR ligands and associates with disease activity in patients with rheumatoid arthritis. Cytokine, 2017, 97, 187-192.	3.2	10
11	Sirt1 regulates canonical TCF-β signalling to control fibroblast activation and tissue fibrosis. Annals of the Rheumatic Diseases, 2016, 75, 226-233.	0.9	115
12	Tribbles homologue 3 stimulates canonical TGF-β signalling to regulate fibroblast activation and tissue fibrosis. Annals of the Rheumatic Diseases, 2016, 75, 609-616.	0.9	38
13	High levels of metastasis-inducing S100A4 protein and treatment outcome in early rheumatoid arthritis: data from the PERAC cohort. Biomarkers, 2015, 20, 47-51.	1.9	12
14	S100A4 amplifies TGF-Î <sup>2</sup> -induced fibroblast activation in systemic sclerosis. Annals of the Rheumatic Diseases, 2015, 74, 1748-1755.	0.9	52
15	Interleukin 35 Synovial Fluid Levels Are Associated with Disease Activity of Rheumatoid Arthritis. PLoS ONE, 2015, 10, e0132674.	2.5	28
16	OP0240â€Signal Transducer and Activator of Transcription 3 Regulates Fibroblast Activation as A Downstream Mediator of Transforming Growth Factor-Beta. Annals of the Rheumatic Diseases, 2014, 73, 153.1-153.	0.9	0
17	Metastasis-inducing S100A4 protein is associated with treatment response in patients with recent-onset rheumatoid arthritis. Annals of the Rheumatic Diseases, 2012, 71, A15.2-A15.	0.9	0