

Sara Negri

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/807603/sara-negri-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

37
papers

1,092
citations

16
h-index

33
g-index

40
ext. papers

1,279
ext. citations

5.5
avg, IF

3.48
L-index

#	Paper	IF	Citations
37	Long read sequencing overcomes challenges in the diagnosis of SORD neuropathy.. <i>Journal of the Peripheral Nervous System</i> , 2022 ,	4.7	1
36	Selective Targeting of Cancer-Associated Fibroblasts by Engineered H-Ferritin Nanocages Loaded with Navitoclax. <i>Cells</i> , 2021 , 10,	7.9	9
35	Biallelic mutations in SORD cause a common and potentially treatable hereditary neuropathy with implications for diabetes. <i>Nature Genetics</i> , 2020 , 52, 473-481	36.3	38
34	Association of Hydroxychloroquine With QTc Interval in Patients With COVID-19. <i>Circulation</i> , 2020 , 142, 513-515	16.7	26
33	Development and validation of a simple and versatile method for the quantification of everolimus loaded in H-ferritin nanocages using UHPLC-MS/MS. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020 , 191, 113644	3.5	3
32	Exposure duration and absorbed dose assessment in pesticide-exposed agricultural workers: Implications for risk assessment and modeling. <i>International Journal of Hygiene and Environmental Health</i> , 2019 , 222, 494-502	6.9	11
31	Everolimus Nanoformulation in Biological Nanoparticles Increases Drug Responsiveness in Resistant and Low-Responsive Breast Cancer Cell Lines. <i>Pharmaceutics</i> , 2019 , 11,	6.4	10
30	Paclitaxel-Loaded Silk Fibroin Nanoparticles: Method Validation by UHPLC-MS/MS to Assess an Exogenous Approach to Load Cytotoxic Drugs. <i>Pharmaceutics</i> , 2019 , 11,	6.4	10
29	Validation of cleaning procedures used in an Italian Hospital Pharmacy for antineoplastic drug decontamination: a new tool for industrial hygiene. <i>Medicina Del Lavoro</i> , 2019 , 110, 93-101	1.9	2
28	Urinary levels of metal elements in the non-smoking general population in Italy: SIVR study 2012-2015. <i>Toxicology Letters</i> , 2018 , 298, 177-185	4.4	19
27	A new, sensitive and versatile assay for quantitative determination of Fluoro-Balanine (AFBA) in human urine by using the reversed-phase ultrahigh performance-tandem mass spectrometry (rp-UHPLC-MS/MS) system. <i>Toxicology Letters</i> , 2018 , 298, 164-170	4.4	1
26	Monitoring Surface Contamination by Antineoplastic Drugs in Italian Hospitals: Performance-Based Hygienic Guidance Values (HGVs) Project. <i>Annals of Work Exposures and Health</i> , 2017 , 61, 994-1002	2.4	16
25	Methodology to define biological reference values in the environmental and occupational fields: the contribution of the Italian Society for Reference Values (SIVR). <i>Medicina Del Lavoro</i> , 2017 , 108, 138-148	1.9	3
24	Pharmaceutical Industries Air Quality. <i>Comprehensive Analytical Chemistry</i> , 2016 , 589-621	1.9	1
23	Exposure to perfluorinated compounds: in vitro study on thyroid cells. <i>Environmental Science and Pollution Research</i> , 2015 , 22, 2287-94	5.1	27
22	Determination of Glucocorticoids in UPLC-MS in Environmental Samples from an Occupational Setting. <i>International Journal of Analytical Chemistry</i> , 2015 , 2015, 678579	1.4	2
21	Subclinical abnormalities in sarcoplasmic reticulum Ca(2+) release promote eccentric myocardial remodeling and pump failure death in response to pressure overload. <i>Journal of the American College of Cardiology</i> , 2014 , 63, 1569-79	15.1	37

20	PFOS induces behavioral alterations, including spontaneous hyperactivity that is corrected by dexamfetamine in zebrafish larvae. <i>PLoS ONE</i> , 2014 , 9, e94227	3.7	55
19	P388Reduced pressure overload-induced myocardial remodeling in K201-treated mice with the R4496C cardiac ryanodine receptor mutation. <i>Cardiovascular Research</i> , 2014 , 103, S71.2-S71	9.9	
18	Trabectedin and indole-3-carbinol combination in heavily pretreated metastatic breast cancer: Results of a pilot clinical study.. <i>Journal of Clinical Oncology</i> , 2014 , 32, e12015-e12015	2.2	
17	Exposure to benzene and childhood leukaemia: a pilot case-control study. <i>BMJ Open</i> , 2013 , 3,	3	27
16	Oxidative activity of ammonium persulfate salt on mast cells and basophils: implication in hairdressersTasthma. <i>International Archives of Allergy and Immunology</i> , 2013 , 160, 409-19	3.7	16
15	Prenatal exposure to PFOS or PFOA alters motor function in mice in a sex-related manner. <i>Neurotoxicity Research</i> , 2011 , 19, 452-61	4.3	91
14	Short communication: flecainide exerts an antiarrhythmic effect in a mouse model of catecholaminergic polymorphic ventricular tachycardia by increasing the threshold for triggered activity. <i>Circulation Research</i> , 2011 , 109, 291-5	15.7	84
13	Dimethylformamide metabolism following self-harm using a veterinary euthanasia product. <i>Clinical Toxicology</i> , 2010 , 48, 725-9	2.9	9
12	Perfluorooctane sulfonate and perfluorooctanoic acid in surgical thyroid specimens of patients with thyroid diseases. <i>Thyroid</i> , 2009 , 19, 1407-12	6.2	21
11	Evaluation of urinary biomarkers of exposure to benzene: correlation with blood benzene and influence of confounding factors. <i>International Archives of Occupational and Environmental Health</i> , 2009 , 82, 985-95	3.2	56
10	Saliva as an analytical tool to measure occupational exposure to toluene. <i>International Archives of Occupational and Environmental Health</i> , 2008 , 81, 1021-8	3.2	14
9	Arrhythmogenesis in catecholaminergic polymorphic ventricular tachycardia: insights from a RyR2 R4496C knock-in mouse model. <i>Circulation Research</i> , 2006 , 99, 292-8	15.7	256
8	Mercapturic acids of styrene in man: comparability of the results obtained by LC/MS/MS and by HPLC-fluorimeter, and stability of samples under different storage conditions. <i>Toxicology Letters</i> , 2006 , 162, 225-33	4.4	7
7	Determination of perfluorooctanoic acid and perfluorooctanesulfonate in human tissues by liquid chromatography/single quadrupole mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2006 , 20, 2728-34	2.2	134
6	High-pressure liquid chromatographic-mass spectrometric determination of sorbic acid in urine: verification of formation of trans,trans-muconic acid. <i>Chemico-Biological Interactions</i> , 2005 , 153-154, 243-6	5	20
5	Determination of urinary S-phenylmercapturic acid, a specific metabolite of benzene, by liquid chromatography/single quadrupole mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2005 , 19, 1139-44	2.2	24
4	A field method for sampling toluene in end-exhaled air, as a biomarker of occupational exposure: correlation with other exposure indices. <i>Industrial Health</i> , 2004 , 42, 226-34	2.5	21
3	Urinary determination of N-acetyl- S-(N-methylcarbamoyl)cysteine and N-methylformamide in workers exposed to N, N-dimethylformamide. <i>International Archives of Occupational and Environmental Health</i> , 2002 , 75, 445-52	3.2	15

- | | | | |
|---|---|-----|----|
| 2 | Measurement of urinary N-acetyl-S-(N-methylcarbamoyl)cysteine by high-performance liquid chromatography with direct ultraviolet detection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2002 , 778, 231-6 | 3.2 | 7 |
| 1 | Trichloroethylene in urine as biological exposure index. <i>Industrial Health</i> , 2001 , 39, 225-30 | 2.5 | 15 |