

# Chen-Te Chou

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8081196/publications.pdf>

Version: 2024-02-01

16  
papers

149  
citations

1163117

8  
h-index

1199594

12  
g-index

16  
all docs

16  
docs citations

16  
times ranked

237  
citing authors

#	ARTICLE	IF	CITATIONS
1	Gallbladder torsion: preoperative diagnosis by MDCT. <i>Abdominal Imaging</i> , 2007, 32, 657-659.	2.0	25
2	Non-Invasive Evaluation of Hepatic Fibrosis: The Diagnostic Performance of Magnetic Resonance Elastography in Patients with Viral Hepatitis B or C. <i>PLoS ONE</i> , 2015, 10, e0140068.	2.5	24
3	Prospective Comparison of the Diagnostic Performance of Magnetic Resonance Elastography with Acoustic Radiation Force Impulse Elastography for Pre-operative Staging of Hepatic Fibrosis in Patients with Hepatocellular Carcinoma. <i>Ultrasound in Medicine and Biology</i> , 2017, 43, 2783-2790.	1.5	15
4	Thoroscopic Lobectomy Produces Long-Term Survival Similar to That with Open Lobectomy in Cases of Non-Small Cell Lung Carcinoma: A Propensity-Matched Analysis Using a Population-Based Cancer Registry. <i>Journal of Thoracic Oncology</i> , 2016, 11, 1326-1334.	1.1	12
5	Feasibility of measuring spleen stiffness with MR elastography and splenic volume to predict hepatic fibrosis stage. <i>PLoS ONE</i> , 2019, 14, e0217876.	2.5	12
6	Comparison of the efficacy of Gd-EOB-DTPA-enhanced magnetic resonance imaging and magnetic resonance elastography in the detection and staging of hepatic fibrosis. <i>Medicine (United States)</i> , 2017, 96, e8339.	1.0	10
7	Hemobilia after CT-guided radiofrequency ablation of liver tumors: frequency, risk factors, and clinical significance. <i>Abdominal Radiology</i> , 2019, 44, 337-345.	2.1	10
8	Diagnostic value of spleen stiffness by magnetic resonance elastography for prediction of esophageal varices in cirrhotic patients. <i>Abdominal Radiology</i> , 2021, 46, 526-533.	2.1	9
9	Gastric Subserosal Hematoma Developing from Focal Pancreatitis: A Case Report. <i>Kaohsiung Journal of Medical Sciences</i> , 2009, 25, 45-48.	1.9	8
10	Diagnostic Accuracy of Acoustic Radiation Force Impulse (ARFI) and Wisteria floribunda Agglutinin-Positive Mac-2-Binding Protein (WFA <sup>+</sup> -M2BP) in Patients with Chronic Liver Disease. <i>Medical Science Monitor</i> , 2019, 25, 7169-7174.	1.1	7
11	Dynamic-contrast-enhanced magnetic resonance imaging of cirrhotic liver parenchyma: A comparison between gadolinium-diethylenetriamine pentaacetic acid and gadolinium-ethoxybenzyl-diethylenetriamine pentaacetic acid. <i>Journal of the Chinese Medical Association</i> , 2015, 78, 666-672.	1.4	5
12	The short-term safety and efficacy of TANDEM microspheres of various sizes and doxorubicin loading concentrations for hepatocellular carcinoma treatment. <i>Scientific Reports</i> , 2021, 11, 12277.	3.3	5
13	Prognostic Role of Serum Wisteria Floribunda Agglutinin-Positive Mac-2 Binding Protein Level in Early Stage Hepatocellular Carcinoma. <i>Scientific Reports</i> , 2020, 10, 5651.	3.3	3
14	Using Glissonian sheath as an alternative way to prevent biliary stricture in living-donor liver transplantation. <i>Asian Journal of Surgery</i> , 2021, 44, 742-748.	0.4	3
15	Efficacy of gadoxetic acid-enhanced magnetic resonance cholangiography compared with T2-weighted magnetic resonance cholangiography in patients with liver cirrhosis. <i>Journal of the Chinese Medical Association</i> , 2015, 78, 726-731.	1.4	1
16	Bilateral retrocaval ureters with IVC duplication. <i>Abdominal Imaging</i> , 2006, 31, 596.	2.0	0