
Chen-Te Chou

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8081196/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Diagnostic value of spleen stiffness by magnetic resonance elastography for prediction of esophageal varices in cirrhotic patients. Abdominal Radiology, 2021, 46, 526-533.	2.1	9
2	Using Glissonian sheath as an alternative way to prevent biliary stricture in living-donor liver transplantation. Asian Journal of Surgery, 2021, 44, 742-748.	0.4	3
3	The short-term safety and efficacy of TANDEM microspheres of various sizes and doxorubicin loading concentrations for hepatocellular carcinoma treatment. Scientific Reports, 2021, 11, 12277.	3.3	5
4	Prognostic Role of Serum Wisteria Floribunda Agglutinin-Positive Mac-2 Binding Protein Level in Early Stage Hepatocellular Carcinoma. Scientific Reports, 2020, 10, 5651.	3.3	3
5	Hemobilia after CT-guided radiofrequency ablation of liver tumors: frequency, risk factors, and clinical significance. Abdominal Radiology, 2019, 44, 337-345.	2.1	10
6	Feasibility of measuring spleen stiffness with MR elastography and splenic volume to predict hepatic fibrosis stage. PLoS ONE, 2019, 14, e0217876.	2.5	12
7	Diagnostic Accuracy of Acoustic Radiation Force Impulse (ARFI) and Wisteria floribunda Agglutinin-Positive Mac-2-Binding Protein (WFAâº-M2BP) in Patients with Chronic Liver Disease. Medical Science Monitor, 2019, 25, 7169-7174.	1.1	7
8	Prospective Comparison of the Diagnostic Performance of Magnetic Resonance Elastography with Acoustic Radiation Force Impulse Elastography for Pre-operative Staging of Hepatic Fibrosis in Patients with Hepatocellular Carcinoma. Ultrasound in Medicine and Biology, 2017, 43, 2783-2790.	1.5	15
9	Comparison of the efficacy of Gd-EOB-DTPA-enhanced magnetic resonance imaging and magnetic resonance elastography in the detection and staging of hepatic fibrosis. Medicine (United States), 2017, 96, e8339.	1.0	10
10	Thoracoscopic Lobectomy Produces Long-Term Survival Similar to That with Open Lobectomy in Cases of Non–Small Cell Lung Carcinoma: AAPropensity-Matched Analysis Using a Population-Based Cancer Registry. Journal of Thoracic Oncology, 2016, 11, 1326-1334.	1.1	12
11	Non-Invasive Evaluation of Hepatic Fibrosis: The Diagnostic Performance of Magnetic Resonance Elastography in Patients with Viral Hepatitis B or C. PLoS ONE, 2015, 10, e0140068.	2.5	24
12	Efficacy of gadoxetic acid-enhanced magnetic resonance cholangiography compared with T2-weighted magnetic resonance cholangiography in patients with liver cirrhosis. Journal of the Chinese Medical Association, 2015, 78, 726-731.	1.4	1
13	Dynamic-contrast-enhanced magnetic resonance imaging of cirrhotic liver parenchyma: A comparison between gadolinium–diethylenetriamine pentaacetic acid and gadolinium–ethoxybenzyl–diethylenetriamine pentaacetic acid. Journal of the Chinese Medical Association. 2015. 78. 666-672.	1.4	5
14	Gastric Subserosal Hematoma Developing from Focal Pancreatitis: A Case Report. Kaohsiung Journal of Medical Sciences, 2009, 25, 45-48.	1.9	8
15	Gallbladder torsion: preoperative diagnosis by MDCT. Abdominal Imaging, 2007, 32, 657-659.	2.0	25
16	Bilateral retrocaval ureters with IVC duplication. Abdominal Imaging, 2006, 31, 596.	2.0	0