Ivonne M C M Rietjens

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8088148/ivonne-m-c-m-rietjens-publications-by-year.pdf
Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

411 papers	13,605 citations	60 h-index	93 g-index
424 ext. papers	15,175 ext. citations	4.7 avg, IF	6.35 L-index

#	Paper	IF	Citations
411	In Vitro Methodologies to Study the Role of Advanced Glycation End Products (AGEs) in Neurodegeneration.. *Nutrients*, **2022**, 14,	6.7	1
410	Developmental toxicity testing of unsubstituted and methylated 4- and 5-ring polycyclic aromatic hydrocarbons using the zebrafish embryotoxicity test.. *Toxicology in Vitro*, **2022**, 80, 105312	3.6	2
409	The effect of alkyl substitution on the oxidative metabolism and mutagenicity of phenanthrene.. *Archives of Toxicology*, **2022**, 96, 1109-1131	5.8	1
408	Inter-individual variation in chlorpyrifos toxicokinetics characterized by physiologically based kinetic (PBK) and Monte Carlo simulation comparing human liver microsome and Supersome cytochromes P450 (CYP)-specific kinetic data as model input.. *Archives of Toxicology*, **2022**, 96, 1387	5.8	
407	Identification of phosphodiesterase type-5 (PDE-5) inhibitors in herbal supplements using a tiered approach and associated consumer risk.. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, **2022**, 1-12	3.2	0
406	Letter to the Editor on Bil et al. 2021 "Risk Assessment of Per- and Polyfluoroalkyl Substance Mixtures: A Relative Potency Factor Approach".. *Environmental Toxicology and Chemistry*, **2022**, 41, 7-12	3.8	2
405	The Role of Kinetics as Key Determinant in Toxicity of Pyrrolizidine Alkaloids and Their N-Oxides. *Planta Medica*, **2021**,	3.1	2
404	In vitro and in silico study on consequences of combined exposure to the food-borne alkenylbenzenes estragole and safrole. *Toxicology in Vitro*, **2021**, 79, 105290	3.6	
403	Use of Physiologically Based Pharmacokinetic Modeling to Predict Human Gut Microbial Conversion of Daidzein to S-Equol. *Journal of Agricultural and Food Chemistry*, **2021**,	5.7	4
402	PBK Model-Based Prediction of Intestinal Microbial and Host Metabolism of Zearalenone and Consequences for its Estrogenicity. *Molecular Nutrition and Food Research*, **2021**, 65, e2100443	5.9	5
401	Physiologically based kinetic modelling based prediction of in vivo rat and human acetylcholinesterase (AChE) inhibition upon exposure to diazinon. *Archives of Toxicology*, **2021**, 95, 1573-1593	5.8	1
400	Prediction of dose-dependent in vivo acetylcholinesterase inhibition by profenofos in rats and humans using physiologically based kinetic (PBK) modeling-facilitated reverse dosimetry. *Archives of Toxicology*, **2021**, 95, 1287-1301	5.8	2
399	Assessment of the in vitro developmental toxicity of diethylstilbestrol and estradiol in the zebrafish embryotoxicity test. *Toxicology in Vitro*, **2021**, 72, 105088	3.6	3
398	An in vitro model for microbial fructoselysine degradation shows substantial interindividual differences in metabolic capacities of human fecal slurries. *Toxicology in Vitro*, **2021**, 72, 105078	3.6	4
397	Novel advances in biotransformation and bioactivation research - 2020 year in review. *Drug Metabolism Reviews*, **2021**, 53, 384-433	7	1
396	PDE-5 inhibitors in selected herbal supplements from the Ghanaian market for better erectile function as tested by a bioassay. *Toxicology in Vitro*, **2021**, 73, 105130	3.6	1
395	Incorporating renal excretion via the OCT2 transporter in physiologically based kinetic modelling to predict in vivo kinetics of mepiquat in rat. *Toxicology Letters*, **2021**, 343, 34-43	4.4	4

394	Next generation risk assessment of human exposure to anti-androgens using newly defined comparator compound values. *Toxicology in Vitro*, **2021**, 73, 105132	3.6	3
393	Estragole DNA adduct accumulation in human liver HepaRG cells upon repeated in vitro exposure. *Toxicology Letters*, **2021**, 337, 1-6	4.4	0
392	Novel testing strategy for prediction of rat biliary excretion of intravenously administered estradiol-17β-glucuronide. *Archives of Toxicology*, **2021**, 95, 91-102	5.8	
391	A chemical-specific adjustment factor for human interindividual differences in kinetics for glutamates (E620-625). *Food and Chemical Toxicology*, **2021**, 147, 111910	4.7	0
390	Interindividual Differences in Human In Vitro Intestinal Microbial Conversion of Green Tea (-)-Epigallocatechin-3-O-Gallate and Consequences for Activation of Nrf2 Mediated Gene Expression. *Molecular Nutrition and Food Research*, **2021**, 65, 2000934	5.9	5
389	Organophosphate and carbamate pesticide residues and accompanying risks in commonly consumed vegetables in Kenya. *Food Additives and Contaminants: Part B Surveillance*, **2021**, 14, 48-58	3.3	14
388	Risk characterisation of constituents present in jamu to promote its safe use. *Critical Reviews in Toxicology*, **2021**, 51, 183-191	5.7	1
387	Predicting the in vivo developmental toxicity of benzo[a]pyrene (BaP) in rats by an in vitro-in silico approach. *Archives of Toxicology*, **2021**, 95, 3323-3340	5.8	3
386	Interaction between food-borne mycotoxins and gut microbiota: A review. *Food Control*, **2021**, 126, 107998	6.2	3
385	FEMA GRAS assessment of natural flavor complexes: Eucalyptus oil and other cyclic ether-containing flavoring ingredients. *Food and Chemical Toxicology*, **2021**, 155, 112357	4.7	3
384	FEMA GRAS assessment of natural flavor complexes: Origanum oil, thyme oil and related phenol derivative-containing flavoring ingredients. *Food and Chemical Toxicology*, **2021**, 155, 112378	4.7	1
383	Developmental toxicity testing of the fume condensate extracts of bitumen and oxidized asphalt in a series of in vitro alternative assays. *Toxicology in Vitro*, **2021**, 75, 105195	3.6	1
382	Predicting the Acute Liver Toxicity of Aflatoxin B1 in Rats and Humans by an In Vitro-In Silico Testing Strategy. *Molecular Nutrition and Food Research*, **2020**, 64, e2000063	5.9	7
381	Integrating in vitro data and physiologically based kinetic modeling-facilitated reverse dosimetry to predict human cardiotoxicity of methadone. *Archives of Toxicology*, **2020**, 94, 2809-2827	5.8	9
380	Monocrotaline-induced liver toxicity in rat predicted by a combined in vitro physiologically based kinetic modeling approach. *Archives of Toxicology*, **2020**, 94, 3281-3295	5.8	6
379	Mode of action-based risk assessment of genotoxic carcinogens. *Archives of Toxicology*, **2020**, 94, 1787-1877	5.8	46
378	Cellular levels and molecular dynamics simulations of estragole DNA adducts point at inefficient repair resulting from limited distortion of the double-stranded DNA helix. *Archives of Toxicology*, **2020**, 94, 1349-1365	5.8	4
377	The safety evaluation of food flavoring substances: the role of genotoxicity studies. *Critical Reviews in Toxicology*, **2020**, 50, 1-27	5.7	12

376	Induction of peroxisome proliferator activated receptor (PPAR) mediated gene expression and inhibition of induced nitric oxide production by Maerua subcordata (Gilg) DeWolf. *BMC Complementary Medicine and Therapies*, **2020**, 20, 80	2.9	3
375	Detection of pyrrolizidine alkaloids in jamu available on the Indonesian market and accompanying safety assessment for human consumption. *Food and Chemical Toxicology*, **2020**, 138, 111230	4.7	6
374	Novel advances in biotransformation and bioactivation research-2019 year in review. *Drug Metabolism Reviews*, **2020**, 52, 333-365	7	1
373	Use of Physiologically Based Kinetic Modeling to Predict Rat Gut Microbial Metabolism of the Isoflavone Daidzein to S-Equol and Its Consequences for ER Activation. *Molecular Nutrition and Food Research*, **2020**, 64, e1900912	5.9	12
372	Estrogen receptor alpha (ER)-mediated coregulator binding and gene expression discriminates the toxic ER agonist diethylstilbestrol (DES) from the endogenous ER agonist 17 estradiol (E2). *Cell Biology and Toxicology*, **2020**, 36, 417-435	7.4	6
371	The use of adverse outcome pathways in the safety evaluation of food additives. *Archives of Toxicology*, **2020**, 94, 959-966	5.8	11
370	Development of a Generic Physiologically Based Kinetic Model to Predict In Vivo Uterotrophic Responses Induced by Estrogenic Chemicals in Rats Based on In Vitro Bioassays. *Toxicological Sciences*, **2020**, 173, 19-31	4.4	4
369	Evaluation of in vitro models of stem cell-derived cardiomyocytes to screen for potential cardiotoxicity of chemicals. *Toxicology in Vitro*, **2020**, 67, 104891	3.6	5
368	Pyrrolizidine alkaloids in food and phytomedicine: Occurrence, exposure, toxicity, mechanisms, and risk assessment - A review. *Food and Chemical Toxicology*, **2020**, 136, 111107	4.7	36
367	Combining In Vitro Data and Physiologically Based Kinetic Modeling Facilitates Reverse Dosimetry to Define In Vivo Dose-Response Curves for Bixin- and Crocetin-Induced Activation of PPAR in Humans. *Molecular Nutrition and Food Research*, **2020**, 64, e1900880	5.9	4
366	The role of metabolism in the developmental toxicity of polycyclic aromatic hydrocarbon-containing extracts of petroleum substances. *Journal of Applied Toxicology*, **2020**, 40, 330-341	4.1	8
365	FEMA GRAS assessment of natural flavor complexes: Cinnamomum and Myroxylon-derived flavoring ingredients. *Food and Chemical Toxicology*, **2020**, 135, 110949	4.7	9
364	In vitro metabolism of naphthalene and its alkylated congeners by human and rat liver microsomes via alkyl side chain or aromatic oxidation. *Chemico-Biological Interactions*, **2020**, 315, 108905	5	8
363	FEMA GRAS assessment of natural flavor complexes: Lavender, Guaiac Coriander-derived and related flavoring ingredients. *Food and Chemical Toxicology*, **2020**, 145, 111584	4.7	1
362	Interindividual Differences in Human Intestinal Microbial Conversion of (-)-Epicatechin to Bioactive Phenolic Compounds. *Journal of Agricultural and Food Chemistry*, **2020**,	5.7	18
361	An model to quantify interspecies differences in kinetics for intestinal microbial bioactivation and detoxification of zearalenone. *Toxicology Reports*, **2020**, 7, 938-946	4.8	6
360	FEMA GRAS assessment of natural flavor complexes: Clove, cinnamon leaf and West Indian bay leaf-derived flavoring ingredients. *Food and Chemical Toxicology*, **2020**, 145, 111585	4.7	11
359	Defining in vivo dose-response curves for kidney DNA adduct formation of aristolochic acid I in rat, mouse and human by an in vitro and physiologically based kinetic modeling approach. *Journal of Applied Toxicology*, **2020**, 40, 1647-1660	4.1	1

358	Molecular Dynamics and Quantification of Safrole DNA Adducts Reveal DNA Adduct Persistence Due to Limited DNA Distortion Resulting in Inefficient Repair. *Chemical Research in Toxicology*, **2020** , 33, 2298-2309	4	4
357	FEMA GRAS assessment of natural flavor complexes: Mint, buchu, dill and caraway derived flavoring ingredients. *Food and Chemical Toxicology*, **2020**, 135, 110870	4.7	12
356	In vitro prenatal developmental toxicity induced by some petroleum substances is mediated by their 3- to 7-ring PAH constituent with a potential role for the aryl hydrocarbon receptor (AhR). *Toxicology Letters*, **2019**, 315, 64-76	4.4	15
355	Integrating physiologically based kinetic (PBK) and Monte Carlo modelling to predict inter-individual and inter-ethnic variation in bioactivation and liver toxicity of lasiocarpine. *Archives of Toxicology*, **2019**, 93, 2943-2960	5.8	5
354	Use of an in vitro-in silico testing strategy to predict inter-species and inter-ethnic human differences in liver toxicity of the pyrrolizidine alkaloids lasiocarpine and riddelliine. *Archives of Toxicology*, **2019**, 93, 801-818	5.8	22
353	Levels of methyleugenol and eugenol in instant herbal beverages available on the Indonesian market and related risk assessment. *Food and Chemical Toxicology*, **2019**, 125, 467-478	4.7	19
352	The in vivo developmental toxicity of diethylstilbestrol (DES) in rat evaluated by an alternative testing strategy. *Archives of Toxicology*, **2019**, 93, 2021-2033	5.8	8
351	Physiologically based kinetic modelling-facilitated reverse dosimetry to predict in vivo red blood cell acetylcholinesterase inhibition following exposure to chlorpyrifos in the Caucasian and Chinese population. *Toxicological Sciences*, **2019**,	4.4	8
350	Role of toxicokinetics and alternative testing strategies in pyrrolizidine alkaloid toxicity and risk assessment; state-of-the-art and future perspectives. *Food and Chemical Toxicology*, **2019**, 131, 110572	4.7	8
349	Hazard assessment of Maerua subcordata (Gilg) DeWolf. for selected endpoints using a battery of in vitro tests. *Journal of Ethnopharmacology*, **2019**, 241, 111978	5	1
348	Biotransformation and bioactivation reactions - 2018 literature highlights. *Drug Metabolism Reviews*, **2019**, 51, 121-161	7	3
347	Induction of EpRE-mediated gene expression by a series of mediterranean botanicals and their constituents. *Journal of Ethnopharmacology*, **2019**, 240, 111940	5	1
346	Risk assessment of intake of pyrrolizidine alkaloids from herbal teas and medicines following realistic exposure scenarios. *Food and Chemical Toxicology*, **2019**, 130, 142-153	4.7	15
345	Selecting the dose metric in reverse dosimetry based QIVIVE : Reply to ℞Comment on ℞Use of an in vitro-in silico testing strategy to predict inter-species and inter-ethnic human differences in liver toxicity of the pyrrolizidine alkaloids lasiocarpine and riddelliine℞ by Ning et al., Arch Toxicol doi: https://doi.org/10.1007/s00204-019-02397-7℞ Arch Toxicol doi: https://doi.org/10.1007/s00204	5.8	5
344	Effects of Maerua subcordata (Gilg) DeWolf on electrophile-responsive element (EpRE)-mediated gene expression in vitro. *PLoS ONE*, **2019**, 14, e0215155	3.7	2
343	Novel approaches to derive points of departure for food chemical risk assessment. *Current Opinion in Food Science*, **2019**, 27, 139-144	9.8	3
342	Characterizing the coverage of critical effects relevant in the safety evaluation of food additives by AOPs. *Archives of Toxicology*, **2019**, 93, 2115-2125	5.8	10
341	Prediction of in vivo genotoxicity of lasiocarpine and riddelliine in rat liver using a combined in vitro-physiologically based kinetic modelling-facilitated reverse dosimetry approach. *Archives of Toxicology*, **2019**, 93, 2385-2395	5.8	10

340	Aflatoxin B1 in nixtamalized maize in Mexico; occurrence and accompanying risk assessment. *Toxicology Reports*, **2019**, 6, 1135-1142	4.8	11
339	Risk assessment of herbal supplements containing ingredients that are genotoxic and carcinogenic. *Critical Reviews in Toxicology*, **2019**, 49, 567-579	5.7	6
338	FEMA GRAS assessment of natural flavor complexes: Citrus-derived flavoring ingredients. *Food and Chemical Toxicology*, **2019**, 124, 192-218	4.7	18
337	The effects of all-trans retinoic acid on estrogen receptor signaling in the estrogen-sensitive MCF/BUS subline. *Journal of Receptor and Signal Transduction Research*, **2018**, 38, 112-121	2.6	4
336	The safety evaluation of food flavouring substances: the role of metabolic studies. *Toxicology Research*, **2018**, 7, 618-646	2.6	12
335	Updated procedure for the safety evaluation of natural flavor complexes used as ingredients in food. *Food and Chemical Toxicology*, **2018**, 113, 171-178	4.7	17
334	Use of physiologically based kinetic modelling-facilitated reverse dosimetry to convert in vitro cytotoxicity data to predicted in vivo liver toxicity of lasiocarpine and riddelliine in rat. *Food and Chemical Toxicology*, **2018**, 116, 216-226	4.7	24
333	Risk assessment of aflatoxin B1 exposure from maize and peanut consumption in Indonesia using the margin of exposure and liver cancer risk estimation approaches. *Food and Chemical Toxicology*, **2018**, 113, 134-144	4.7	34
332	A comparison of the embryonic stem cell test and whole embryo culture assay combined with the BeWo placental passage model for predicting the embryotoxicity of azoles. *Toxicology Letters*, **2018**, 286, 10-21	4.4	18
331	Exposure assessment of process-related contaminants in food by biomarker monitoring. *Archives of Toxicology*, **2018**, 92, 15-40	5.8	27
330	Effects of Systematic Variation in Size and Surface Coating of Silver Nanoparticles on Their In Vitro Toxicity to Macrophage RAW 264.7 Cells. *Toxicological Sciences*, **2018**, 162, 79-88	4.4	26
329	Natural occurrence of genotoxic and carcinogenic alkenylbenzenes in Indonesian jamu and evaluation of consumer risks. *Food and Chemical Toxicology*, **2018**, 118, 53-67	4.7	13
328	Risk assessment of genotoxic and carcinogenic alkenylbenzenes in botanical containing products present on the Chinese market. *Food and Chemical Toxicology*, **2018**, 115, 344-357	4.7	10
327	Perceived versus real toxicological safety of pangasius catfish: a review modifying market perspectives. *Reviews in Aquaculture*, **2018**, 10, 123-134	8.9	12
326	Towards a generic physiologically based kinetic model to predict in vivo uterotrophic responses in rats by reverse dosimetry of in vitro estrogenicity data. *Archives of Toxicology*, **2018**, 92, 1075-1088	5.8	16
325	Risk Assessment Paradigm for Glutamate. *Annals of Nutrition and Metabolism*, **2018**, 73 Suppl 5, 53-64	4.5	11
324	Use of proteomics to detect sex-related differences in effects of toxicants: implications for using proteomics in toxicology. *Critical Reviews in Toxicology*, **2018**, 48, 666-681	5.7	5
323	Soy supplementation: Impact on gene expression in different tissues of ovariectomized rats and evaluation of the rat model to predict (post)menopausal health effect. *Toxicology Reports*, **2018**, 5, 1087-1097	4.8	2

322	Impact of nanoparticle surface functionalization on the protein corona and cellular adhesion, uptake and transport. *Journal of Nanobiotechnology*, **2018**, 16, 70	9.4	35
321	The Role of Endocrine and Dioxin-Like Activity of Extracts of Petroleum Substances in Developmental Toxicity as Detected in a Panel of CALUX Reporter Gene Assays. *Toxicological Sciences*, **2018**, 164, 576-591	4.4	20
320	Biotransformation and bioactivation reactions - 2017 literature highlights. *Drug Metabolism Reviews*, **2018**, 50, 221-255	7	6
319	Determination and risk assessment of naturally occurring genotoxic and carcinogenic alkenylbenzenes in basil-containing sauce of pesto. *Toxicology Reports*, **2017**, 4, 1-8	4.8	16
318	Physiologically based kinetic modeling of hesperidin metabolism and its use to predict in vivo effective doses in humans. *Molecular Nutrition and Food Research*, **2017**, 61, 1600894	5.9	16
317	Embryotoxic and pharmacologic potency ranking of six azoles in the rat whole embryo culture by morphological and transcriptomic analysis. *Toxicology and Applied Pharmacology*, **2017**, 322, 15-26	4.6	14
316	Development of a Combined In Vitro Physiologically Based Kinetic (PBK) and Monte Carlo Modelling Approach to Predict Interindividual Human Variation in Phenol-Induced Developmental Toxicity. *Toxicological Sciences*, **2017**, 157, 365-376	4.4	12
315	Marine biotoxins and associated outbreaks following seafood consumption: Prevention and surveillance in the 21st century. *Global Food Security*, **2017**, 15, 11-21	8.3	39
314	Biotransformation and bioactivation reactions - 2016 literature highlights. *Drug Metabolism Reviews*, **2017**, 49, 285-317	7	5
313	Use of physiologically based kinetic modeling-facilitated reverse dosimetry of in vitro toxicity data for prediction of in vivo developmental toxicity of tebuconazole in rats. *Toxicology Letters*, **2017**, 266, 85-93	4.4	22
312	In vitro bioassays to evaluate beneficial and adverse health effects of botanicals: promises and pitfalls. *Drug Discovery Today*, **2017**, 22, 1187-1200	8.8	9
311	In vitro-in silico-based analysis of the dose-dependent in vivo oestrogenicity of the soy phytoestrogen genistein in humans. *British Journal of Pharmacology*, **2017**, 174, 2739-2757	8.6	15
310	Determination and risk assessment of naturally occurring genotoxic and carcinogenic alkenylbenzenes in nutmeg-based plant food supplements. *Journal of Applied Toxicology*, **2017**, 37, 1254-1264	4.1	14
309	Study on inter-ethnic human differences in bioactivation and detoxification of estragole using physiologically based kinetic modeling. *Archives of Toxicology*, **2017**, 91, 3093-3108	5.8	5
308	Risk assessment for pyrrolizidine alkaloids detected in (herbal) teas and plant food supplements. *Regulatory Toxicology and Pharmacology*, **2017**, 86, 292-302	3.4	37
307	A transcriptomic approach for evaluating the relative potency and mechanism of action of azoles in the rat Whole Embryo Culture. *Toxicology*, **2017**, 392, 96-105	4.4	7
306	The Regulatory Framework Across International Jurisdictions for Risks Associated with Consumption of Botanical Food Supplements. *Comprehensive Reviews in Food Science and Food Safety*, **2017**, 16, 821-834	16.4	29
305	Risk assessment of combined exposure to alkenylbenzenes through consumption of plant food supplements containing parsley and dill. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, **2017**, 34, 2201-2211	3.2	10

304	Characterization of the differential coregulator binding signatures of the Retinoic Acid Receptor subtypes upon (ant)agonist action. *Biochimica Et Biophysica Acta - Proteins and Proteomics*, **2017**, 1865, 1195-1206	4	4
303	Prenatal developmental toxicity testing of petroleum substances: Application of the mouse embryonic stem cell test (EST) to compare in vitro potencies with potencies observed in vivo. *Toxicology in Vitro*, **2017**, 44, 303-312	3.6	18
302	Undesired Plant-Derived Components in Food **2017**, 379-424		6
301	Integrating in vitro data and physiologically based kinetic (PBK) modelling to assess the in vivo potential developmental toxicity of a series of phenols. *Archives of Toxicology*, **2017**, 91, 2119-2133	5.8	28
300	Safety evaluation of substituted thiophenes used as flavoring ingredients. *Food and Chemical Toxicology*, **2017**, 99, 40-59	4.7	12
299	Use of Physiologically Based Kinetic Modeling-Based Reverse Dosimetry to Predict in Vivo Toxicity from in Vitro Data. *Chemical Research in Toxicology*, **2017**, 30, 114-125	4	55
298	The potential health effects of dietary phytoestrogens. *British Journal of Pharmacology*, **2017**, 174, 1263-1280	8	226
297	Physiologically based kinetic modeling of the bioactivation of myristicin. *Archives of Toxicology*, **2017**, 91, 713-734	5.8	22
296	Risks to human and animal health related to the presence of deoxynivalenol and its acetylated and modified forms in food and feed. *EFSA Journal*, **2017**, 15, e04718	2.3	132
295	Risk assessment of plant food supplements and other herbal products containing aristolochic acids using the margin of exposure (MOE) approach. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, **2017**, 34, 135-144	3.2	10
294	Level of Alkenylbenzenes in Parsley and Dill Based Teas and Associated Risk Assessment Using the Margin of Exposure Approach. *Journal of Agricultural and Food Chemistry*, **2016**, 64, 8640-8646	5.7	15
293	Biotransformation and bioactivation reactions - 2015 literature highlights. *Drug Metabolism Reviews*, **2016**, 48, 113-38	7	10
292	Evaluation of Interindividual Human Variation in Bioactivation and DNA Adduct Formation of Estragole in Liver Predicted by Physiologically Based Kinetic/Dynamic and Monte Carlo Modeling. *Chemical Research in Toxicology*, **2016**, 29, 659-68	4	15
291	Effect of Glucuronidation on the Potential of Kaempferol to Inhibit Serine/Threonine Protein Kinases. *Journal of Agricultural and Food Chemistry*, **2016**, 64, 1256-63	5.7	6
290	Mode of action based risk assessment of the botanical food-borne alkenylbenzene apiol from parsley using physiologically based kinetic (PBK) modelling and read-across from safrole. *Food and Chemical Toxicology*, **2016**, 89, 138-50	4.7	18
289	Extended evaluation on the ES-D3 cell differentiation assay combined with the BeWo transport model, to predict relative developmental toxicity of triazole compounds. *Archives of Toxicology*, **2016**, 90, 1225-37	5.8	13
288	FEMA expert panel review of p-mentha-1,8-dien-7-al genotoxicity testing results. *Food and Chemical Toxicology*, **2016**, 98, 201-209	4.7	6
287	Predicting points of departure for risk assessment based on in vitro cytotoxicity data and physiologically based kinetic (PBK) modeling: The case of kidney toxicity induced by aristolochic acid I. *Food and Chemical Toxicology*, **2016**, 92, 104-16	4.7	28

286	Progenitor-derived hepatocyte-like (B-13/H) cells metabolise 1Phydroxyestragole to a genotoxic species via a SULT2B1-dependent mechanism. *Toxicology Letters*, **2016**, 243, 98-110	4.4	8
285	Flusilazole induces spatio-temporal expression patterns of retinoic acid-, differentiation- and sterol biosynthesis-related genes in the rat Whole Embryo Culture. *Reproductive Toxicology*, **2016**, 64, 77-85	3.4	11
284	Combining an in vitro reporter gene assay with metabolomics to identify tomato phytochemicals responsible for inducing electrophile-responsive element (EpRE)-mediated gene transcription. *Metabolomics*, **2015**, 11, 302-311	4.7	1
283	Plasma bioavailability and changes in PBMC gene expression after treatment of ovariectomized rats with a commercial soy supplement. *Toxicology Reports*, **2015**, 2, 308-321	4.8	2
282	In vitro gastrointestinal digestion increases the translocation of polystyrene nanoparticles in an in vitro intestinal co-culture model. *Nanotoxicology*, **2015**, 9, 886-94	5.3	61
281	Quercetin tests negative for genotoxicity in transcriptome analyses of liver and small intestine of mice. *Food and Chemical Toxicology*, **2015**, 81, 34-39	4.7	13
280	Progress and future of in vitro models to study translocation of nanoparticles. *Archives of Toxicology*, **2015**, 89, 1469-95	5.8	95
279	The effect of quercetin and kaempferol aglycones and glucuronides on peroxisome proliferator-activated receptor-gamma (PPAR-) *Food and Function*, **2015**, 6, 1098-107	6.1	20
278	Bioavailability and biodistribution of differently charged polystyrene nanoparticles upon oral exposure in rats. *Journal of Nanoparticle Research*, **2015**, 17, 231	2.3	68
277	Exploration of new functional endpoints in neuro-2a cells for the detection of the marine biotoxins saxitoxin, palytoxin and tetrodotoxin. *Toxicology in Vitro*, **2015**, 30, 341-7	3.6	12
276	The effect of glucuronidation on isoflavone induced estrogen receptor (ER) and ER mediated coregulator interactions. *Journal of Steroid Biochemistry and Molecular Biology*, **2015**, 154, 245-53	5.1	17
275	Matrix-derived combination effects influencing absorption, distribution, metabolism and excretion (ADME) of food-borne toxic compounds: implications for risk assessment. *Toxicology Research*, **2015**, 4, 23-35	2.6	13
274	In vitro detection of cardiotoxins or neurotoxins affecting ion channels or pumps using beating cardiomyocytes as alternative for animal testing. *Toxicology in Vitro*, **2015**, 29, 281-8	3.6	10
273	Use of the ES-D3 cell differentiation assay, combined with the BeWo transport model, to predict relative in vivo developmental toxicity of antifungal compounds. *Toxicology in Vitro*, **2015**, 29, 320-8	3.6	23
272	Translocation of differently sized and charged polystyrene nanoparticles in in vitro intestinal cell models of increasing complexity. *Nanotoxicology*, **2015**, 9, 453-61	5.3	83
271	Prediction of in vivo developmental toxicity of all-trans-retinoic acid based on in vitro toxicity data and in silico physiologically based kinetic modeling. *Archives of Toxicology*, **2015**, 89, 1135-48	5.8	48
270	Direct comparison of metabolic health effects of the flavonoids quercetin, hesperetin, epicatechin, apigenin and anthocyanins in high-fat-diet-fed mice. *Genes and Nutrition*, **2015**, 10, 469	4.3	62
269	Translocation of positively and negatively charged polystyrene nanoparticles in an in vitro placental model. *Toxicology in Vitro*, **2015**, 29, 1701-10	3.6	30

268	Use of physiologically based kinetic (PBK) modeling to study interindividual human variation and species differences in plasma concentrations of quercetin and its metabolites. *Biochemical Pharmacology*, **2015**, 98, 690-702	6	12
267	Evaluation of the interindividual human variation in bioactivation of methyleugenol using physiologically based kinetic modeling and Monte Carlo simulations. *Toxicology and Applied Pharmacology*, **2015**, 283, 117-26	4.6	10
266	Effects of silver nanoparticles (NM-300K) on Lumbricus rubellus earthworms and particle characterization in relevant test matrices including soil. *Environmental Toxicology and Chemistry*, **2014**, 33, 743-52	3.8	74
265	Conversion of major soy isoflavone glucosides and aglycones in in vitro intestinal models. *Molecular Nutrition and Food Research*, **2014**, 58, 503-15	5.9	37
264	Active pharmaceutical ingredients detected in herbal food supplements for weight loss sampled on the Dutch market. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, **2014**, 31, 1783-93	3.2	20
263	Role of membrane disturbance and oxidative stress in the mode of action underlying the toxicity of differently charged polystyrene nanoparticles. *RSC Advances*, **2014**, 4, 19321-19330	3.7	39
262	Impact of structural and metabolic variations on the toxicity and carcinogenicity of hydroxy- and alkoxy-substituted allyl- and propenylbenzenes. *Chemical Research in Toxicology*, **2014**, 27, 1092-103	4	36
261	Predicting individual responses to pravastatin using a physiologically based kinetic model for plasma cholesterol concentrations. *Journal of Pharmacokinetics and Pharmacodynamics*, **2014**, 41, 351-62	2.7	5
260	The natural basil flavonoid nevadensin protects against a methyleugenol-induced marker of hepatocarcinogenicity in male F344 rat. *Food and Chemical Toxicology*, **2014**, 74, 28-34	4.7	18
259	P-gp efflux pump inhibition potential of common environmental contaminants determined in vitro. *Environmental Toxicology and Chemistry*, **2014**, 33, 804-13	3.8	12
258	Identification of coregulators influenced by estrogen receptor subtype specific binding of the ER antagonists 4-hydroxytamoxifen and fulvestrant. *Chemico-Biological Interactions*, **2014**, 220, 222-30	5	8
257	Extending an in vitro panel for estrogenicity testing: the added value of bioassays for measuring antiandrogenic activities and effects on steroidogenesis. *Toxicological Sciences*, **2014**, 141, 78-89	4.4	21
256	Chemical analysis of estragole in fennel based teas and associated safety assessment using the Margin of Exposure (MOE) approach. *Food and Chemical Toxicology*, **2014**, 65, 147-54	4.7	21
255	A physiologically based kinetic (PBK) model describing plasma concentrations of quercetin and its metabolites in rats. *Biochemical Pharmacology*, **2014**, 89, 287-99	6	13
254	Cell proliferation and modulation of interaction of estrogen receptors with coregulators induced by ER and ER agonists. *Journal of Steroid Biochemistry and Molecular Biology*, **2014**, 143, 376-85	5.1	13
253	Detection of marine neurotoxins in food safety testing using a multielectrode array. *Molecular Nutrition and Food Research*, **2014**, 58, 2369-78	5.9	42
252	In vitro nanoparticle toxicity to rat alveolar cells and coelomocytes from the earthworm Lumbricus rubellus. *Nanotoxicology*, **2014**, 8, 28-37	5.3	25
251	Formation of genotoxic compounds by medium pressure ultraviolet treatment of nitrate-rich water. *Desalination and Water Treatment*, **2014**, 52, 6275-6281		16

250	Towards an integrated in vitro strategy for estrogenicity testing. *Journal of Applied Toxicology*, **2014**, 34, 1031-40	4.1	14
249	GRASr2 evaluation of aliphatic acyclic and alicyclic terpenoid tertiary alcohols and structurally related substances used as flavoring ingredients. *Journal of Food Science*, **2014**, 79, R428-41	3.4	13
248	Marine neurotoxins: state of the art, bottlenecks, and perspectives for mode of action based methods of detection in seafood. *Molecular Nutrition and Food Research*, **2014**, 58, 87-100	5.9	31
247	Malabaricone C-containing mace extract inhibits safrole bioactivation and DNA adduct formation both in vitro and in vivo. *Food and Chemical Toxicology*, **2014**, 66, 373-84	4.7	11
246	Developmental toxicity of thyroid-active compounds in a zebrafish embryotoxicity test. *ALTEX: Alternatives To Animal Experimentation*, **2014**, 31, 303-17	4.3	25
245	Identification of thyroid hormone receptor active compounds using a quantitative high-throughput screening platform. *Current Chemical Genomics and Translational Medicine*, **2014**, 8, 36-46		18
244	A physiologically based in silico model for trans-2-hexenal detoxification and DNA adduct formation in human including interindividual variation indicates efficient detoxification and a negligible genotoxicity risk. *Archives of Toxicology*, **2013**, 87, 1725-37	5.8	13
243	Assessment of an in vitro transport model using BeWo b30 cells to predict placental transfer of compounds. *Archives of Toxicology*, **2013**, 87, 1661-9	5.8	52
242	Correlation between activation of PPARγ and resistin downregulation in a mouse adipocyte cell line by a series of thiazolidinediones. *Toxicology in Vitro*, **2013**, 27, 1425-32	3.6	0
241	Consumer and farmer safety evaluation of application of botanical pesticides in black pepper crop protection. *Food and Chemical Toxicology*, **2013**, 56, 483-90	4.7	18
240	Role of surface charge in bioavailability and biodistribution of tri-block copolymer nanoparticles in rats after oral exposure. *Toxicology Research*, **2013**, 2, 187	2.6	4
239	Induction of peroxisome proliferator-activated receptor γ (PPARγ)-mediated gene expression by tomato (Solanum lycopersicum L.) extracts. *Journal of Agricultural and Food Chemistry*, **2013**, 61, 3419-27	5.7	29
238	Mechanisms underlying the dualistic mode of action of major soy isoflavones in relation to cell proliferation and cancer risks. *Molecular Nutrition and Food Research*, **2013**, 57, 100-13	5.9	81
237	Matrix-derived combination effect and risk assessment for estragole from basil-containing plant food supplements (PFS). *Food and Chemical Toxicology*, **2013**, 62, 32-40	4.7	13
236	Inhibition of methyleugenol bioactivation by the herb-based constituent nevadensin and prediction of possible in vivo consequences using physiologically based kinetic modeling. *Food and Chemical Toxicology*, **2013**, 59, 564-71	4.7	18
235	A low-density DNA microchip for the detection of (anti-)estrogenic compounds and their relative potencies. *Analytical Biochemistry*, **2013**, 435, 83-92	3.1	3
234	Human T47D-ERβ breast cancer cells with tetracycline-dependent ERβ expression reflect ERα/ERβ ratios in rat and human breast tissue. *Toxicology in Vitro*, **2013**, 27, 1753-61	3.6	6
233	Surface charge-specific cytotoxicity and cellular uptake of tri-block copolymer nanoparticles. *Nanotoxicology*, **2013**, 7, 71-84	5.3	51

232	Behaviour of silver nanoparticles and silver ions in an in vitro human gastrointestinal digestion model. *Nanotoxicology*, **2013**, 7, 1198-210	5.3	165
231	Cytotoxicity of surface-functionalized silicon and germanium nanoparticles: the dominant role of surface charges. *Nanoscale*, **2013**, 5, 4870-83	7.7	141
230	Surface charge-specific interactions between polymer nanoparticles and ABC transporters in Caco-2 cells. *Journal of Nanoparticle Research*, **2013**, 15, 1	2.3	13
229	Combining in vitro embryotoxicity data with physiologically based kinetic (PBK) modelling to define in vivo dose-response curves for developmental toxicity of phenol in rat and human. *Archives of Toxicology*, **2013**, 87, 1709-23	5.8	49
228	Robust array-based coregulator binding assay predicting ER agonist potency and generating binding profiles reflecting ligand structure. *Chemical Research in Toxicology*, **2013**, 26, 336-46	4	24
227	In vivo validation and physiologically based biokinetic modeling of the inhibition of SULT-mediated estragole DNA adduct formation in the liver of male Sprague-Dawley rats by the basil flavonoid nevadensin. *Molecular Nutrition and Food Research*, **2013**, 57, 1969-78	5.9	20
226	Quercetin induces hepatic lipid omega-oxidation and lowers serum lipid levels in mice. *PLoS ONE*, **2013**, 8, e51588	3.7	55
225	A 155-plex high-throughput in vitro coregulator binding assay for (anti-)estrogenicity testing evaluated with 23 reference compounds. *ALTEX: Alternatives To Animal Experimentation*, **2013**, 30, 145-57	4.3	22
224	Physiologically based kinetic modeling of bioactivation and detoxification of the alkenylbenzene methyleugenol in human as compared with rat. *Toxicology and Applied Pharmacology*, **2012**, 260, 271-84	4.6	32
223	Alkaloids in the human food chain--natural occurrence and possible adverse effects. *Molecular Nutrition and Food Research*, **2012**, 56, 30-52	5.9	104
222	Interference of flavonoids with enzymatic assays for the determination of free fatty acid and triglyceride levels. *Analytical and Bioanalytical Chemistry*, **2012**, 402, 1389-92	4.4	12
221	A physiologically based in silico model for trans-2-hexenal detoxification and DNA adduct formation in rat. *Chemical Research in Toxicology*, **2012**, 25, 2630-41	4	11
220	Physiologically based kinetic models for the alkenylbenzene elemicin in rat and human and possible implications for risk assessment. *Chemical Research in Toxicology*, **2012**, 25, 2352-67	4	28
219	Inhibition of cellular efflux pumps involved in multi xenobiotic resistance (MXR) in echinoid larvae as a possible mode of action for increased ecotoxicological risk of mixtures. *Ecotoxicology*, **2012**, 21, 2276-87	2.9	32
218	Bovine liver slices: A multifunctional in vitro model to study the prohormone dehydroepiandrosterone (DHEA). *Toxicology in Vitro*, **2012**, 26, 1014-21	3.6	11
217	Relative embryotoxic potency of p-substituted phenols in the embryonic stem cell test (EST) and comparison to their toxic potency in vivo and in the whole embryo culture (WEC) assay. *Toxicology Letters*, **2012**, 213, 235-42	4.4	17
216	Proliferation assays for estrogenicity testing with high predictive value for the in vivo uterotrophic effect. *Journal of Steroid Biochemistry and Molecular Biology*, **2012**, 128, 98-106	5.1	16
215	Induction of electrophile-responsive element (EpRE)-mediated gene expression by tomato extracts in vitro. *Food Chemistry*, **2012**, 135, 1166-72	8.5	7

214	Toward in vitro biomarkers for developmental toxicity and their extrapolation to the in vivo situation. *Expert Opinion on Drug Metabolism and Toxicology*, **2012**, 8, 11-27	5.5	25
213	Refined hazard characterization of 3-MCPD using benchmark dose modeling. *European Journal of Lipid Science and Technology*, **2012**, 114, 1140-1147	3	4
212	Cytotoxicity and cellular uptake of tri-block copolymer nanoparticles with different size and surface characteristics. *Particle and Fibre Toxicology*, **2012**, 9, 11	8.4	55
211	Safety assessment of smoke flavouring primary products by the European Food Safety Authority. *Trends in Food Science and Technology*, **2012**, 27, 97-108	15.3	12
210	A state-of-the-art overview of the effect of metabolic conjugation on the biological activity of flavonoids. *Food and Function*, **2012**, 3, 1008-18	6.1	47
209	Physiologically based biokinetic (PBBK) modeling of safrole bioactivation and detoxification in humans as compared with rats. *Toxicological Sciences*, **2012**, 128, 301-16	4.4	25
208	Matrix modulation of the bioactivation of estragole by constituents of different alkenylbenzene-containing herbs and spices and physiologically based biokinetic modeling of possible in vivo effects. *Toxicological Sciences*, **2012**, 129, 174-87	4.4	17
207	A physiologically based in silico kinetic model predicting plasma cholesterol concentrations in humans. *Journal of Lipid Research*, **2012**, 53, 2734-46	6.3	23
206	In vivo validation of DNA adduct formation by estragole in rats predicted by physiologically based biodynamic modelling. *Mutagenesis*, **2012**, 27, 653-63	2.8	27
205	Update of risk assessments of main marine biotoxins in the European Union. *Toxicon*, **2011**, 58, 336-54	2.8	57
204	The FEMA GRAS assessment of aliphatic and aromatic terpene hydrocarbons used as flavor ingredients. *Food and Chemical Toxicology*, **2011**, 49, 2471-94	4.7	47
203	A physiologically-based kinetic model for the prediction of plasma cholesterol concentrations in the mouse. *Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids*, **2011**, 1811, 333-42	5	6
202	Physiologically based biokinetic model of bioactivation and detoxification of the alkenylbenzene methyleugenol in rat. *Toxicology in Vitro*, **2011**, 25, 267-85	3.6	33
201	Relative developmental toxicity potencies of retinoids in the embryonic stem cell test compared with their relative potencies in in vivo and two other in vitro assays for developmental toxicity. *Toxicology Letters*, **2011**, 203, 1-8	4.4	23
200	Physiologically based biokinetic (PBBK) model for safrole bioactivation and detoxification in rats. *Chemical Research in Toxicology*, **2011**, 24, 818-34	4	23
199	Levels of Genotoxic and Carcinogenic Ingredients in Plant Food Supplements and Associated Risk Assessment. *Food and Nutrition Sciences (Print)*, **2011**, 02, 989-1010	0.4	53
198	Evaluation of research activities and research needs to increase the impact and applicability of alternative testing strategies in risk assessment practice. *Regulatory Toxicology and Pharmacology*, **2011**, 61, 105-14	3.4	28
197	Flavonoids and alkenylbenzenes: New concepts in bioactivation studies. *Chemico-Biological Interactions*, **2011**, 192, 87-95	5	14

196	Tutorial on physiologically based kinetic modeling in molecular nutrition and food research. *Molecular Nutrition and Food Research*, **2011**, 55, 941-56	5.9	39
195	Interaction of hesperetin glucuronide conjugates with human BCRP, MRP2 and MRP3 as detected in membrane vesicles of overexpressing baculovirus-infected Sf9 cells. *Biopharmaceutics and Drug Disposition*, **2011**, 32, 530-5	1.7	26
194	Safety assessment of plant food supplements (PFS). *Food and Function*, **2011**, 2, 760-8	6.1	28
193	Stable reporter cell lines for peroxisome proliferator-activated receptor (PPAR)-mediated modulation of gene expression. *Analytical Biochemistry*, **2011**, 414, 77-83	3.1	44
192	Quantitative proteomics and transcriptomics addressing the estrogen receptor subtype-mediated effects in T47D breast cancer cells exposed to the phytoestrogen genistein. *Molecular and Cellular Proteomics*, **2011**, 10, M110.002170	7.6	29
191	Quantitative comparison between in vivo DNA adduct formation from exposure to selected DNA-reactive carcinogens, natural background levels of DNA adduct formation and tumour incidence in rodent bioassays. *Mutagenesis*, **2011**, 26, 605-18	2.8	38
190	Phase II metabolism of hesperetin by individual UDP-glucuronosyltransferases and sulfotransferases and rat and human tissue samples. *Drug Metabolism and Disposition*, **2010**, 38, 617-25	4	73
189	Evaluation of human interindividual variation in bioactivation of estragole using physiologically based biokinetic modeling. *Toxicological Sciences*, **2010**, 113, 337-48	4.4	19
188	Stereoselective conjugation, transport and bioactivity of s- and R-hesperetin enantiomers in vitro. *Journal of Agricultural and Food Chemistry*, **2010**, 58, 6119-25	5.7	39
187	Superinduction of estrogen receptor mediated gene expression in luciferase based reporter gene assays is mediated by a post-transcriptional mechanism. *Journal of Steroid Biochemistry and Molecular Biology*, **2010**, 122, 204-11	5.1	45
186	Systematic construction of a conceptual minimal model of plasma cholesterol levels based on knockout mouse phenotypes. *Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids*, **2010**, 1801, 646-54	5	9
185	The use of in vitro toxicity data and physiologically based kinetic modeling to predict dose-response curves for in vivo developmental toxicity of glycol ethers in rat and man. *Toxicological Sciences*, **2010**, 118, 470-84	4.4	94
184	Application of bioassays in toxicological hazard, risk and impact assessments of dredged sediments. *Marine Pollution Bulletin*, **2010**, 60, 2026-42	6.7	30
183	A physiologically based biodynamic (PBBD) model for estragole DNA binding in rat liver based on in vitro kinetic data and estragole DNA adduct formation in primary hepatocytes. *Toxicology and Applied Pharmacology*, **2010**, 245, 57-66	4.6	52
182	Identification of nevadensin as an important herb-based constituent inhibiting estragole bioactivation and physiology-based biokinetic modeling of its possible in vivo effect. *Toxicology and Applied Pharmacology*, **2010**, 245, 179-90	4.6	40
181	Decrease of intracellular pH as possible mechanism of embryotoxicity of glycol ether alkoxyacetic acid metabolites. *Toxicology and Applied Pharmacology*, **2010**, 245, 236-43	4.6	14
180	The effect of co-administered flavonoids on the metabolism of hesperetin and the disposition of its metabolites in Caco-2 cell monolayers. *Molecular Nutrition and Food Research*, **2010**, 54, 851-60	5.9	42
179	In silico methods for physiologically based biokinetic models describing bioactivation and detoxification of coumarin and estragole: implications for risk assessment. *Molecular Nutrition and Food Research*, **2010**, 54, 195-207	5.9	29

178	Safety assessment of botanicals and botanical preparations used as ingredients in food supplements: testing an European Food Safety Authority-tiered approach. *Molecular Nutrition and Food Research*, **2010**, 54, 175-85	5.9	28
177	Role of surface charge and oxidative stress in cytotoxicity of organic monolayer-coated silicon nanoparticles towards macrophage NR8383 cells. *Particle and Fibre Toxicology*, **2010**, 7, 25	8.4	195
176	Relative developmental toxicity of glycol ether alkoxy acid metabolites in the embryonic stem cell test as compared with the in vivo potency of their parent compounds. *Toxicological Sciences*, **2009**, 110, 117-24	4.4	57
175	Estrogenicity of food-associated estrogenic compounds in the fetuses of female transgenic mice upon oral and IP maternal exposure. *Reproductive Toxicology*, **2009**, 27, 133-9	3.4	7
174	Use of physiologically based biokinetic (PBBK) modeling to study estragole bioactivation and detoxification in humans as compared with male rats. *Toxicological Sciences*, **2009**, 110, 255-69	4.4	61
173	An n-3 PUFA-rich microalgal oil diet protects to a similar extent as a fish oil-rich diet against AOM-induced colonic aberrant crypt foci in F344 rats. *Food and Chemical Toxicology*, **2009**, 47, 316-20	4.7	32
172	Quantitative structure-activity relationship modeling of the toxicity of organothiophosphate pesticides to Daphnia magna and Cyprinus carpio. *Chemosphere*, **2009**, 75, 1531-8	8.4	40
171	Potency of isothiocyanates to induce luciferase reporter gene expression via the electrophile-responsive element from murine glutathione S-transferase Ya. *Toxicology in Vitro*, **2009**, 23, 617-21	3.6	9
170	On the number of EINECS compounds that can be covered by (Q)SAR models for acute toxicity. *Toxicology Letters*, **2009**, 184, 67-72	4.4	19
169	Food-associated estrogenic compounds induce estrogen receptor-mediated luciferase gene expression in transgenic male mice. *Chemico-Biological Interactions*, **2008**, 174, 126-33	5	17
168	Cultivation of the heart urchin Echinocardium cordatum and validation of its use in marine toxicity testing for environmental risk assessment. *Journal of Experimental Marine Biology and Ecology*, **2008**, 364, 11-18	2.1	21
167	A physiologically based biokinetic (PBBK) model for estragole bioactivation and detoxification in rat. *Toxicology and Applied Pharmacology*, **2008**, 231, 248-59	4.6	66
166	Risk assessment of botanicals and botanical preparations intended for use in food and food supplements: emerging issues. *Toxicology Letters*, **2008**, 180, 131-6	4.4	59
165	Low dose estragole and 1?-hydroxyestragole induced DNA adduct formation. *Toxicology Letters*, **2008**, 180, S73	4.4	
164	Role of catechin quinones in the induction of EpRE-mediated gene expression. *Chemical Research in Toxicology*, **2008**, 21, 2352-60	4	38
163	Phytoestrogen-mediated inhibition of proliferation of the human T47D breast cancer cells depends on the ERalpha/ERbeta ratio. *Journal of Steroid Biochemistry and Molecular Biology*, **2008**, 112, 171-8	5.1	83
162	Shifted concentration dependency of EpRE- and XRE-mediated gene expression points at monofunctional EpRE-mediated induction by flavonoids at physiologically relevant concentrations. *Toxicology in Vitro*, **2008**, 22, 921-6	3.6	6
161	Differences in simulated liver concentrations of toxic coumarin metabolites in rats and different human populations evaluated through physiologically based biokinetic (PBBK) modeling. *Toxicology in Vitro*, **2008**, 22, 1890-901	3.6	26

160	Effects of flavonoid mixtures on the transport of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) through Caco-2 monolayers: an in vitro and kinetic modeling approach to predict the combined effects on transporter inhibition. *Food and Chemical Toxicology*, **2008**, 46, 557-66	4.7	11
159	Activation of EpRE-mediated gene transcription by quercetin glucuronides depends on their deconjugation. *Food and Chemical Toxicology*, **2008**, 46, 2128-34	4.7	22
158	Basil extract inhibits the sulfotransferase mediated formation of DNA adducts of the procarcinogen 1Phydroxyestragole by rat and human liver S9 homogenates and in HepG2 human hepatoma cells. *Food and Chemical Toxicology*, **2008**, 46, 2296-302	4.7	43
157	The FEMA GRAS assessment of alpha,beta-unsaturated aldehydes and related substances used as flavor ingredients. *Food and Chemical Toxicology*, **2008**, 46, 2935-67	4.7	40
156	Quercetin increases the bioavailability of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in rats. *Food and Chemical Toxicology*, **2008**, 46, 3422-8	4.7	6
155	Promises and pitfalls of quantitative structure-activity relationship approaches for predicting metabolism and toxicity. *Chemical Research in Toxicology*, **2008**, 21, 2229-36	4	68
154	QSAR models for predicting in vivo aquatic toxicity of chlorinated alkanes to fish. *Chemical Research in Toxicology*, **2008**, 21, 739-45	4	26
153	pH-Dependent radical scavenging capacity of green tea catechins. *Journal of Agricultural and Food Chemistry*, **2008**, 56, 816-23	5.7	101
152	Metabolism and transport of the citrus flavonoid hesperetin in Caco-2 cell monolayers. *Drug Metabolism and Disposition*, **2008**, 36, 1794-802	4	119
151	The influence of fruit and vegetable consumption and genetic variation on NAD(P)H:quinone oxidoreductase (NQO1) phenotype in an endoscopy-based population. *Nutrition and Cancer*, **2008**, 60, 204-15	2.8	3
150	Influence of cellular ERalpha/ERbeta ratio on the ERalpha-agonist induced proliferation of human T47D breast cancer cells. *Toxicological Sciences*, **2008**, 105, 303-11	4.4	98
149	Transcriptome and proteome profiling of colon mucosa from quercetin fed F344 rats point to tumor preventive mechanisms, increased mitochondrial fatty acid degradation and decreased glycolysis. *Proteomics*, **2008**, 8, 45-61	4.8	57
148	Human cytochrome p450 enzyme specificity for the bioactivation of estragole and related alkenylbenzenes. *Chemical Research in Toxicology*, **2007**, 20, 798-806	4	61
147	Molecular determinants of xenobiotic metabolism: QM/MM simulation of the conversion of 1-chloro-2,4-dinitrobenzene catalyzed by M1-1 glutathione S-transferase. *Biochemistry*, **2007**, 46, 6353-63	3.2	28
146	Human glutathione S-transferase-mediated glutathione conjugation of curcumin and efflux of these conjugates in Caco-2 cells. *Chemical Research in Toxicology*, **2007**, 20, 1895-902	4	47
145	Pathway and single gene analyses of inhibited Caco-2 differentiation by ascorbate-stabilized quercetin suggest enhancement of cellular processes associated with development of colon cancer. *Molecular Nutrition and Food Research*, **2007**, 51, 1031-45	5.9	11
144	Development of an on-line high performance liquid chromatography detection system for human cytochrome P450 1A2 inhibitors in extracts of natural products. *Journal of Chromatography A*, **2007**, 1141, 81-9	4.5	20
143	A case study on Bangka Island, Indonesia on the habits and consequences of pesticide use in pepper plantations. *Environmental Toxicology*, **2007**, 22, 405-14	4.2	7

142	The effect of quercetin phase II metabolism on its MRP1 and MRP2 inhibiting potential. *Biochemical Pharmacology*, **2007**, 74, 345-51	6	68
141	Tandem mass spectrometry analysis of N2-(trans-Isoestragol-3Pyl)-2Pdeoxyguanosine as a strategy to study species differences in sulfotransferase conversion of the proximate carcinogen 1Phydroxyestragole. *Chemical Research in Toxicology*, **2007**, 20, 991-8	4	55
140	Enforcement of the ban on aristolochic acids in Chinese traditional herbal preparations on the Dutch market. *Analytical and Bioanalytical Chemistry*, **2007**, 389, 263-75	4.4	45
139	Glutathione S-transferase phenotypes in relation to genetic variation and fruit and vegetable consumption in an endoscopy-based population. *Carcinogenesis*, **2007**, 28, 848-57	4.6	9
138	A comparative study on the effect of algal and fish oil on viability and cell proliferation of Caco-2 cells. *Food and Chemical Toxicology*, **2007**, 45, 716-24	4.7	28
137	Consequences of quercetin methylation for its covalent glutathione and DNA adduct formation. *Chemico-Biological Interactions*, **2006**, 160, 193-203	5	28
136	Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity. *Biochemical Pharmacology*, **2006**, 72, 217-26	6	41
135	An in vitro and in silico study on the flavonoid-mediated modulation of the transport of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) through Caco-2 monolayers. *Toxicology and Applied Pharmacology*, **2006**, 217, 204-15	4.6	18
134	Application of a continuous bioreactor cascade to study the effect of linoleic acid on hybridoma cell physiology. *Biotechnology and Bioengineering*, **2006**, 95, 370-83	4.9	4
133	Estrogenic potency of food-packaging-associated plasticizers and antioxidants as detected in ERalpha and ERbeta reporter gene cell lines. *Journal of Agricultural and Food Chemistry*, **2006**, 54, 4407-16	5.7	67
132	Future of toxicology--low-dose toxicology and risk--benefit analysis. *Chemical Research in Toxicology*, **2006**, 19, 977-81	4	32
131	Bioavailability of genistein and its glycoside genistin as measured in the portal vein of freely moving unanesthetized rats. *Journal of Agricultural and Food Chemistry*, **2006**, 54, 8006-12	5.7	52
130	Pro-oxidant activity of flavonoids induces EpRE-mediated gene expression. *Chemical Research in Toxicology*, **2006**, 19, 1499-505	4	162
129	Human cytochrome p450 enzymes of importance for the bioactivation of methyleugenol to the proximate carcinogen 1Phydroxymethyleugenol. *Chemical Research in Toxicology*, **2006**, 19, 111-6	4	59
128	Differential induction of electrophile-responsive element-regulated genes by n-3 and n-6 polyunsaturated fatty acids. *FEBS Letters*, **2006**, 580, 4587-90	3.8	17
127	Flavonoid-mediated inhibition of intestinal ABC transporters may affect the oral bioavailability of drugs, food-borne toxic compounds and bioactive ingredients. *Biomedicine and Pharmacotherapy*, **2006**, 60, 508-19	7.5	124
126	Myricetin stimulates the absorption of the pro-carcinogen PhIP. *Cancer Letters*, **2006**, 231, 36-42	9.9	15
125	Modulatory effects of quercetin on proliferation and differentiation of the human colorectal cell line Caco-2. *Cancer Letters*, **2006**, 238, 248-59	9.9	41

124	T-screen to quantify functional potentiating, antagonistic and thyroid hormone-like activities of poly halogenated aromatic hydrocarbons (PHAHs). *Toxicology in Vitro*, **2006**, 20, 490-8	3.6	77
123	In vivo relevance of two critical levels for NAD(P)H:quinone oxidoreductase (NQO1)-mediated cellular protection against electrophile toxicity found in vitro. *Toxicology in Vitro*, **2006**, 20, 594-600	3.6	11
122	Quercetin, but not its glycosidated conjugate rutin, inhibits azoxymethane-induced colorectal carcinogenesis in F344 rats. *Journal of Nutrition*, **2006**, 136, 2862-7	4.1	56
121	Quantum chemistry based quantitative structure-activity relationships for modeling the (sub)acute toxicity of substituted mononitrobenzenes in aquatic systems. *Environmental Toxicology and Chemistry*, **2006**, 25, 2313-21	3.8	16
120	Radical scavenging capacity of wine anthocyanins is strongly pH-dependent. *Journal of Agricultural and Food Chemistry*, **2005**, 53, 5526-34	5.7	74
119	Formation of transient covalent protein and DNA adducts by quercetin in cells with and without oxidative enzyme activity. *Chemical Research in Toxicology*, **2005**, 18, 1907-16	4	39
118	Tissue distribution of quercetin in rats and pigs. *Journal of Nutrition*, **2005**, 135, 1718-25	4.1	340
117	Quantitative structure activity relationship studies on the flavonoid mediated inhibition of multidrug resistance proteins 1 and 2. *Biochemical Pharmacology*, **2005**, 69, 699-708	6	154
116	Reversal of in vitro cellular MRP1 and MRP2 mediated vincristine resistance by the flavonoid myricetin. *Biochemical Pharmacology*, **2005**, 69, 1657-65	6	47
115	Inhibition of multidrug resistance proteins MRP1 and MRP2 by a series of alpha,beta-unsaturated carbonyl compounds. *Biochemical Pharmacology*, **2005**, 69, 1879-90	6	56
114	Human NAD(P)H:quinone oxidoreductase inhibition by flavonoids in living cells. *Free Radical Biology and Medicine*, **2005**, 39, 257-65	7.8	20
113	Flavonoids and alkenylbenzenes: mechanisms of mutagenic action and carcinogenic risk. *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, **2005**, 574, 124-38	3.3	183
112	Molecular mechanisms of toxicity of important food-borne phytotoxins. *Molecular Nutrition and Food Research*, **2005**, 49, 131-58	5.9	84
111	The stimulation of cell proliferation by quercetin is mediated by the estrogen receptor. *Molecular Nutrition and Food Research*, **2005**, 49, 763-71	5.9	92
110	The definition of hormesis and its implications for in vitro to in vivo extrapolation and risk assessment. *Critical Reviews in Toxicology*, **2005**, 35, 603-7	5.7	30
109	Metabolism of ATP-binding cassette drug transporter inhibitors: complicating factor for multidrug resistance. *Expert Opinion on Drug Metabolism and Toxicology*, **2005**, 1, 219-32	5.5	13
108	Structural requirements for the flavonoid-mediated modulation of glutathione S-transferase P1-1 and GS-X pump activity in MCF7 breast cancer cells. *Biochemical Pharmacology*, **2004**, 67, 1607-17	6	94
107	The effect of catechol O-methylation on radical scavenging characteristics of quercetin and luteolin--a mechanistic insight. *Free Radical Research*, **2004**, 38, 639-47	4	58

106	Human cytochrome p450 enzyme specificity for bioactivation of safrole to the proximate carcinogen 1Phydroxysafrole. *Chemical Research in Toxicology*, **2004**, 17, 1245-50	4	56
105	Identification of 14 quercetin phase II mono- and mixed conjugates and their formation by rat and human phase II in vitro model systems. *Chemical Research in Toxicology*, **2004**, 17, 1520-30	4	138
104	Rapid yeast estrogen bioassays stably expressing human estrogen receptors alpha and beta, and green fluorescent protein: a comparison of different compounds with both receptor types. *Journal of Steroid Biochemistry and Molecular Biology*, **2004**, 91, 99-109	5.1	154
103	Safety evaluation of neem (Azadirachta indica) derived pesticides. *Journal of Ethnopharmacology*, **2004**, 94, 25-41	5	137
102	Quenching of quercetin quinone/quinone methides by different thiolate scavengers: stability and reversibility of conjugate formation. *Chemical Research in Toxicology*, **2003**, 16, 822-31	4	62
101	Inhibition of human glutathione S-transferase P1-1 by the flavonoid quercetin. *Chemico-Biological Interactions*, **2003**, 145, 139-48	5	80
100	Degradation of 3,4-dichloro- and 3,4-difluoroaniline by Pseudomonas fluorescens 26-K. *Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes*, **2003**, 38, 121-32	2.2	44
99	Interplay between MRP inhibition and metabolism of MRP inhibitors: the case of curcumin. *Chemical Research in Toxicology*, **2003**, 16, 1642-51	4	96
98	Biphasic modulation of cell proliferation by quercetin at concentrations physiologically relevant in humans. *Cancer Letters*, **2003**, 200, 41-7	9.9	121
97	The role of quinone reductase (NQO1) and quinone chemistry in quercetin cytotoxicity. *Toxicology in Vitro*, **2003**, 17, 423-31	3.6	24
96	Ab Initio QM/MM Modeling of the Hydroxylation Step in p-Hydroxybenzoate Hydroxylase. *Journal of Physical Chemistry B*, **2003**, 107, 2118-2126	3.4	67
95	Conversion of 2-fluoromuconate to cis-dienelactone by purified enzymes of Rhodococcus opacus 1cp. *Applied and Environmental Microbiology*, **2003**, 69, 5636-42	4.8	11
94	A physiological threshold for protection against menadione toxicity by human NAD(P)H:quinone oxidoreductase (NQO1) in Chinese hamster ovary (CHO) cells. *Biochemical Pharmacology*, **2002**, 64, 1597-603	6	39
93	Flavoenzyme-Catalyzed Oxygenations and Oxidations of Phenolic Compounds. *Advanced Synthesis and Catalysis*, **2002**, 344, 1023-1035	5.6	47
92	Reductive deamination as a new step in the anaerobic microbial degradation of halogenated anilines. *FEMS Microbiology Letters*, **2002**, 209, 307-12	2.9	26
91	Regioselectivity of phase II metabolism of luteolin and quercetin by UDP-glucuronosyl transferases. *Chemical Research in Toxicology*, **2002**, 15, 662-70	4	196
90	Isolation and characterization of a microperoxidase-8 with a modified histidine axial ligand. *Journal of Biological Inorganic Chemistry*, **2002**, 7, 870-8	3.7	6
89	Identification of the human P450 enzymes involved in the in vitro metabolism of the synthetic steroidal hormones Org 4060 and Org 30659. *Xenobiotica*, **2002**, 32, 109-18	2	2

88	Quantum mechanical/molecular mechanical free energy simulations of the glutathione S-transferase (M1-1) reaction with phenanthrene 9,10-oxide. *Journal of the American Chemical Society*, **2002**, 124, 9926-36	16.4	76
87	Identification of o-quinone/quinone methide metabolites of quercetin in a cellular in vitro system. *FEBS Letters*, **2002**, 520, 30-4	3.8	76
86	The pro-oxidant chemistry of the natural antioxidants vitamin C, vitamin E, carotenoids and flavonoids. *Environmental Toxicology and Pharmacology*, **2002**, 11, 321-33	5.8	243
85	The regioselectivity of glutathione adduct formation with flavonoid quinone/quinone methides is pH-dependent. *Chemical Research in Toxicology*, **2002**, 15, 343-51	4	71
84	Structure-activity study on the quinone/quinone methide chemistry of flavonoids. *Chemical Research in Toxicology*, **2001**, 14, 398-408	4	128
83	The influence of pH on antioxidant properties and the mechanism of antioxidant action of hydroxyflavones. *Free Radical Biology and Medicine*, **2001**, 31, 869-81	7.8	282
82	The interplay of glutathione-related processes in antioxidant defense. *Environmental Toxicology and Pharmacology*, **2001**, 10, 141-52	5.8	219
81	Computer-modeling-based QSARs for analyzing experimental data on biotransformation and toxicity. *Toxicology in Vitro*, **2001**, 15, 539-51	3.6	55
80	Excretion and metabolism of desogestrel in healthy postmenopausal women. *Journal of Steroid Biochemistry and Molecular Biology*, **2001**, 78, 471-80	5.1	10
79	Transformation of the insecticide teflubenzuron by microorganisms. *Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes*, **2001**, 36, 559-67	2.2	11
78	Structure activity relationships for the chemical behaviour and toxicity of electrophilic quinones/quinone methides. *Advances in Experimental Medicine and Biology*, **2001**, 500, 11-21	3.6	7
77	Heme-(hydro)peroxide mediated O- and N-dealkylation. A study with microperoxidase. *FEBS Journal*, **2000**, 267, 6673-8		25
76	The mercapturic acid biotransformation pathway of hexachlorobenzene is not involved in the induction of splenomegaly, or skin and lung lesions in the Brown Norway rat. *Archives of Toxicology*, **2000**, 74, 609-17	5.8	5
75	Identification of fluoropyrogallols as new intermediates in biotransformation of monofluorophenols in Rhodococcus opacus 1cp. *Applied and Environmental Microbiology*, **2000**, 66, 2148-53	4.8	20
74	Peroxidase-catalyzed formation of quercetin quinone methide-glutathione adducts. *Archives of Biochemistry and Biophysics*, **2000**, 378, 224-33	4.1	145
73	A mechanism for oxygen exchange between ligated oxometalloporphinates and bulk water. *Biochemical and Biophysical Research Communications*, **2000**, 272, 551-6	3.4	11
72	Excretion balance and metabolism of the progestagen Org 30659 in healthy postmenopausal women. *Journal of Steroid Biochemistry and Molecular Biology*, **2000**, 73, 39-48	5.1	2
71	Modelling flavin and substrate substituent effects on the activation barrier and rate of oxygen transfer by p-hydroxybenzoate hydroxylase. *FEBS Letters*, **2000**, 478, 197-201	3.8	24

70	A Quantum Mechanical/Molecular Mechanical Study of the Hydroxylation of Phenol and Halogenated Derivatives by Phenol Hydroxylase. *Journal of the American Chemical Society*, **2000**, 122, 8728-8738	16.4	76
69	Characterization of different commercial soybean peroxidase preparations and use of the enzyme for N-demethylation of methyl N-methylanthranilate To produce the food flavor methylanthranilate. *Journal of Agricultural and Food Chemistry*, **2000**, 48, 1949-54	5.7	22
68	Regioselectivity and reversibility of the glutathione conjugation of quercetin quinone methide. *Chemical Research in Toxicology*, **2000**, 13, 185-91	4	115
67	Combined quantum mechanical and molecular mechanical reaction pathway calculation for aromatic hydroxylation by p-hydroxybenzoate-3-hydroxylase. *Journal of Molecular Graphics and Modelling*, **1999**, 17, 163-75, 214	2.8	34
66	Preferential oxidative dehalogenation upon conversion of 2-halophenols by Rhodococcus opacus 1G. *FEMS Microbiology Letters*, **1999**, 181, 73-82	2.9	24
65	The effect of iron to manganese substitution on microperoxidase 8 catalysed peroxidase and cytochrome P450 type of catalysis. *Journal of Biological Inorganic Chemistry*, **1999**, 4, 274-83	3.7	34
64	Differential substrate behaviour of phenol and aniline derivatives during conversion by horseradish peroxidase. *BBA - Proteins and Proteomics*, **1999**, 1435, 22-9		21
63	TEAC antioxidant activity of 4-hydroxybenzoates. *Free Radical Biology and Medicine*, **1999**, 27, 1427-36	7.8	71
62	Computational methods in flavin research. *Methods in Molecular Biology*, **1999**, 131, 207-28	1.4	5
61	Quantitative structure activity relationships for the electron transfer reactions of Anabaena PCC 7119 ferredoxin-NADP+ oxidoreductase with nitrobenzene and nitrobenzimidazolone derivatives: mechanistic implications. *FEBS Letters*, **1999**, 450, 44-8	3.8	5
60	Prooxidant toxicity of polyphenolic antioxidants to HL-60 cells: description of quantitative structure-activity relationships. *FEBS Letters*, **1999**, 462, 392-6	3.8	174
59	Antioxidant activities of carotenoids: quantitative relationships between theoretical calculations and experimental literature data. *Free Radical Research*, **1999**, 30, 233-40	4	29
58	Characterization of Iron(III) Tetramesitylporphyrin and Microperoxidase-8 Incorporated into the Molecular Sieve MCM-41. *Inorganic Chemistry*, **1999**, 38, 4901-4905	5.1	27
57	Reversible formation of high-valent-iron-oxo porphyrin intermediates in heme-based catalysis: revisiting the kinetic model for horseradish peroxidase. *Inorganica Chimica Acta*, **1998**, 275-276, 98-105	2.7	25
56	19F NMR study on the biodegradation of fluorophenols by various Rhodococcus species. *Biodegradation*, **1998**, 9, 475-86	4.1	44
55	Oxygen exchange with water in heme-oxo intermediates during H2O2-driven oxygen incorporation in aromatic hydrocarbons catalyzed by microperoxidase-8. *FEBS Journal*, **1998**, 253, 659-68		26
54	Quantitative structure/activity relationship for the rate of conversion of C4-substituted catechols by catechol-1,2-dioxygenase from Pseudomonas putida (arvilla) C1. *FEBS Journal*, **1998**, 257, 92-100		16
53	Molecular orbital study of porphyrin-substrate interactions in cytochrome P450 catalysed aromatic hydroxylation of substituted anilines. *Biophysical Chemistry*, **1998**, 73, 189-203	3.5	12

52	Quantitative structure activity relationships for the conversion of nitrobenzimidazolones and nitrobenzimidazoles by DT-diaphorase: implications for the kinetic mechanism. *FEBS Letters*, **1998**, 427, 325-9	3.8	7
51	Calculated ionisation potentials determine the oxidation of vanillin precursors by lignin peroxidase. *FEBS Letters*, **1998**, 430, 390-2	3.8	23
50	GSTP1-1 stereospecifically catalyzes glutathione conjugation of ethacrynic acid. *FEBS Letters*, **1998**, 441, 153-7	3.8	11
49	Correlation of Calculated Activation Energies with Experimental Rate Constants for an Enzyme Catalyzed Aromatic Hydroxylation. *Journal of the American Chemical Society*, **1998**, 120, 7641-7642	16.4	52
48	Conversion of pentahalogenated phenols by microperoxidase-8/H2O2 to benzoquinone-type products. *Chemical Research in Toxicology*, **1998**, 11, 1319-25	4	19
47	Occurrence of the NIH shift upon the cytochrome P450-catalyzed in vivo and in vitro aromatic ring hydroxylation of fluorobenzenes. *Chemical Research in Toxicology*, **1998**, 11, 503-12	4	48
46	5-Fluorouracil in colorectal cancer: rationale and clinical results of frequently used schedules. *Anti-Cancer Drugs*, **1998**, 9, 371-80	2.4	12
45	19F nuclear magnetic resonance as a tool to investigate microbial degradation of fluorophenols to fluorocatechols and fluoromuconates. *Applied and Environmental Microbiology*, **1998**, 64, 1256-63	4.8	39
44	Influence of substituents in fluorobenzene derivatives on the cytochrome P450-catalyzed hydroxylation at the adjacent ortho aromatic carbon center. *Chemical Research in Toxicology*, **1997**, 10, 279-88	4	14
43	Cytochrome P450-catalyzed oxidation of halobenzene derivatives. *Chemical Research in Toxicology*, **1997**, 10, 629-35	4	66
42	Microperoxidase/H2O2-mediated alkoxylating dehalogenation of halophenol derivatives in alcoholic media. *Proceedings of the National Academy of Sciences of the United States of America*, **1997**, 94, 4295-9	11.5	22
41	MP8-dependent oxidative dehalogenation: evidence for the direct formation of 1,4-benzoquinone from 4-fluorophenol by a peroxidase-type of reaction pathway. *Chemico-Biological Interactions*, **1997**, 104, 147-64	5	14
40	Metabolism of the insecticide teflubenzuron in rats. *Xenobiotica*, **1997**, 27, 801-17	2	5
39	5-Fluorouracil metabolite patterns in viable and necrotic tumor areas of murine colon carcinoma determined by 19F NMR spectroscopy. *Magnetic Resonance in Medicine*, **1996**, 36, 445-50	4.4	12
38	Regioselectivity and quantitative structure-activity relationships for the conjugation of a series of fluoronitrobenzenes by purified glutathione S-transferase enzymes from rat and man. *Chemical Research in Toxicology*, **1996**, 9, 638-46	4	36
37	Influence of the halogen-substituent pattern of fluoronitrobenzenes on their biotransformation and capacity to induce methemoglobinemia. *Toxicology and Applied Pharmacology*, **1996**, 139, 71-83	4.6	6
36	Relationships between the regioselectivity of the hydroxylation of C4-substituted 2-fluoroaniline derivatives and their toxic endpoints. *Toxicology and Applied Pharmacology*, **1996**, 141, 403-15	4.6	6
35	Interaction between the substrate and the high-valent-iron-oxo porphyrin cofactor as a possible factor influencing the regioselectivity of cytochrome P450 catalysed aromatic ring hydroxylation of 3-fluoro(methyl)anilines. *Chemico-Biological Interactions*, **1996**, 99, 129-46	5	9

34	Comparative MO-QSAR studies in various species including man. *Chemico-Biological Interactions*, **1996**, 100, 187-201	5	9
33	Molecular orbital study of the hydroxylation of benzene and monofluorobenzene catalysed by iron-oxo porphyrin □cation radical complexes. *Journal of Biological Inorganic Chemistry*, **1996**, 1, 192-204	3.7	29
32	The influence of the peptide chain on the kinetics and stability of microperoxidases. *FEBS Journal*, **1996**, 241, 215-20		21
31	Microperoxidase/H2O2-catalyzed aromatic hydroxylation proceeds by a cytochrome-P-450-type oxygen-transfer reaction mechanism. *FEBS Journal*, **1996**, 240, 232-8		63
30	Different metabolic pathways of 2,5-difluoronitrobenzene and 2,5-difluoroaminobenzene compared to molecular orbital substrate characteristics. *Chemico-Biological Interactions*, **1995**, 94, 49-72	5	13
29	Quantitative structure-activity relationships based on computer calculated parameters for the overall rate of glutathione S-transferase catalyzed conjugation of a series of fluoronitrobenzenes. *Chemical Research in Toxicology*, **1995**, 8, 481-8	4	29
28	The effect of varying halogen substituent patterns on the cytochrome P450 catalysed dehalogenation of 4-halogenated anilines to 4-aminophenol metabolites. *Biochemical Pharmacology*, **1995**, 49, 1235-48	6	44
27	1H NMR T1 relaxation rate study on substrate orientation of fluoromethylanilines in the active sites of microsomal and purified cytochromes P450 1A1 and 2B1. *FEBS Letters*, **1995**, 368, 279-84	3.8	9
26	Conversion of phenol derivatives to hydroxylated products by phenol hydroxylase from Trichosporon cutaneum. A comparison of regioselectivity and rate of conversion with calculated molecular orbital substrate characteristics. *FEBS Journal*, **1995**, 227, 284-91		45
25	A spectrophotometric assay for the detection of 2-aminophenols in biological samples. *Analytical Biochemistry*, **1994**, 220, 165-71	3.1	6
24	Catabolism of 4-hydroxybenzoate in Candida parapsilosis proceeds through initial oxidative decarboxylation by a FAD-dependent 4-hydroxybenzoate 1-hydroxylase. *FEMS Microbiology Letters*, **1994**, 121, 207-15	2.9	18
23	NADPH-cytochrome reductase catalysed redox cycling of 1,4-benzoquinone; hampered at physiological conditions, initiated at increased pH values. *Biochemical Pharmacology*, **1994**, 47, 1949-55	6	22
22	Influence of the type of halogen substituent on in vivo and in vitro phase II metabolism of 2-fluoro-4-halophenol metabolites formed from 3-halo-fluorobenzenes. *Xenobiotica*, **1994**, 24, 759-74	2	5
21	Regioselectivity of cytochrome P-450 catalyzed hydroxylation of fluorobenzenes predicted by calculated frontier orbital substrate characteristics. *Biochemistry*, **1993**, 32, 4801-12	3.2	81
20	Cytochrome P450-mediated oxidation of pentafluorophenol to tetrafluorobenzoquinone as the primary reaction product. *Chemical Research in Toxicology*, **1993**, 6, 674-80	4	40
19	Kinetics of cytochromes P-450 IA1 and IIB1 in reconstituted systems with dilauroyl- and distearoyl-glycerophosphocholine. *FEBS Journal*, **1993**, 215, 373-81		7
18	19F-NMR study on the pH-dependent regioselectivity and rate of the ortho-hydroxylation of 3-fluorophenol by phenol hydroxylase from Trichosporon cutaneum. Implications for the reaction mechanism. *FEBS Journal*, **1993**, 218, 345-53		17
17	A specific interaction between NADPH-cytochrome reductase and phosphatidylserine and phosphatidylinositol. *FEBS Journal*, **1993**, 218, 1021-9		14

16	The effect of age on inducibility of various types of rat liver cytochrome P-450. *Xenobiotica*, **1992**, 22, 515-22	2	24
15	Experimental and theoretical study on the redox cycling of resorufin by solubilized and membrane-bound NADPH-cytochrome reductase. *Chemical Research in Toxicology*, **1992**, 5, 268-73	4	15
14	A new hypothesis for the mechanism for cytochrome P-450 dependent aerobic conversion of hexahalogenated benzenes to pentahalogenated phenols. *Chemical Research in Toxicology*, **1992**, 5, 10-9	4	34
13	The involvement of primary and secondary metabolism in the covalent binding of 1,2- and 1,4-dichlorobenzenes. *Chemico-Biological Interactions*, **1992**, 84, 259-75	5	24
12	Rat liver microsomal metabolism of 2-halogenated 4-methylanilines. *Xenobiotica*, **1992**, 22, 1403-23	2	5
11	Interlaboratory comparison of microsomal ethoxyresorufin and pentoxyresorufin O-dealkylation determinations: standardization of assay conditions. *Archives of Toxicology*, **1992**, 66, 237-44	5.8	70
10	Frontier orbital study on the 4-hydroxybenzoate-3-hydroxylase-dependent activity with benzoate derivatives. *FEBS Journal*, **1992**, 206, 479-84		42
9	Reaction pathways for biodehalogenation of fluorinated anilines. *FEBS Journal*, **1990**, 194, 945-54		24
8	Development of a 19F-n.m.r. method for studies on the in vivo and in vitro metabolism of 2-fluoroaniline. *Xenobiotica*, **1990**, 20, 657-70	2	37
7	Microsomal metabolism of fluoroanilines. *Xenobiotica*, **1989**, 19, 1297-305	2	44
6	Methyl linoleate ozonide as a substrate for rat glutathione S-transferases: reaction pathway and isoenzyme selectivity. *Chemico-Biological Interactions*, **1989**, 69, 269-78	5	14
5	On the role of phospholipids in the reconstituted cytochrome P-450 system. A model study using dilauroyl and distearoyl glycerophosphocholine. *FEBS Journal*, **1989**, 181, 309-16		17
4	Impact Mechanisms of Ozone at Cell Level. *Studies in Environmental Science*, **1989**, 35, 513-522		
3	Fathobiochemical Effects in Rat Lung Related to Episodic Ozone Exposure. *Studies in Environmental Science*, **1989**, 35, 723-732		
2	Methyl linoleate ozonide: a substrate for rat glutathione S-transferases. *European Journal of Drug Metabolism and Pharmacokinetics*, **1987**, 12, 275-7	2.7	2
1	The role of glutathione and changes in thiol homeostasis in cultured lung cells exposed to ozone. *Toxicology*, **1985**, 35, 207-17	4.4	28