

John Giesy

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8106090/publications.pdf>

Version: 2024-02-01

829
papers

48,385
citations

2093

100
h-index

4101

175
g-index

840
all docs

840
docs citations

840
times ranked

28109
citing authors

#	ARTICLE	IF	CITATIONS
1	Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife.. Environmental Health Perspectives, 1998, 106, 775-792.	2.8	2,883
2	Global Distribution of Perfluorooctane Sulfonate in Wildlife. Environmental Science & Technology, 2001, 35, 1339-1342.	4.6	2,216
3	Pharmaceuticals and Personal Care Products in the Environment: What Are the Big Questions?. Environmental Health Perspectives, 2012, 120, 1221-1229.	2.8	1,033
4	Peer Reviewed: Perfluorochemical Surfactants in the Environment. Environmental Science & Technology, 2002, 36, 146A-152A.	4.6	913
5	Accumulation of Perfluorooctane Sulfonate in Marine Mammals. Environmental Science & Technology, 2001, 35, 1593-1598.	4.6	454
6	Perfluorinated Compounds in Aquatic Organisms at Various Trophic Levels in a Great Lakes Food Chain. Archives of Environmental Contamination and Toxicology, 2005, 48, 559-566.	2.1	432
7	Removal of antibiotics from wastewater by sewage treatment facilities in Hong Kong and Shenzhen, China. Water Research, 2008, 42, 395-403.	5.3	421
8	Dioxin-Like and Non-Dioxin-Like Toxic Effects of Polychlorinated Biphenyls (PCBs): Implications For Risk Assessment. Critical Reviews in Toxicology, 1998, 28, 511-569.	1.9	401
9	Perfluorooctanesulfonate and Related Fluorinated Hydrocarbons in Marine Mammals, Fishes, and Birds from Coasts of the Baltic and the Mediterranean Seas. Environmental Science & Technology, 2002, 36, 3210-3216.	4.6	380
10	Species-Specific Recombinant Cell Lines as Bioassay Systems for the Detection of 2,3,7,8-Tetrachlorodibenzo-p-dioxin-like Chemicals. Fundamental and Applied Toxicology, 1996, 30, 194-203.	1.9	369
11	Perfluorinated Compounds in Coastal Waters of Hong Kong, South China, and Korea. Environmental Science & Technology, 2004, 38, 4056-4063.	4.6	368
12	Analytical Methods for Detection of Selected Estrogenic Compounds in Aqueous Mixtures. Environmental Science & Technology, 1999, 33, 2814-2820.	4.6	367
13	Identification and Quantification of Estrogen Receptor Agonists in Wastewater Effluents. Environmental Science & Technology, 2001, 35, 3620-3625.	4.6	326
14	2-Chloro-s-Triazine Herbicides Induce Aromatase (CYP19) Activity in H295R Human Adrenocortical Carcinoma Cells: A Novel Mechanism for Estrogenicity?. Toxicological Sciences, 2000, 54, 121-127.	1.4	315
15	Perfluorooctanesulfonate and Related Fluorochemicals in Human Blood Samples from China. Environmental Science & Technology, 2006, 40, 715-720.	4.6	308
16	Toxicity Reference Values for the Toxic Effects of Polychlorinated Biphenyls to Aquatic Mammals. Human and Ecological Risk Assessment (HERA), 2000, 6, 181-201.	1.7	291
17	Chemical-Activated Luciferase Gene Expression (CALUX): A Novel in Vitro Bioassay for Ah Receptor Active Compounds in Sediments and Pore Water. Fundamental and Applied Toxicology, 1996, 33, 149-160.	1.9	283
18	Perfluorooctane Sulfonate in Fish-Eating Water Birds Including Bald Eagles and Albatrosses. Environmental Science & Technology, 2001, 35, 3065-3070.	4.6	275

#	ARTICLE	IF	CITATIONS
19	Health Risks in Infants Associated with Exposure to Perfluorinated Compounds in Human Breast Milk from Zhoushan, China. <i>Environmental Science & Technology</i> , 2006, 40, 2924-2929.	4.6	253
20	Derivation and application of relative potency estimates based on in vitro bioassay results. <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 2835-2843.	2.2	248
21	Biological impact of phthalates. <i>Toxicology Letters</i> , 2013, 217, 50-58.	0.4	247
22	Occurrence of Butyltin Compounds in Human Blood. <i>Environmental Science & Technology</i> , 1999, 33, 1776-1779.	4.6	241
23	Comparison of Ah Receptor-Mediated Luciferase and Ethoxyresorufin-O-deethylase Induction in H4IIE Cells: Implications for Their Use as Bioanalytical Tools for the Detection of Polyhalogenated Aromatic Hydrocarbons. <i>Toxicology and Applied Pharmacology</i> , 1996, 137, 316-325.	1.3	234
24	Endocrine disruption and consequences of chronic exposure to ibuprofen in Japanese medaka (<i>Oryzias latipes</i>). <i>Environmental Health Perspectives</i> , 2008, 116, 98, 256-264.	1.9	234
25	Relative Potencies of Individual Polychlorinated Naphthalenes and Halowax Mixtures To Induce Ah Receptor-Mediated Responses. <i>Environmental Science & Technology</i> , 2000, 34, 3153-3158.	4.6	233
26	Characterization of the H4IIE rat hepatoma cell bioassay as a tool for assessing toxic potency of planar halogenated hydrocarbons in environmental samples. <i>Environmental Science & Technology</i> , 1991, 25, 87-92.	4.6	232
27	Effects of Atrazine on Fish, Amphibians, and Aquatic Reptiles: A Critical Review. <i>Critical Reviews in Toxicology</i> , 2008, 38, 721-772.	1.9	226
28	Characterization and Distribution of Trace Organic Contaminants in Sediment from Masan Bay, Korea. 1. Instrumental Analysis. <i>Environmental Science & Technology</i> , 1999, 33, 4199-4205.	4.6	225
29	Spatial and Temporal Distribution of Polycyclic Aromatic Hydrocarbons in Sediments from Michigan Inland Lakes. <i>Environmental Science & Technology</i> , 2005, 39, 4700-4706.	4.6	221
30	Effects of chloro-s-triazine herbicides and metabolites on aromatase activity in various human cell lines and on vitellogenin production in male carp hepatocytes.. <i>Environmental Health Perspectives</i> , 2001, 109, 1027-1031.	2.8	219
31	Relative Potencies of Individual Polychlorinated Naphthalenes to Induce Dioxin-Like Responses in Fish and Mammalian In Vitro Bioassays. <i>Archives of Environmental Contamination and Toxicology</i> , 2000, 39, 273-281.	2.1	216
32	Removal of Phosphate from Eutrophic Lakes through Adsorption by in Situ Formation of Magnesium Hydroxide from Diatomite. <i>Environmental Science & Technology</i> , 2014, 48, 582-590.	4.6	213
33	Trans-Placental Transfer of Thirteen Perfluorinated Compounds and Relations with Fetal Thyroid Hormones. <i>Environmental Science & Technology</i> , 2011, 45, 7465-7472.	4.6	212
34	Origin of Hydroxylated Brominated Diphenyl Ethers: Natural Compounds or Man-Made Flame Retardants?. <i>Environmental Science & Technology</i> , 2009, 43, 7536-7542.	4.6	209
35	Ecological risk assessment of fifty pharmaceuticals and personal care products (PPCPs) in Chinese surface waters: A proposed multiple-level system. <i>Environment International</i> , 2020, 136, 105454.	4.8	203
36	Gene Expression Profiles in Rat Liver Treated With Perfluorooctanoic Acid (PFOA). <i>Toxicological Sciences</i> , 2006, 89, 93-107.	1.4	202

#	ARTICLE	IF	CITATIONS
37	Peer Reviewed: Analytical Challenges Hamper Perfluoroalkyl Research. <i>Environmental Science & Technology</i> , 2004, 38, 248A-255A.	4.6	201
38	Ecological risk assessment of atrazine in North American surface waters. <i>Environmental Toxicology and Chemistry</i> , 2013, 32, 10-11.	2.2	199
39	Relative potencies of individual polycyclic aromatic hydrocarbons to induce dioxinlike and estrogenic responses in three cell lines. <i>Environmental Toxicology</i> , 2002, 17, 128-137.	2.1	194
40	Perfluorooctanesulfonate and Related Fluorinated Hydrocarbons in Mink and River Otters from the United States. <i>Environmental Science & Technology</i> , 2002, 36, 2566-2571.	4.6	193
41	Effect of perinatal and postnatal bisphenol A exposure to the regulatory circuits at the hypothalamus-pituitary-gonadal axis of CD-1 mice. <i>Reproductive Toxicology</i> , 2011, 31, 409-417.	1.3	189
42	Inhibition of Gap Junctional Intercellular Communication by Perfluorinated Compounds in Rat Liver and Dolphin Kidney Epithelial Cell Lines in Vitro and Sprague-Dawley Rats in Vivo. <i>Toxicological Sciences</i> , 2002, 68, 429-436.	1.4	188
43	Vertical Profile of Polychlorinated Dibenzo-p-dioxins, Dibenzofurans, Naphthalenes, Biphenyls, Polycyclic Aromatic Hydrocarbons, and Alkylphenols in a Sediment Core from Tokyo Bay, Japan. <i>Environmental Science & Technology</i> , 2000, 34, 3560-3567.	4.6	173
44	Human adrenocarcinoma (H295R) cells for rapid in vitro determination of effects on steroidogenesis: Hormone production. <i>Toxicology and Applied Pharmacology</i> , 2006, 217, 114-124.	1.3	169
45	Polybrominated diphenyl ethers and their hydroxylated/methoxylated analogs: Environmental sources, metabolic relationships, and relative toxicities. <i>Marine Pollution Bulletin</i> , 2011, 63, 179-188.	2.3	169
46	Response of larval <i>Xenopus laevis</i> to atrazine: Assessment of growth, metamorphosis, and gonadal and laryngeal morphology. <i>Environmental Toxicology and Chemistry</i> , 2003, 22, 396-405.	2.2	167
47	Deformities in birds of the Great Lakes region. Assigning causality. <i>Environmental Science & Technology</i> , 1994, 28, 128A-135A.	4.6	166
48	Isomer-Specific Analysis and Toxic Evaluation of Polychlorinated Naphthalenes in Soil, Sediment, and Biota Collected near the Site of a Former Chlor-Alkali Plant. <i>Environmental Science & Technology</i> , 1998, 32, 2507-2514.	4.6	161
49	Organophosphate Esters in Sediment of the Great Lakes. <i>Environmental Science & Technology</i> , 2017, 51, 1441-1449.	4.6	161
50	Perfluorinated Compounds and Total and Extractable Organic Fluorine in Human Blood Samples from China. <i>Environmental Science & Technology</i> , 2008, 42, 8140-8145.	4.6	160
51	Freshwater Sediment Toxicity Bioassessment: Rationale for Species Selection and Test Design. <i>Journal of Great Lakes Research</i> , 1989, 15, 539-569.	0.8	159
52	Assessment of the Effects of Chemicals on the Expression of Ten Steroidogenic Genes in the H295R Cell Line Using Real-Time PCR. <i>Toxicological Sciences</i> , 2004, 81, 78-89.	1.4	159
53	Cancer risk assessments of Hong Kong soils contaminated by polycyclic aromatic hydrocarbons. <i>Journal of Hazardous Materials</i> , 2013, 261, 770-776.	6.5	158
54	Comparison of arsenic and antimony biogeochemical behavior in water, soil and tailings from Xikuangshan, China. <i>Science of the Total Environment</i> , 2016, 539, 97-104.	3.9	157

#	ARTICLE	IF	CITATIONS
55	The occurrence of selected antibiotics in Hong Kong coastal waters. <i>Marine Pollution Bulletin</i> , 2007, 54, 1287-1293.	2.3	155
56	Characterization of Organic Phosphorus in Lake Sediments by Sequential Fractionation and Enzymatic Hydrolysis. <i>Environmental Science & Technology</i> , 2013, 47, 7679-7687.	4.6	155
57	Effects of waterborne exposure of 17 β -estradiol on secondary sex characteristics and gonads of fathead minnows (<i>Pimephales promelas</i>). <i>Aquatic Toxicology</i> , 1999, 47, 129-145.	1.9	154
58	Polychloronaphthalenes and Other Dioxin-like Compounds in Arctic and Antarctic Marine Food Webs. <i>Environmental Science & Technology</i> , 2002, 36, 3490-3496.	4.6	145
59	Hydroxylated Polybrominated Diphenyl Ethers and Bisphenol A in Pregnant Women and Their Matching Fetuses: Placental Transfer and Potential Risks. <i>Environmental Science & Technology</i> , 2010, 44, 5233-5239.	4.6	143
60	Effects of land use on concentrations of metals in surface soils and ecological risk around Guanting Reservoir, China. <i>Environmental Geochemistry and Health</i> , 2007, 29, 459-471.	1.8	142
61	Alberta oil sands development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 951-952.	3.3	138
62	Cell bioassays for detection of aryl hydrocarbon (AhR) and estrogen receptor (ER) mediated activity in environmental samples. <i>Environmental Science and Pollution Research</i> , 2000, 7, 159-171.	2.7	137
63	Nonylphenol Isomers Differ in Estrogenic Activity. <i>Environmental Science & Technology</i> , 2006, 40, 5147-5153.	4.6	136
64	Trace Organic Contaminants in Sediment and Water from Ulsan Bay and Its Vicinity, Korea. <i>Archives of Environmental Contamination and Toxicology</i> , 2001, 40, 141-150.	2.1	134
65	What level of estrogenic activity determined by in vitro assays in municipal waste waters can be considered as safe?. <i>Environment International</i> , 2014, 64, 98-109.	4.8	134
66	Ecological Risk of Nonylphenol in China Surface Waters Based on Reproductive Fitness. <i>Environmental Science & Technology</i> , 2014, 48, 1256-1262.	4.6	132
67	Effects-Directed Analysis of Dissolved Organic Compounds in Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2015, 49, 12395-12404.	4.6	132
68	Concentrations and Profiles of Polychlorinated Naphthalene Congeners in Eighteen Technical Polychlorinated Biphenyl Preparations. <i>Environmental Science & Technology</i> , 2000, 34, 4236-4241.	4.6	131
69	Polychlorinated Naphthalenes and Polychlorinated Biphenyls in Fishes from Michigan Waters Including the Great Lakes. <i>Environmental Science & Technology</i> , 2000, 34, 566-572.	4.6	129
70	Toxicity of untreated and ozone-treated oil sands process-affected water (OSPW) to early life stages of the fathead minnow (<i>Pimephales promelas</i>). <i>Water Research</i> , 2012, 46, 6359-6368.	5.3	128
71	Avian Toxicity Reference Values for Perfluorooctane Sulfonate. <i>Environmental Science & Technology</i> , 2005, 39, 9357-9362.	4.6	127
72	In vitro profiling of the endocrine disrupting potency of organochlorine pesticides. <i>Toxicology Letters</i> , 2008, 183, 65-71.	0.4	127

#	ARTICLE	IF	CITATIONS
73	Involvement of activating ERK1/2 through G protein coupled receptor 30 and estrogen receptor $\hat{\pm}/\hat{t}^2$ in low doses of bisphenol A promoting growth of Sertoli TM4 cells. <i>Toxicology Letters</i> , 2014, 226, 81-89.	0.4	126
74	Real-time PCR array to study effects of chemicals on the Hypothalamicâ€Pituitaryâ€Gonadal axis of the Japanese medaka. <i>Aquatic Toxicology</i> , 2008, 88, 173-182.	1.9	124
75	The photoenhanced toxicity of anthracene to juvenile sunfish (<i>Lepomis</i> spp.). <i>Aquatic Toxicology</i> , 1985, 6, 133-146.	1.9	123
76	Development of a marine fish model for studying in vivo molecular responses in ecotoxicology. <i>Aquatic Toxicology</i> , 2008, 86, 131-141.	1.9	122
77	Cell bioassays for detection of aryl hydrocarbon (AhR) and estrogen receptor (ER) mediated activity in environmental samples. <i>Marine Pollution Bulletin</i> , 2002, 45, 3-16.	2.3	121
78	Ball milling synthesis of covalent organic framework as a highly active photocatalyst for degradation of organic contaminants. <i>Journal of Hazardous Materials</i> , 2019, 369, 494-502.	6.5	121
79	Ecological risk assessment of arsenic and metals in sediments of coastal areas of northern Bohai and Yellow Seas, China. <i>Ambio</i> , 2010, 39, 367-375.	2.8	120
80	Perfluorinated compounds in surface waters from Northern China: Comparison to level of industrialization. <i>Environment International</i> , 2012, 42, 37-46.	4.8	120
81	Contribution of known endocrine disrupting substances to the estrogenic activity in Tama River water samples from Japan using instrumental analysis and in vitro reporter gene assay. <i>Water Research</i> , 2004, 38, 4491-4501.	5.3	119
82	Effects of Waterborne Exposure to 4-Nonylphenol and Nonylphenol Ethoxylate on Secondary Sex Characteristics and Gonads of Fathead Minnows (<i>Pimephales promelas</i>). <i>Environmental Research</i> , 1999, 80, S122-S137.	3.7	118
83	Concentrations of neonicotinoid insecticides in honey, pollen and honey bees (<i>Apis mellifera</i> L.) in central Saskatchewan, Canada. <i>Chemosphere</i> , 2016, 144, 2321-2328.	4.2	117
84	Malformations of the endangered Chinese sturgeon, <i>Acipenser sinensis</i> , and its causal agent. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 9339-9344.	3.3	116
85	Dietary exposure of mink to carp from Saginaw Bay, Michigan. 1. Effects on reproduction and survival, and the potential risks to wild mink populations. <i>Archives of Environmental Contamination and Toxicology</i> , 1995, 28, 334-43.	2.1	114
86	Bisphenol A Disrupts Steroidogenesis in Human H295R Cells. <i>Toxicological Sciences</i> , 2011, 121, 320-327.	1.4	114
87	Effects of 20 PBDE metabolites on steroidogenesis in the H295R cell line. <i>Toxicology Letters</i> , 2008, 176, 230-238.	0.4	113
88	Effects of Prochloraz or Propylthiouracil on the Cross-Talk between the HPG, HPA, and HPT Axes in Zebrafish. <i>Environmental Science & Technology</i> , 2011, 45, 769-775.	4.6	113
89	Polychlorinated naphthalenes, biphenyls, dibenzoâ€dioxins, and dibenzofurans as well as polycyclic aromatic hydrocarbons and alkylphenols in sediment from the Detroit and Rouge Rivers, Michigan, USA. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 1878-1889.	2.2	109
90	Effects of atrazine on metamorphosis, growth, laryngeal and gonadal development, aromatase activity, and sex steroid concentrations in <i>Xenopus laevis</i> . <i>Ecotoxicology and Environmental Safety</i> , 2005, 62, 160-173.	2.9	109

#	ARTICLE	IF	CITATIONS
91	Responses of the Proteome and Metabolome in Livers of Zebrafish Exposed Chronically to Environmentally Relevant Concentrations of Microcystin-LR. <i>Environmental Science & Technology</i> , 2017, 51, 596-607.	4.6	109
92	Atrazine concentrations, gonadal gross morphology and histology in ranid frogs collected in Michigan agricultural areas. <i>Aquatic Toxicology</i> , 2006, 76, 230-245.	1.9	108
93	Polychlorinated Dibenzo-p-dioxin and Dibenzofuran Concentration Profiles in Sediments and Flood-Plain Soils of the Tittabawassee River, Michigan. <i>Environmental Science & Technology</i> , 2003, 37, 468-474.	4.6	107
94	Environmentally Relevant Concentrations of the Flame Retardant Tris(1,3-dichloro-2-propyl) Phosphate Inhibit Growth of Female Zebrafish and Decrease Fecundity. <i>Environmental Science & Technology</i> , 2015, 49, 14579-14587.	4.6	107
95	Tissue Distribution and Maternal Transfer of Poly- and Perfluorinated Compounds in Chinese Sturgeon (<i>Acipenser sinensis</i>): Implications for Reproductive Risk. <i>Environmental Science & Technology</i> , 2010, 44, 1868-1874.	4.6	106
96	Alkaline Digestion and Solid Phase Extraction Method for Perfluorinated Compounds in Mussels and Oysters from South China and Japan. <i>Archives of Environmental Contamination and Toxicology</i> , 2006, 50, 240-248.	2.1	105
97	Shifts in production of perfluoroalkyl acids affect emissions and concentrations in the environment of the Xiaoqing River Basin, China. <i>Journal of Hazardous Materials</i> , 2016, 307, 55-63.	6.5	104
98	Cytotoxicity of Ag, Au and Ag-Au bimetallic nanoparticles prepared using golden rod (<i>Solidago</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 46	1.6	104
99	The Impact of Metallic Coagulants on the Removal of Organic Compounds from Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2011, 45, 8452-8459.	4.6	103
100	Frequency Distributions of Trace Metal Concentrations in Five Freshwater Fishes. <i>Transactions of the American Fisheries Society</i> , 1977, 106, 393-403.	0.6	102
101	H4IIIE rat hepatoma cell bioassay-derived 2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents in colonial fish-eating waterbird eggs from the Great Lakes. <i>Archives of Environmental Contamination and Toxicology</i> , 1991, 21, 91-101.	2.1	102
102	Humic Acids Reduce Bioaccumulation of Some Polycyclic Aromatic Hydrocarbons. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1983, 40, s63-s69.	0.7	101
103	Perfluorooctane Sulfonate in Oysters, <i>Crassostrea virginica</i> , from the Gulf of Mexico and the Chesapeake Bay, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 2002, 42, 313-318.	2.1	101
104	Relative Potencies of Individual Chlorinated and Brominated Polycyclic Aromatic Hydrocarbons for Induction of Aryl Hydrocarbon Receptor-Mediated Responses. <i>Environmental Science & Technology</i> , 2009, 43, 2159-2165.	4.6	101
105	Amino Acid Sequence of the Ligand-Binding Domain of the Aryl Hydrocarbon Receptor 1 Predicts Sensitivity of Wild Birds to Effects of Dioxin-Like Compounds. <i>Toxicological Sciences</i> , 2013, 131, 139-152.	1.4	101
106	In Vitro Vitellogenin Production by Carp (<i>Cyprinus carpio</i>) Hepatocytes as a Screening Method for Determining (Anti)Estrogenic Activity of Xenobiotics. <i>Toxicology and Applied Pharmacology</i> , 1999, 157, 68-76.	1.3	100
107	Quantification of rainbow trout (<i>Oncorhynchus mykiss</i>) zona radiata and vitellogenin mRNA levels using real-time PCR after in vivo treatment with estradiol-17 β or \pm -zearalenol. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2000, 75, 109-119.	1.2	98
108	Interconversion of Hydroxylated and Methoxylated Polybrominated Diphenyl Ethers in Japanese Medaka. <i>Environmental Science & Technology</i> , 2010, 44, 8729-8735.	4.6	98

#	ARTICLE	IF	CITATIONS
109	Polyhalogenated Carbazoles in Sediments of Lake Michigan: A New Discovery. <i>Environmental Science & Technology</i> , 2014, 48, 12807-12815.	4.6	98
110	Occurrence of Thyroid Hormone Activities in Drinking Water from Eastern China: Contributions of Phthalate Esters. <i>Environmental Science & Technology</i> , 2012, 46, 1811-1818.	4.6	97
111	Controlling Air Pollution from Straw Burning in China Calls for Efficient Recycling. <i>Environmental Science & Technology</i> , 2012, 46, 7934-7936.	4.6	97
112	Challenges of using blooms of <i>Microcystis</i> spp. in animal feeds: A comprehensive review of nutritional, toxicological and microbial health evaluation. <i>Science of the Total Environment</i> , 2021, 764, 142319.	3.9	97
113	Quantitative RT-PCR Methods for Evaluating Toxicant-Induced Effects on Steroidogenesis Using the H295R Cell Line. <i>Environmental Science & Technology</i> , 2005, 39, 2777-2785.	4.6	96
114	The use of biomarkers in ecological risk assessment: recommendations from the Christchurch conference on Biomarkers in Ecotoxicology. <i>Biomarkers</i> , 2001, 6, 1-6.	0.9	95
115	Removal of Estrogenic Activity from Municipal Waste Landfill Leachate Assessed with a Bioassay Based on Reporter Gene Expression. <i>Environmental Science & Technology</i> , 2003, 37, 3430-3434.	4.6	95
116	Perfluorinated compounds in estuarine and coastal areas of north Bohai Sea, China. <i>Marine Pollution Bulletin</i> , 2011, 62, 1905-1914.	2.3	95
117	Uptake of planar polychlorinated biphenyls and 2,3,7,8-substituted polychlorinated dibenzofurans and dibenzo-p-dioxins by birds nesting in the lower fox river and Green Bay, Wisconsin, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 1993, 24, 332-344.	2.1	94
118	Simultaneous quantification of multiple classes of phenolic compounds in blood plasma by liquid chromatography-electrospray tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2010, 1217, 506-513.	1.8	94
119	Testicular Signaling Is the Potential Target of Perfluorooctanesulfonate-Mediated Subfertility in Male Mice ¹ . <i>Biology of Reproduction</i> , 2011, 84, 1016-1023.	1.2	93
120	Interactions between aryl hydrocarbon receptor (AhR) and hypoxia signaling pathways. <i>Environmental Toxicology and Pharmacology</i> , 2001, 10, 17-27.	2.0	92
121	Polychlorinated Naphthalenes, -Biphenyls, -Dibenzo-p-dioxins, and -Dibenzofurans in Double-Crested Cormorants and Herring Gulls from Michigan Waters of the Great Lakes. <i>Environmental Science & Technology</i> , 2001, 35, 441-447.	4.6	91
122	Identification of genes responsive to PFOS using gene expression profiling. <i>Environmental Toxicology and Pharmacology</i> , 2005, 19, 57-70.	2.0	91
123	Comparison of approaches to quantify SARS-CoV-2 in wastewater using RT-qPCR: Results and implications from a collaborative inter-laboratory study in Canada. <i>Journal of Environmental Sciences</i> , 2021, 107, 218-229.	3.2	91
124	Genotoxicity of Several Polybrominated Diphenyl Ethers (PBDEs) and Hydroxylated PBDEs, and Their Mechanisms of Toxicity. <i>Environmental Science & Technology</i> , 2011, 45, 5003-5008.	4.6	90
125	China's Soil Pollution Control: Choices and Challenges. <i>Environmental Science & Technology</i> , 2016, 50, 13181-13183.	4.6	90
126	Bioaccumulation and Toxic Potential of Extremely Hydrophobic Polychlorinated Biphenyl Congeners in Biota Collected at a Superfund Site Contaminated with Aroclor 1268. <i>Environmental Science & Technology</i> , 1998, 32, 1214-1221.	4.6	89

#	ARTICLE	IF	CITATIONS
127	Predicting Water Quality Criteria for Protecting Aquatic Life from Physicochemical Properties of Metals or Metalloids. <i>Environmental Science & Technology</i> , 2013, 47, 446-453.	4.6	89
128	Bioaccumulation characteristics of perfluoroalkyl acids (PFAAs) in coastal organisms from the west coast of South Korea. <i>Chemosphere</i> , 2015, 129, 157-163.	4.2	89
129	Hydroxylated Polychlorinated Biphenyl Metabolites Are Anti-estrogenic in a Stably Transfected Human Breast Adenocarcinoma (MCF7) Cell Line. <i>Toxicology and Applied Pharmacology</i> , 1997, 144, 363-376.	1.3	88
130	Butyltin Residues in Southern Sea Otters (<i>Enhydra lutris nereis</i>) Found Dead along California Coastal Waters. <i>Environmental Science & Technology</i> , 1998, 32, 1169-1175.	4.6	88
131	EFFECTS OF AIR CELL INJECTION OF PERFLUOROCTANE SULFONATE BEFORE INCUBATION ON DEVELOPMENT OF THE WHITE LEGHORN CHICKEN (<i>GALLUS DOMESTICUS</i>) EMBRYO. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 227.	2.2	88
132	Alkylphenols, polycyclic aromatic hydrocarbons, and organochlorines in sediment from Lake Shihwa, Korea: Instrumental and bioanalytical characterization. <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 2424-2432.	2.2	87
133	Effects of perfluorooctane sulfonate on mallard and northern bobwhite quail exposed chronically via the diet. <i>Environmental Toxicology and Pharmacology</i> , 2007, 23, 1-9.	2.0	87
134	The H295R system for evaluation of endocrine-disrupting effects. <i>Ecotoxicology and Environmental Safety</i> , 2006, 65, 293-305.	2.9	86
135	Contaminants in fishes from Great Lakes-influenced sections and above dams of three Michigan rivers. II: Implications for health of mink. <i>Archives of Environmental Contamination and Toxicology</i> , 1994, 27, 213-23.	2.1	84
136	Bioaccumulation profiles of polychlorinated biphenyl congeners and organochlorine pesticides in Ganges river dolphins. <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 1511-1520.	2.2	83
137	Occurrence of Estrogenic Compounds in and Removal by a Swine Farm Waste Treatment Plant. <i>Environmental Science & Technology</i> , 2006, 40, 7896-7902.	4.6	83
138	Ecogenomics of Zooplankton Community Reveals Ecological Threshold of Ammonia Nitrogen. <i>Environmental Science & Technology</i> , 2017, 51, 3057-3064.	4.6	83
139	ALKYLPHENOLS, POLYCYCLIC AROMATIC HYDROCARBONS, AND ORGANOCHLORINES IN SEDIMENT FROM LAKE SHIHWA, KOREA: INSTRUMENTAL AND BIOANALYTICAL CHARACTERIZATION. <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 2424.	2.2	83
140	Responses of the Medaka HPG Axis PCR Array and Reproduction to Prochloraz and Ketoconazole. <i>Environmental Science & Technology</i> , 2008, 42, 6762-6769.	4.6	82
141	In vitro bioassays for detecting dioxin-like activity " Application potentials and limits of detection, a review. <i>Science of the Total Environment</i> , 2014, 487, 37-48.	3.9	82
142	Traditional and new POPs in environments along the Bohai and Yellow Seas: An overview of China and South Korea. <i>Chemosphere</i> , 2017, 169, 503-515.	4.2	82
143	Linking the molecular composition of autochthonous dissolved organic matter to source identification for freshwater lake ecosystems by combination of optical spectroscopy and FT-ICR-MS analysis. <i>Science of the Total Environment</i> , 2020, 703, 134764.	3.9	82
144	Combined cytotoxicity of polystyrene nanoplastics and phthalate esters on human lung epithelial A549 cells and its mechanism. <i>Ecotoxicology and Environmental Safety</i> , 2021, 213, 112041.	2.9	82

#	ARTICLE	IF	CITATIONS
145	Plasma concentrations of estradiol and testosterone, gonadal aromatase activity and ultrastructure of the testis in <i>Xenopus laevis</i> exposed to estradiol or atrazine. <i>Aquatic Toxicology</i> , 2005, 72, 383-396.	1.9	81
146	Untargeted Identification of Organo-Bromine Compounds in Lake Sediments by Ultrahigh-Resolution Mass Spectrometry with the Data-Independent Precursor Isolation and Characteristic Fragment Method. <i>Analytical Chemistry</i> , 2015, 87, 10237-10246.	3.2	81
147	Effects of Tris(1,3-dichloro-2-propyl) Phosphate on Growth, Reproduction, and Gene Transcription of <i>Daphnia magna</i> at Environmentally Relevant Concentrations. <i>Environmental Science & Technology</i> , 2015, 49, 12975-12983.	4.6	81
148	InÂvitro characterization of the effectiveness of enhanced sewage treatment processes to eliminate endocrine activity of hospital effluents. <i>Water Research</i> , 2013, 47, 1545-1557.	5.3	80
149	Spatial and Temporal Trends of Polyhalogenated Carbazoles in Sediments of Upper Great Lakes: Insights into Their Origin. <i>Environmental Science & Technology</i> , 2017, 51, 89-97.	4.6	80
150	Characterization and Distribution of Trace Organic Contaminants in Sediment from Masan Bay, Korea. 2. In Vitro Gene Expression Assays. <i>Environmental Science & Technology</i> , 1999, 33, 4206-4211.	4.6	79
151	Review of the effects of endocrine-disrupting chemicals in birds. <i>Pure and Applied Chemistry</i> , 2003, 75, 2287-2303.	0.9	78
152	Bioassay-Derived 2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin Equivalents in PCB-Containing Extracts from the Flesh and Eggs of Lake Michigan Chinook Salmon (<i>Oncorhynchus tshawytscha</i>) and Possible Implications for Reproduction. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1991, 48, 1685-1690.	0.7	77
153	Effect of Ozonation on the Estrogenicity and Androgenicity of Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2011, 45, 6268-6274.	4.6	77
154	Two Years after the Hebei Spirit Oil Spill: Residual Crude-Derived Hydrocarbons and Potential AhR-Mediated Activities in Coastal Sediments. <i>Environmental Science & Technology</i> , 2012, 46, 1406-1414.	4.6	77
155	Effectiveness of Ozonation Treatment in Eliminating Toxicity of Oil Sands Process-Affected Water to <i>Chironomus dilutus</i> . <i>Environmental Science & Technology</i> , 2012, 46, 486-493.	4.6	77
156	The OECD validation program of the H295R steroidogenesis assay: Phase 3. Final inter-laboratory validation study. <i>Environmental Science and Pollution Research</i> , 2011, 18, 503-515.	2.7	76
157	Organophosphorus insecticides in honey, pollen and bees (<i>Apis mellifera</i> L.) and their potential hazard to bee colonies in Egypt. <i>Ecotoxicology and Environmental Safety</i> , 2015, 114, 1-8.	2.9	76
158	Bisphenol A modulates colorectal cancer protein profile and promotes the metastasis via induction of epithelial to mesenchymal transitions. <i>Archives of Toxicology</i> , 2015, 89, 1371-1381.	1.9	75
159	Cu/Cu ₂ O/CuO loaded on the carbon layer derived from novel precursors with amazing catalytic performance. <i>Science of the Total Environment</i> , 2016, 571, 380-387.	3.9	75
160	Identification and Quantitation of Nonylphenol Ethoxylates and Nonylphenol in Fish Tissues from Michigan. <i>Environmental Science & Technology</i> , 2001, 35, 10-13.	4.6	74
161	Support of Science-Based Decisions Concerning the Evaluation of the Toxicology of Mixtures: A New Beginning. <i>Regulatory Toxicology and Pharmacology</i> , 2002, 36, 34-39.	1.3	73
162	Historical trends of inorganic and organic fluorine in sediments of Lake Michigan. <i>Chemosphere</i> , 2014, 114, 203-209.	4.2	73

#	ARTICLE	IF	CITATIONS
163	Kinetics and biotransformation of benzo(a)pyrene in <i>Chironomus riparius</i> . <i>Archives of Environmental Contamination and Toxicology</i> , 1982, 11, 25-31.	2.1	72
164	Acute and Chronic Effects of Perfluorobutane Sulfonate (PFBS) on the Mallard and Northern Bobwhite Quail. <i>Archives of Environmental Contamination and Toxicology</i> , 2008, 54, 535-545.	2.1	72
165	Competitive interaction between soil-derived humic acid and phosphate on goethite. <i>Applied Geochemistry</i> , 2013, 36, 125-131.	1.4	72
166	Multi-pathway assessment of human health risk posed by polycyclic aromatic hydrocarbons. <i>Environmental Geochemistry and Health</i> , 2015, 37, 587-601.	1.8	72
167	Bioaccumulation, Biotransformation, and Toxicity of BDE-47, 6-OH-BDE-47, and 6-MeO-BDE-47 in Early Life-Stages of Zebrafish (<i>Danio rerio</i>). <i>Environmental Science & Technology</i> , 2015, 49, 1823-1833.	4.6	72
168	Hydroxylated and methylsulfonyl polychlorinated biphenyl metabolites in albatrosses from Midway Atoll, North Pacific Ocean. <i>Environmental Toxicology and Chemistry</i> , 1998, 17, 1620-1625.	2.2	71
169	Organochlorine Pesticides, Polychlorinated Biphenyls, and Butyltin Compounds in Blubber and Livers of Stranded California Sea Lions, Elephant Seals, and Harbor Seals from Coastal California, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 2001, 41, 90-99.	2.1	71
170	Effects of Atrazine on Metamorphosis, Growth, and Gonadal Development in the Green Frog (<i>Rana</i>). <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 1000-1007.	1.1	71
171	Activation of the aryl hydrocarbon receptor by berberine in HepG2 and H4IIE cells: Biphasic effect on CYP1A1. <i>Biochemical Pharmacology</i> , 2005, 70, 925-936.	2.0	71
172	Horizontal and Vertical Distribution of Estrogenic Activities in Sediments and Waters from Tokyo Bay, Japan. <i>Archives of Environmental Contamination and Toxicology</i> , 2005, 48, 209-216.	2.1	71
173	In vitro endocrine disruption and TCDD-like effects of three novel brominated flame retardants: TBPH, TBB, & TBCO. <i>Toxicology Letters</i> , 2013, 223, 252-259.	0.4	71
174	Deformities, PCBs, and TCDD-Equivalents in Double-Crested Cormorants (<i>Phalacrocorax auritus</i>) and Caspian Terns (<i>Hydroprogne caspia</i>) of the Upper Great Lakes 1986-1991: Testing a Cause-Effect Hypothesis. <i>Journal of Great Lakes Research</i> , 1996, 22, 172-197.	0.8	70
175	Perfluoroalkyl Acids in the Egg Yolk of Birds from Lake Shihwa, Korea. <i>Environmental Science & Technology</i> , 2008, 42, 5821-5827.	4.6	70
176	Predicting toxic potencies of metal oxide nanoparticles by means of nano-QSARs. <i>Nanotoxicology</i> , 2016, 10, 1207-1214.	1.6	70
177	Hydroxylated and methoxylated polybrominated diphenyl ethers in blood plasma of humans in Hong Kong. <i>Environment International</i> , 2012, 47, 66-72.	4.8	69
178	Bioaccumulation of microcystins (MCs) in four fish species from Lake Taihu, China: Assessment of risks to humans. <i>Science of the Total Environment</i> , 2014, 487, 224-232.	3.9	69
179	Depuration kinetics and tissue disposition of PFOA and PFOS in white leghorn chickens (<i>Gallus gallus</i>) administered by subcutaneous implantation. <i>Ecotoxicology and Environmental Safety</i> , 2009, 72, 26-36.	2.9	68
180	Halogenated POPs and PAHs in Blood Plasma of Hong Kong Residents. <i>Environmental Science & Technology</i> , 2011, 45, 1630-1637.	4.6	68

#	ARTICLE	IF	CITATIONS
181	Perfluorinated Compounds in Water, Sediment and Soil from Guanting Reservoir, China. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2011, 87, 74-79.	1.3	68
182	Accumulation and Biotransformation of BDE-47 by Zebrafish Larvae and Teratogenicity and Expression of Genes along the Hypothalamus-Pituitary-Thyroid Axis. <i>Environmental Science & Technology</i> , 2012, 46, 12943-12951.	4.6	68
183	Transcriptional Responses of the Brain-Gonadal-Liver Axis of Fathead Minnows Exposed to Untreated and Ozone-Treated Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2012, 46, 9701-9708.	4.6	68
184	Risks posed by trace organic contaminants in coastal sediments in the Pearl River Delta, China. <i>Marine Pollution Bulletin</i> , 2005, 50, 1036-1049.	2.3	67
185	Gonadal Development of Larval Male <i>Xenopus laevis</i> Exposed to Atrazine in Outdoor Microcosms. <i>Environmental Science & Technology</i> , 2005, 39, 5255-5261.	4.6	67
186	Using feathers to assess risk of mercury and selenium to bald eagle reproduction in the Great Lakes region. <i>Archives of Environmental Contamination and Toxicology</i> , 1994, 27, 294-298.	2.1	66
187	Comparison of fathead minnow ovary explant and H295R cell-based steroidogenesis assays for identifying endocrine-active chemicals. <i>Ecotoxicology and Environmental Safety</i> , 2007, 68, 20-32.	2.9	66
188	Cyanobacteria blooms produce teratogenic retinoic acids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 9477-9482.	3.3	66
189	Effects of exposure to oil sands process-affected water from experimental reclamation ponds on <i>Chironomus dilutus</i> . <i>Water Research</i> , 2012, 46, 1662-1672.	5.3	66
190	Concentrations of organochlorine pesticides (OCPs) in human blood plasma from Hong Kong: Markers of exposure and sources from fish. <i>Environment International</i> , 2013, 54, 18-25.	4.8	66
191	Regulation of engineered nanomaterials: current challenges, insights and future directions. <i>Environmental Science and Pollution Research</i> , 2018, 25, 3060-3077.	2.7	66
192	Novel Insights into the Kinetics, Evolved Gases, and Mechanisms for Biomass (Sugar Cane Residue) Pyrolysis. <i>Environmental Science & Technology</i> , 2019, 53, 13495-13505.	4.6	66
193	PLASMA SEX STEROID CONCENTRATIONS AND GONADAL AROMATASE ACTIVITIES IN AFRICAN CLAWED FROGS (<i>XENOPUS LAEVIS</i>) FROM SOUTH AFRICA. <i>Environmental Toxicology and Chemistry</i> , 2004, 23, 1996.	2.2	65
194	Effects of Atrazine on CYP19 Gene Expression and Aromatase Activity in Testes and on Plasma Sex Steroid Concentrations of Male African Clawed Frogs (<i>Xenopus laevis</i>). <i>Toxicological Sciences</i> , 2005, 86, 273-280.	1.4	65
195	Tissue Concentrations of Polybrominated Compounds in Chinese Sturgeon (<i>Acipenser sinensis</i>): Origin, Hepatic Sequestration, and Maternal Transfer. <i>Environmental Science & Technology</i> , 2010, 44, 5781-5786.	4.6	64
196	Genotoxicity and Endocrine-Disruption Potentials of Sediment near an Oil Spill Site: Two Years after the Hebei Spirit Oil Spill. <i>Environmental Science & Technology</i> , 2011, 45, 7481-7488.	4.6	64
197	Estrogen-, androgen- and aryl hydrocarbon receptor mediated activities in passive and composite samples from municipal waste and surface waters. <i>Environment International</i> , 2013, 59, 372-383.	4.8	64
198	Effects of 3,3',4,4',5-pentachlorobiphenyl (PCB 126) and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) injected into the yolks of chicken (<i>Gallus domesticus</i>) eggs prior to incubation. <i>Archives of Environmental Contamination and Toxicology</i> , 1996, 31, 404-409.	2.1	63

#	ARTICLE	IF	CITATIONS
199	Novel trends in endocrine disruptor testing: the H295R Steroidogenesis Assay for identification of inducers and inhibitors of hormone production. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 390, 287-291.	1.9	63
200	Responses of earthworms and microbial communities in their guts to Triclosan. <i>Chemosphere</i> , 2017, 168, 1194-1202.	4.2	63
201	Occurrences and Fates of Hydroxylated Polybrominated Diphenyl Ethers in Marine Sediments in Relation to Trophodynamics. <i>Environmental Science & Technology</i> , 2012, 46, 2148-2155.	4.6	62
202	Integrated assessment of contaminated sediments in the lower Fox River and Green Bay, Wisconsin. <i>Ecotoxicology and Environmental Safety</i> , 1992, 23, 46-63.	2.9	61
203	Polychlorinated biphenyls and chlorinated insecticides in plasma of Caspian terns: Relationships with age, productivity, and colony site tenacity in the great lakes. <i>Archives of Environmental Contamination and Toxicology</i> , 1993, 24, 320-331.	2.1	61
204	Characterization of dioxin-like activity of sediments from a Czech River Basin. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 2768-2777.	2.2	61
205	Perfluoroalkyl Acids in Marine Organisms from Lake Shihwa, Korea. <i>Archives of Environmental Contamination and Toxicology</i> , 2009, 57, 552-560.	2.1	61
206	Estrogenic potencies of several environmental pollutants, as determined by vitellogenin induction in a carp hepatocyte assay. <i>Toxicological Sciences</i> , 1999, 50, 206-213.	1.4	60
207	Effects of sulfathiazole, oxytetracycline and chlortetracycline on steroidogenesis in the human adrenocarcinoma (H295R) cell line and freshwater fish <i>Oryzias latipes</i> . <i>Journal of Hazardous Materials</i> , 2010, 182, 494-502.	6.5	60
208	Relative importance of food and water sources to cadmium uptake by <i>Gambusia affinis</i> (poeciliidae). <i>Environmental Research</i> , 1978, 16, 326-332.	3.7	59
209	Reproductive Responses of Common Carp (<i>Cyprinus carpio</i>) Exposed in Cages to Influent of the Las Vegas Wash in Lake Mead, Nevada, from Late Winter to Early Spring. <i>Environmental Science & Technology</i> , 2004, 38, 6385-6395.	4.6	59
210	The endocrine disrupting potential of sediments from the Upper Danube River (Germany) as revealed by in vitro bioassays and chemical analysis. <i>Environmental Science and Pollution Research</i> , 2011, 18, 446-460.	2.7	59
211	Phorate-induced oxidative stress, DNA damage and transcriptional activation of p53 and caspase genes in male Wistar rats. <i>Toxicology and Applied Pharmacology</i> , 2012, 259, 54-65.	1.3	59
212	Environmental geochemical and spatial/temporal behavior of total and speciation of antimony in typical contaminated aquatic environment from Xikuangshan, China. <i>Microchemical Journal</i> , 2018, 137, 181-189.	2.3	59
213	Removal of antimonate (Sb(V)) and antimonite (Sb(III)) from aqueous solutions by coagulation-flocculation-sedimentation (CFS): Dependence on influencing factors and insights into removal mechanisms. <i>Science of the Total Environment</i> , 2018, 644, 1277-1285.	3.9	59
214	2,3,7,8-Tetrachlorodibenzo-p-dioxin equivalents in tissues of birds at Green Bay, Wisconsin, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 1993, 24, 345-354.	2.1	58
215	Thyroid hormone disrupting activities associated with phthalate esters in water sources from Yangtze River Delta. <i>Environment International</i> , 2012, 42, 117-123.	4.8	58
216	Shape-dependent toxicity of alumina nanoparticles in rat astrocytes. <i>Science of the Total Environment</i> , 2019, 690, 158-166.	3.9	58

#	ARTICLE	IF	CITATIONS
217	In Vitro Antiestrogenic Effects of Aryl Methyl Sulfone Metabolites of Polychlorinated Biphenyls and 2,2-Bis(4-chlorophenyl)-1,1-dichloroethene on 17beta-Estradiol-Induced Gene Expression in Several Bioassay Systems. <i>Toxicological Sciences</i> , 2002, 69, 362-372.	1.4	57
218	Effects of chronic dietary exposure to environmentally relevant concentrations to 2,3,7,8-tetrachlorodibenzo-p-dioxin on survival, growth, reproduction and biochemical responses of female rainbow trout (<i>Oncorhynchus mykiss</i>). <i>Aquatic Toxicology</i> , 2002, 59, 35-53.	1.9	57
219	Modulation of steroidogenic gene expression and hormone production of H295R cells by pharmaceuticals and other environmentally active compounds. <i>Toxicology and Applied Pharmacology</i> , 2007, 225, 142-153.	1.3	57
220	Assessment of chemical effects on aromatase activity using the H295R cell line. <i>Environmental Science and Pollution Research</i> , 2010, 17, 1137-1148.	2.7	57
221	Families of Nuclear Receptors in Vertebrate Models: Characteristic and Comparative Toxicological Perspective. <i>Scientific Reports</i> , 2015, 5, 8554.	1.6	57
222	Influence of natural organic matter on the bioavailability and preservation of organic phosphorus in lake sediments. <i>Chemical Geology</i> , 2015, 397, 51-60.	1.4	57
223	Prediction of Concentrations of 2,3,7,8-Tetrachlorodibenzo-p-dioxin Equivalents from Total Concentrations of Polychlorinated Biphenyls in Fish Fillets. <i>Environmental Science & Technology</i> , 1992, 26, 1151-1159.	4.6	56
224	Assessing the Toxicity of Naphthenic Acids Using a Microbial Genome Wide Live Cell Reporter Array System. <i>Environmental Science & Technology</i> , 2011, 45, 1984-1991.	4.6	56
225	Dietary intake of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) from fish and meat by residents of Nanjing, China. <i>Environment International</i> , 2012, 42, 138-143.	4.8	56
226	Sonochemical degradation of perfluorinated chemicals in aqueous film-forming foams. <i>Journal of Hazardous Materials</i> , 2016, 317, 275-283.	6.5	56
227	Tiered probabilistic assessment of organohalogen compounds in the Han River and Danjiangkou Reservoir, central China. <i>Science of the Total Environment</i> , 2017, 586, 163-173.	3.9	56
228	Efficient removal of both antimonite ($Sb(III)$) and antimonate ($Sb(V)$) from environmental water using titanate nanotubes and nanoparticles. <i>Environmental Science: Nano</i> , 2019, 6, 834-850.	2.2	56
229	Caspian Tern Reproduction in the Saginaw Bay Ecosystem Following a 100-Year Flood Event. <i>Journal of Great Lakes Research</i> , 1993, 19, 96-108.	0.8	55
230	Effects of exposure to municipal wastewater in situ on the reproductive physiology of the fathead minnow (<i>Pimephales promelas</i>). <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 2001-2012.	2.2	55
231	Environmental fate and bioavailability of agent orange and its associated dioxin during the vietnam war. <i>Environmental Science and Pollution Research</i> , 2004, 11, 359-370.	2.7	55
232	Pharmacokinetics and Acute Lethality of Perfluorooctanesulfonate (PFOS) to Juvenile Mallard and Northern Bobwhite. <i>Archives of Environmental Contamination and Toxicology</i> , 2006, 50, 411-420.	2.1	55
233	Cetyltrimethylammonium Bromide-Coated Fe_3O_4 Magnetic Nanoparticles for Analysis of 15 Trace Polycyclic Aromatic Hydrocarbons in Aquatic Environments by Ultrapformance, Liquid Chromatography With Fluorescence Detection. <i>Analytical Chemistry</i> , 2015, 87, 7667-7675.	3.2	55
234	Refocusing on Nonpriority Toxic Metals in the Aquatic Environment in China. <i>Environmental Science & Technology</i> , 2017, 51, 3117-3118.	4.6	55

#	ARTICLE	IF	CITATIONS
235	Prenatal transfer of decabromodiphenyl ether (BDE-209) results in disruption of the thyroid system and developmental toxicity in zebrafish offspring. <i>Aquatic Toxicology</i> , 2017, 190, 46-52.	1.9	55
236	Effects of ultraviolet radiation on the primary production of natural phytoplankton assemblages in Lake Michigan. <i>Ecotoxicology and Environmental Safety</i> , 1991, 22, 345-361.	2.9	54
237	Analysis of trace organic contaminants in sediment, pore water, and water samples from Onsan Bay, Korea: Instrumental analysis and in vitro gene expression assay. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 1796-1803.	2.2	54
238	Cytochrome P4501A Induction by 2,3,7,8-Tetrachlorodibenzo-p-Dioxin and Two Chlorinated Dibenzofurans in Primary Hepatocyte Cultures of Three Avian Species. <i>Toxicological Sciences</i> , 2010, 113, 380-391.	1.4	54
239	Sequence and In Vitro Function of Chicken, Ring-Necked Pheasant, and Japanese Quail AHR1 Predict In Vivo Sensitivity to Dioxins. <i>Environmental Science & Technology</i> , 2012, 46, 2967-2975.	4.6	54
240	Biota-sediment accumulation factor (BSAF), bioaccumulation factor (BAF), and contaminant levels in prey fish to indicate the extent of PAHs and OCPs contamination in eggs of waterbirds. <i>Environmental Science and Pollution Research</i> , 2013, 20, 8425-8434.	2.7	54
241	Isomer-Specific Accumulation of Perfluorooctanesulfonate from (N-Ethyl) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 507 Td <i>Environmental Science & Technology</i> , 2014, 48, 1058-1066.	4.6	54
242	Europe-wide survey of estrogenicity in wastewater treatment plant effluents: the need for the effect-based monitoring. <i>Environmental Science and Pollution Research</i> , 2014, 21, 10970-10982.	2.7	54
243	Contaminants in fishes from great lakes-influenced sections and above dams of three Michigan Rivers: III. Implications for health of bald eagles. <i>Archives of Environmental Contamination and Toxicology</i> , 1995, 29, 309-321.	2.1	53
244	Instrumental and bioanalytical measures of dioxin-like and estrogenic compounds and activities associated with sediment from the Korean coast. <i>Ecotoxicology and Environmental Safety</i> , 2005, 61, 366-379.	2.9	53
245	Preparation and evaluation of a neutral methacrylate-based monolithic column for hydrophilic interaction stationary phase by pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 4611-4617.	1.8	53
246	Comparison of extraction and quantification methods of perfluorinated compounds in human plasma, serum, and whole blood. <i>Analytica Chimica Acta</i> , 2008, 628, 214-221.	2.6	52
247	Chronic Exposure of Marine Medaka (<i>Oryzias melastigma</i>) to 4,5-Dichloro-2-n-octyl-4-isothiazolin-3-one (DCOIT) Reveals Its Mechanism of Action in Endocrine Disruption via the Hypothalamus-Pituitary-Gonadal-Liver (HPGL) Axis. <i>Environmental Science & Technology</i> , 2016, 50, 4492-4501.	4.6	51
248	Polycyclic aromatic hydrocarbons in soils from the Central-Himalaya region: Distribution, sources, and risks to humans and wildlife. <i>Science of the Total Environment</i> , 2016, 556, 12-22.	3.9	51
249	Synthesis of Fe ₃ O ₄ magnetic nanoparticles coated with cationic surfactants and their applications in Sb(V) removal from water. <i>Science of the Total Environment</i> , 2020, 710, 136302.	3.9	51
250	A combined hydraulic and toxicological approach to assess re-suspended sediments during simulated flood events. Part I—multiple biomarkers in rainbow trout. <i>Journal of Soils and Sediments</i> , 2010, 10, 1347-1361.	1.5	50
251	Dioxin-like Potency of HO- and MeO- Analogues of PBDEs™ the Potential Risk through Consumption of Fish from Eastern China. <i>Environmental Science & Technology</i> , 2012, 46, 10781-10788.	4.6	50
252	PAHs in surface sediments from coastal and estuarine areas of the northern Bohai and Yellow Seas, China. <i>Environmental Geochemistry and Health</i> , 2012, 34, 445-456.	1.8	50

#	ARTICLE	IF	CITATIONS
253	Perfluoroalkyl and polyfluoroalkyl substances in sediments from South Bohai coastal watersheds, China. <i>Marine Pollution Bulletin</i> , 2014, 85, 619-627.	2.3	50
254	Perfluorobutanesulfonate Exposure Causes Durable and Transgenerational Dysbiosis of Gut Microbiota in Marine Medaka. <i>Environmental Science and Technology Letters</i> , 2018, 5, 731-738.	3.9	50
255	Global Biomonitoring of Perfluorinated Organics. <i>Scientific World Journal, The</i> , 2001, 1, 627-629.	0.8	49
256	Estrogenic activity in extracts and exudates of cyanobacteria and green algae. <i>Environment International</i> , 2012, 39, 134-140.	4.8	49
257	Toxicity of pentachlorophenol to native aquatic species in the Yangtze River. <i>Environmental Science and Pollution Research</i> , 2012, 19, 609-618.	2.7	49
258	A tiered ecological risk assessment of three chlorophenols in Chinese surface waters. <i>Environmental Science and Pollution Research</i> , 2012, 19, 1544-1554.	2.7	49
259	Metals in agricultural soils and plants in Egypt. <i>Toxicological and Environmental Chemistry</i> , 2014, 96, 730-742.	0.6	49
260	Are styrene oligomers in coastal sediments of an industrial area aryl hydrocarbon-receptor agonists?. <i>Environmental Pollution</i> , 2016, 213, 913-921.	3.7	49
261	Current and historical concentrations of poly and perfluorinated compounds in sediments of the northern Great Lakes " Superior, Huron, and Michigan. <i>Environmental Pollution</i> , 2018, 236, 373-381.	3.7	49
262	Metals associated with organic carbon extracted from Okefenokee Swamp water. <i>Chemical Geology</i> , 1977, 20, 109-120.	1.4	48
263	Butyltin compounds in sediment and fish from the Polish Coast of the Baltic Sea. <i>Environmental Science and Pollution Research</i> , 1999, 6, 200-206.	2.7	48
264	Surfactant-modified flowerlike layered double hydroxide-coated magnetic nanoparticles for preconcentration of phthalate esters from environmental water samples. <i>Journal of Chromatography A</i> , 2015, 1414, 22-30.	1.8	48
265	Combined Transcriptomic and Proteomic Approach to Identify Toxicity Pathways in Early Life Stages of Japanese Medaka (<i>Oryzias latipes</i>) Exposed to 1,2,5,6-Tetrabromocyclooctane (TBCO). <i>Environmental Science & Technology</i> , 2016, 50, 7781-7790.	4.6	48
266	Integration of metabolomics and transcriptomics reveals short-chain chlorinated paraffin-induced hepatotoxicity in male Sprague-Dawley rat. <i>Environment International</i> , 2019, 133, 105231.	4.8	48
267	Cadmium and zinc accumulation and elimination by freshwater crayfish. <i>Archives of Environmental Contamination and Toxicology</i> , 1980, 9, 683-697.	2.1	47
268	Differential Accumulation of Polychlorinated Biphenyl Congeners in the Terrestrial Food Web of the Kalamazoo River Superfund Site, Michigan. <i>Environmental Science & Technology</i> , 2005, 39, 5954-5963.	4.6	47
269	Occurrence of Atrazine and Related Compounds in Sediments of Upper Great Lakes. <i>Environmental Science & Technology</i> , 2016, 50, 7335-7343.	4.6	47
270	Interaction of alkaline phosphatase with minerals and sediments: Activities, kinetics and hydrolysis of organic phosphorus. <i>Colloids and Surfaces A: Physicochemical and Engineering Aspects</i> , 2016, 495, 46-53.	2.3	47

#	ARTICLE	IF	CITATIONS
271	Newly Identified AhR-Active Compounds in the Sediments of an Industrial Area Using Effect-Directed Analysis. <i>Environmental Science & Technology</i> , 2019, 53, 10043-10052.	4.6	47
272	Effects of long-term exposure to pentachlorophenol on the free amino acid pool and energy reserves of the freshwater amphipod <i>Gammarus pseudolimnaeus bousfield</i> (crustacea, amphipoda). <i>Ecotoxicology and Environmental Safety</i> , 1986, 12, 233-251.	2.9	46
273	In vitro induction of ethoxyresorufin O-deethylase and porphyrins by halogenated aromatic hydrocarbons in avian primary hepatocytes. <i>Environmental Toxicology and Chemistry</i> , 1998, 17, 2006-2018.	2.2	46
274	Risk Assessment of Organohalogenated Compounds in Water Bird Eggs from South China. <i>Environmental Science & Technology</i> , 2008, 42, 6296-6302.	4.6	46
275	Transport of short-chain perfluoroalkyl acids from concentrated fluoropolymer facilities to the Daling River estuary, China. <i>Environmental Science and Pollution Research</i> , 2015, 22, 9626-9636.	2.7	46
276	Effects of monovalent and divalent metal cations on the aggregation and suspension of Fe ₃ O ₄ magnetic nanoparticles in aqueous solution. <i>Science of the Total Environment</i> , 2017, 586, 817-826.	3.9	46
277	Distribution and Elimination of Polychlorinated Dibenzo-p-dioxins, Dibenzofurans, Biphenyls, and p,p'-DDE in Tissues of Bald Eagles from the Upper Peninsula of Michigan. <i>Environmental Science & Technology</i> , 2002, 36, 2789-2796.	4.6	45
278	CYTOTOXICITY AND ARYL HYDROCARBON RECEPTOR-MEDIATED ACTIVITY OF N-HETEROCYCLIC POLYCYCLIC AROMATIC HYDROCARBONS: STRUCTURE-ACTIVITY RELATIONSHIPS. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 1291.	2.2	45
279	Sensitivity of Japanese Quail (<i>Coturnix japonica</i>), Common Pheasant (<i>Phasianus colchicus</i>), and White Leghorn Chicken (<i>Gallus gallus domesticus</i>) Embryos to In Ovo Exposure to TCDD, PeCDF, and TCDF. <i>Toxicological Sciences</i> , 2011, 119, 93-103.	1.4	45
280	AhR-mediated potency of sediments and soils in estuarine and coastal areas of the Yellow Sea region: A comparison between Korea and China. <i>Environmental Pollution</i> , 2012, 171, 216-225.	3.7	45
281	pH-dependent aquatic criteria for 2,4-dichlorophenol, 2,4,6-trichlorophenol and pentachlorophenol. <i>Science of the Total Environment</i> , 2012, 441, 125-131.	3.9	45
282	Combined effects of cadmium and fluoranthene on germination, growth and photosynthesis of soybean seedlings. <i>Journal of Environmental Sciences</i> , 2013, 25, 1936-1946.	3.2	45
283	Mutagenic Azo Dyes, Rather Than Flame Retardants, Are the Predominant Brominated Compounds in House Dust. <i>Environmental Science & Technology</i> , 2016, 50, 12669-12677.	4.6	45
284	Site-specific water quality criteria for aquatic ecosystems: A case study of pentachlorophenol for Tai Lake, China. <i>Science of the Total Environment</i> , 2016, 541, 65-73.	3.9	45
285	Untargeted Screening and Distribution of Organo-Bromine Compounds in Sediments of Lake Michigan. <i>Environmental Science & Technology</i> , 2016, 50, 321-330.	4.6	45
286	Combining High-Throughput Sequencing of DNA and Traditional Paleolimnological Techniques To Infer Historical Trends in Cyanobacterial Communities. <i>Environmental Science & Technology</i> , 2018, 52, 6842-6853.	4.6	45
287	Spatial and interspecies differences in concentrations of eight trace elements in wild freshwater fishes at different trophic levels from middle and eastern China. <i>Science of the Total Environment</i> , 2019, 672, 883-892.	3.9	45
288	Cellular alterations in midgut cells of honey bee workers (<i>Apis mellifera</i> L.) exposed to sublethal concentrations of CdO or PbO nanoparticles or their binary mixture. <i>Science of the Total Environment</i> , 2019, 651, 1356-1367.	3.9	45

#	ARTICLE	IF	CITATIONS
289	Structures of Endocrine-Disrupting Chemicals Determine Binding to and Activation of the Estrogen Receptor α and Androgen Receptor. <i>Environmental Science & Technology</i> , 2020, 54, 11424-11433.	4.6	45
290	Next generation per- and poly-fluoroalkyl substances: Status and trends, aquatic toxicity, and risk assessment. , 2022, 1, 117-131.		45
291	Tissue metal concentrations in two crayfish species cohabiting a tennessee cave stream. <i>Oecologia</i> , 1979, 44, 8-12.	0.9	44
292	Copper speciation in soft, acid, humic waters: Effects on copper bioaccumulation by and toxicity to <i>Daphnia magna</i> (Daphniidae). <i>Science of the Total Environment</i> , 1983, 28, 23-36.	3.9	44
293	Quantitative and qualitative characteristics of dissolved organic matter from eight dominant aquatic macrophytes in Lake Dianchi, China. <i>Environmental Science and Pollution Research</i> , 2013, 20, 7413-7423.	2.7	44
294	Predicting the sensitivity of fishes to dioxin-like compounds: possible role of the aryl hydrocarbon receptor (AhR) ligand binding domain. <i>Environmental Science and Pollution Research</i> , 2013, 20, 1219-1224.	2.7	44
295	Sensitivity of early life stages of white sturgeon, rainbow trout, and fathead minnow to copper. <i>Ecotoxicology</i> , 2013, 22, 139-147.	1.1	44
296	Methylated PACs are more potent than their parent compounds: A study of aryl hydrocarbon receptor-mediated activity, degradability, and mixture interactions in the H4IIE assay. <i>Environmental Toxicology and Chemistry</i> , 2018, 37, 1409-1419.	2.2	44
297	Relationships Between Chlorinated Hydrocarbon Concentrations and Rearing Mortality of Chinook Salmon (<i>Oncorhynchus Tshawytscha</i>) Eggs from Lake Michigan. <i>Journal of Great Lakes Research</i> , 1986, 12, 82-98.	0.8	43
298	Toxicity of Sediments and Sediment Pore Waters from the Grand Calumet River-Indiana Harbor, Indiana Area of Concern. <i>Ecotoxicology and Environmental Safety</i> , 1993, 26, 86-112.	2.9	43
299	Pharmaceuticals and Personal Care Products in the Waters of Lake Mead, Nevada. <i>ACS Symposium Series</i> , 2001, , 116-139.	0.5	43
300	Contribution of Synthetic and Naturally Occurring Organobromine Compounds to Bromine Mass in Marine Organisms. <i>Environmental Science & Technology</i> , 2010, 44, 6068-6073.	4.6	43
301	Effect-directed analysis and mixture effects of AhR-active PAHs in crude oil and coastal sediments contaminated by the Hebei Spirit oil spill. <i>Environmental Pollution</i> , 2015, 199, 110-118.	3.7	43
302	Time-dependent inhibitory effects of Tris(1, 3-dichloro-2-propyl) phosphate on growth and transcription of genes involved in the GH/IGF axis, but not the HPT axis, in female zebrafish. <i>Environmental Pollution</i> , 2017, 229, 470-478.	3.7	43
303	The dose makes the poison. <i>Science of the Total Environment</i> , 2018, 621, 649-653.	3.9	43
304	Composition characterization and biotransformation of dissolved, particulate and algae organic phosphorus in eutrophic lakes. <i>Environmental Pollution</i> , 2020, 265, 114838.	3.7	43
305	Congener profile of polychlorinated/brominated dibenzo-p-dioxins and dibenzofurans in soil and sediments collected at a former chlor-alkali plant. <i>Toxicological and Environmental Chemistry</i> , 1998, 67, 135-146.	0.6	42
306	Instrumental and Bioanalytical Measures of Persistent Organochlorines in Blue Mussel (<i>Mytilus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 639, 360-368.	2.1	42

#	ARTICLE	IF	CITATIONS
307	Organochlorine pollutants [corrected] in California sea lions revisited. <i>BMC Ecology</i> , 2002, 2, 11.	3.0	42
308	Sedimentary record of polycyclic aromatic hydrocarbons and DDTs in Dianchi Lake, an urban lake in Southwest China. <i>Environmental Science and Pollution Research</i> , 2013, 20, 5471-5480.	2.7	42
309	Magnetic Nanoparticles Interaction with Humic Acid: In the Presence of Surfactants. <i>Environmental Science & Technology</i> , 2016, 50, 8640-8648.	4.6	42
310	A Cross-species Quantitative Adverse Outcome Pathway for Activation of the Aryl Hydrocarbon Receptor Leading to Early Life Stage Mortality in Birds and Fishes. <i>Environmental Science & Technology</i> , 2018, 52, 7524-7533.	4.6	42
311	Hotspots and trends of covalent organic frameworks (COFs) in the environmental and energy field: Bibliometric analysis. <i>Science of the Total Environment</i> , 2021, 783, 146838.	3.9	42
312	Effects of nonylphenol ethoxylate exposure on reproductive output and bioindicators of environmental estrogen exposure in fathead minnows, <i>Pimephales promelas</i> . <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 510-522.	2.2	41
313	HCH and DDT in Sediments from Marine and Adjacent Riverine Areas of North Bohai Sea, China. <i>Archives of Environmental Contamination and Toxicology</i> , 2010, 59, 71-79.	2.1	41
314	Perfluorinated compounds in a coastal industrial area of Tianjin, China. <i>Environmental Geochemistry and Health</i> , 2012, 34, 301-311.	1.8	41
315	Perfluoroalkyl substances in soils around the Nepali Koshi River: levels, distribution, and mass balance. <i>Environmental Science and Pollution Research</i> , 2014, 21, 9201-9211.	2.7	41
316	Species- and tissue-specific bioaccumulation of arsenicals in various aquatic organisms from a highly industrialized area in the Pohang City, Korea. <i>Environmental Pollution</i> , 2014, 192, 27-35.	3.7	41
317	Sono-chemical treatment of per- and poly-fluoroalkyl compounds in aqueous film-forming foams by use of a large-scale multi-transducer dual-frequency based acoustic reactor. <i>Ultrasonics Sonochemistry</i> , 2018, 45, 213-222.	3.8	41
318	Endocrine disrupting, mutagenic, and teratogenic effects of upper Danube River sediments using effect-directed analysis. <i>Environmental Toxicology and Chemistry</i> , 2012, 31, 1053-1062.	2.2	40
319	Perfluorinated compounds and organochlorine pesticides in soils around Huaihe River: a heavily contaminated watershed in Central China. <i>Environmental Science and Pollution Research</i> , 2013, 20, 3965-3974.	2.7	40
320	Mechanisms of Toxicity of Hydroxylated Polybrominated Diphenyl Ethers (HO-PBDEs) Determined by Toxicogenomic Analysis with a Live Cell Array Coupled with Mutagenesis in <i>Escherichia coli</i> . <i>Environmental Science & Technology</i> , 2014, 48, 5929-5937.	4.6	40
321	Characterization of phosphorus forms in lake macrophytes and algae by solution ³¹ P nuclear magnetic resonance spectroscopy. <i>Environmental Science and Pollution Research</i> , 2016, 23, 7288-7297.	2.7	40
322	Spectroscopic analyses combined with Gaussian and Coats-Redfern models to investigate the characteristics and pyrolysis kinetics of sugarcane residue-derived biochars. <i>Journal of Cleaner Production</i> , 2019, 237, 117855.	4.6	40
323	Changes in phosphoadenylate concentrations and adenylate energy charge as an integrated biochemical measure of stress in invertebrates: The effects of cadmium on the freshwater clam <i>Corbicula fluminea</i> . <i>Toxicological and Environmental Chemistry</i> , 1983, 6, 259-295.	0.6	39
324	Contaminants in fishes from Great Lakes-influenced sections and above dams of three Michigan rivers. I: Concentrations of organo chlorine insecticides, polychlorinated biphenyls, dioxin equivalents, and mercury. <i>Archives of Environmental Contamination and Toxicology</i> , 1994, 27, 202-12.	2.1	39

#	ARTICLE	IF	CITATIONS
325	Low Reproductive Rates of Lake Superior Bald Eagles: Low Food Delivery Rates or Environmental Contaminants?. <i>Journal of Great Lakes Research</i> , 1998, 24, 32-44.	0.8	39
326	Specific binding of hydroxylated polychlorinated biphenyl metabolites and other substances to bovine calf uterine estrogen receptor: structure-binding relationships. <i>Science of the Total Environment</i> , 1999, 233, 141-161.	3.9	39
327	In vitro response of fish and mammalian cells to complex mixtures of polychlorinated naphthalenes, polychlorinated biphenyls, and polycyclic aromatic hydrocarbons. <i>Aquatic Toxicology</i> , 2001, 54, 125-141.	1.9	39
328	In Vitro Bioassay Determination of Dioxin-Like and Estrogenic Activity in Sediment and Water from Ulsan Bay and Its Vicinity, Korea. <i>Archives of Environmental Contamination and Toxicology</i> , 2001, 40, 151-160.	2.1	39
329	Chlorpyrifos: Ecotoxicological Risk Assessment for Birds and Mammals in Corn Agroecosystems. <i>Human and Ecological Risk Assessment (HERA)</i> , 2001, 7, 497-632.	1.7	39
330	In ovo exposure to o,p -DDE affects sexual development but not sexual differentiation in Japanese medaka (<i>Oryzias latipes</i>).. <i>Environmental Health Perspectives</i> , 2003, 111, 29-32.	2.8	39
331	Measurement of estrogenic activity in sediments from Haihe and Dagu River, China. <i>Environment International</i> , 2006, 32, 676-681.	4.8	39
332	Perfluorooctane Sulfonate Increases the Genotoxicity of Cyclophosphamide in the Micronucleus Assay with V79 Cells: Further Proof of Alterations in Cell Membrane Properties Caused by PFOS (3 pp). <i>Environmental Science and Pollution Research</i> , 2007, 14, 85-87.	2.7	39
333	Exposure of Hong Kong residents to PBDEs and their structural analogues through market fish consumption. <i>Journal of Hazardous Materials</i> , 2011, 192, 374-80.	6.5	39
334	Toxicogenomic Mechanisms of 6-HO-BDE-47, 6-MeO-BDE-47, and BDE-47 in <i>E. coli</i> . <i>Environmental Science & Technology</i> , 2012, 46, 1185-1191.	4.6	39
335	Time-dependent relative potency factors for polycyclic aromatic hydrocarbons and their derivatives in the H4IIE luc bioassay. <i>Environmental Toxicology and Chemistry</i> , 2014, 33, 943-953.	2.2	39
336	Bioaccessibility of AhR-active PAHs in sediments contaminated by the Hebei Spirit oil spill: Application of Tenax extraction in effect-directed analysis. <i>Chemosphere</i> , 2016, 144, 706-712.	4.2	39
337	Potential health risks posed by polycyclic aromatic hydrocarbons in muscle tissues of fishes from the Athabasca and Slave Rivers, Canada. <i>Environmental Geochemistry and Health</i> , 2017, 39, 139-160.	1.8	39
338	Influence of blooms of phytoplankton on concentrations of hydrophobic organic chemicals in sediments and snails in a hyper-eutrophic, freshwater lake. <i>Water Research</i> , 2017, 113, 22-31.	5.3	39
339	Glucuronide and sulfate conjugates of tetrabromobisphenol A (TBBPA): Chemical synthesis and correlation between their urinary levels and plasma TBBPA content in voluntary human donors. <i>Environment International</i> , 2017, 98, 46-53.	4.8	39
340	Selenium Bioaccumulation and Hazards in a Fish Community Affected by Coal Fly Ash Effluent. <i>Ecotoxicology and Environmental Safety</i> , 1996, 35, 7-15.	2.9	38
341	Polychlorinated biphenyls, organochlorine pesticides, tris(4-chlorophenyl)methane, and tris(4-chlorophenyl)methanol in livers of small cetaceans stranded along Florida coastal waters, USA. <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 1566-1574.	2.2	38
342	Concentrations and Profiles of Polychlorinated Dibenzo-p-Dioxins and Dibenzofurans in Soils from Korea. <i>Environmental Science & Technology</i> , 2002, 36, 3700-3705.	4.6	38

#	ARTICLE	IF	CITATIONS
343	Alteration of steroidogenesis in H295R cells by organic sediment contaminants and relationships to other endocrine disrupting effects. <i>Environment International</i> , 2006, 32, 749-757.	4.8	38
344	Effects of fluorotelomer alcohol 8:2 FTOH on steroidogenesis in H295R cells: Targeting the cAMP signalling cascade. <i>Toxicology and Applied Pharmacology</i> , 2010, 247, 222-228.	1.3	38
345	Chronic exposure to dietary selenomethionine increases gonadal steroidogenesis in female rainbow trout. <i>Aquatic Toxicology</i> , 2011, 105, 218-226.	1.9	38
346	Photolytic Degradation Products of Two Highly Brominated Flame Retardants Cause Cytotoxicity and mRNA Expression Alterations in Chicken Embryonic Hepatocytes. <i>Environmental Science & Technology</i> , 2014, 48, 12039-12046.	4.6	38
347	Occurrence of additive brominated flame retardants in aquatic organisms from Tai Lake and Yangtze River in Eastern China, 2009-2012. <i>Chemosphere</i> , 2014, 114, 340-346.	4.2	38
348	Dose-dependent compensation responses of the hypothalamic-pituitary-gonadal-liver axis of zebrafish exposed to the fungicide prochloraz. <i>Aquatic Toxicology</i> , 2015, 160, 69-75.	1.9	38
349	Endocrine disrupting potential of PAHs and their alkylated analogues associated with oil spills. <i>Environmental Sciences: Processes and Impacts</i> , 2017, 19, 1117-1125.	1.7	38
350	Genomic instability in adult men involved in processing electronic waste in Northern China. <i>Environment International</i> , 2018, 117, 69-81.	4.8	38
351	Characterization of phosphorus forms in a Eutrophic Lake, China. <i>Science of the Total Environment</i> , 2019, 659, 1437-1447.	3.9	38
352	Relationships Among Concentrations of Individual Polychlorinated Biphenyl (PCB) Congeners, 2,3,7,8-Tetrachlorodibenzo-P-Dioxin Equivalents (TCDD-EQ), and Rearing Mortality of Chinook Salmon (<i>Oncorhynchus Tshawytscha</i>) Eggs from Lake Michigan. <i>Journal of Great Lakes Research</i> , 1992, 18, 108-124.	0.8	37
353	Distribution and source apportionments of polychlorinated biphenyls (PCBs) in mariculture sediments from the Pearl River Delta, South China. <i>Marine Pollution Bulletin</i> , 2011, 63, 516-522.	2.3	37
354	In vitro profiling of endocrine disrupting potency of 2,2,4,4-tetrabromodiphenyl ether (BDE47) and related hydroxylated analogs (HO-PBDEs). <i>Marine Pollution Bulletin</i> , 2011, 63, 287-296.	2.3	37
355	Revised relative potency values for PCDDs, PCDFs, and non-ortho-substituted PCBs for the optimized H4IIe-luc in vitro bioassay. <i>Environmental Science and Pollution Research</i> , 2013, 20, 8590-8599.	2.7	37
356	Concentrations and congener profiles of polybrominated diphenyl ethers (PBDEs) in blood plasma from Hong Kong: Implications for sources and exposure route. <i>Journal of Hazardous Materials</i> , 2013, 261, 253-259.	6.5	37
357	Identification of Thyroid Hormone Disruptors among HO-PBDEs: <i>In Vitro</i> Investigations and Coregulator Involved Simulations. <i>Environmental Science & Technology</i> , 2016, 50, 12429-12438.	4.6	37
358	Characteristics and degradation of carbon and phosphorus from aquatic macrophytes in lakes: Insights from solid-state ¹³ C NMR and solution ³¹ P NMR spectroscopy. <i>Science of the Total Environment</i> , 2016, 543, 746-756.	3.9	37
359	Pore-level visual analysis of heavy oil recovery using chemical-assisted waterflooding process - Use of a new chemical agent. <i>Fuel</i> , 2019, 239, 202-218.	3.4	37
360	Differential responses of gut microbiota of male and female fathead minnow (<i>Pimephales promelas</i>) to a short-term environmentally-relevant, aqueous exposure to benzo[a]pyrene. <i>Chemosphere</i> , 2020, 252, 126461.	4.2	37

#	ARTICLE	IF	CITATIONS
361	Fighting against the second wave of COVID-19: Can honeybee products help protect against the pandemic?. Saudi Journal of Biological Sciences, 2021, 28, 1519-1527.	1.8	37
362	Microfibers Released into the Air from a Household Tumble Dryer. Environmental Science and Technology Letters, 2022, 9, 120-126.	3.9	37
363	Concentrations and Hazard Assessment of Organochlorine Contaminants and Mercury in Smallmouth Bass from a Remote Lake in the Upper Peninsula of Michigan. Archives of Environmental Contamination and Toxicology, 1998, 34, 81-86.	2.1	36
364	Vertical Profiles of Dioxin-like and Estrogenic Activities Associated with a Sediment Core from Tokyo Bay, Japan. Environmental Science & Technology, 2000, 34, 3568-3573.	4.6	36
365	Identification and quantitation method for nonylphenol and lower oligomer nonylphenol ethoxylates in fish tissues. Environmental Toxicology and Chemistry, 2001, 20, 1870-1873.	2.2	36
366	Differential Accumulation of Polychlorinated Biphenyl Congeners in the Aquatic Food Web at the Kalamazoo River Superfund Site, Michigan. Environmental Science & Technology, 2005, 39, 5964-5974.	4.6	36
367	Bioaccumulation of polychlorinated dibenzo-p-dioxins, dibenzofurans, and dioxin-like polychlorinated biphenyls in fishes from the Tittabawassee and Saginaw Rivers, Michigan, USA. Science of the Total Environment, 2010, 408, 2394-2401.	3.9	36
368	Analytical and bioanalytical assessments of organic micropollutants in the Bosna River using a combination of passive sampling, bioassays and multi-residue analysis. Science of the Total Environment, 2019, 650, 1599-1612.	3.9	36
369	Sublethal effects of chronic exposure to CdO or PbO nanoparticles or their binary mixture on the honey bee (<i>Apis mellifera</i> L.). Environmental Science and Pollution Research, 2020, 27, 19004-19015.	2.7	36
370	Trends of Contaminants and Effects in Bald Eagles of the Great Lakes Basin. Environmental Monitoring and Assessment, 1998, 53, 197-212.	1.3	35
371	Perfluorinated compounds in water and sediment from coastal regions of the northern Bohai Sea, China. Chemistry and Ecology, 2011, 27, 165-176.	0.6	35
372	Effects of exposure to 17 β -ethynylestradiol during larval development on growth, sexual differentiation, and abundances of transcripts in the liver of the wood frog (<i>Lithobates sylvaticus</i>). Aquatic Toxicology, 2013, 126, 42-51.	1.9	35
373	Bioassay-directed identification of organic toxicants in water and sediment of Tai Lake, China. Water Research, 2015, 73, 231-241.	5.3	35
374	Biological plausibility as a tool to associate analytical data for micropollutants and effect potentials in wastewater, surface water, and sediments with effects in fishes. Water Research, 2015, 72, 127-144.	5.3	35
375	High Conservation in Transcriptomic and Proteomic Response of White Sturgeon to Equipotent Concentrations of 2,3,7,8-TCDD, PCB 77, and Benzo[a]pyrene. Environmental Science & Technology, 2016, 50, 4826-4835.	4.6	35
376	Toxicity of Detroit River Sediment Interstitial Water to the Bacterium <i>Photobacterium Phosphoreum</i> . Journal of Great Lakes Research, 1988, 14, 502-513.	0.8	34
377	Toxic responses of medaka, D \times R strain, to polychlorinatednaphthalene mixtures after embryonic exposure by in ovo nanoinjection: A partial life cycle assessment. Environmental Toxicology and Chemistry, 2000, 19, 432-440.	2.2	34
378	Pollutants in particulate and gaseous fractions of ambient air interfere with multiple signaling pathways in vitro. Environment International, 2009, 35, 43-49.	4.8	34

#	ARTICLE	IF	CITATIONS
379	Population-specific incidence of testicular ovarian follicles in <i>Xenopus laevis</i> from South Africa: A potential issue in endocrine testing. <i>Aquatic Toxicology</i> , 2009, 95, 10-16.	1.9	34
380	Extinction Risk of Exploited Wild Roach (<i>Rutilus rutilus</i>) Populations Due to Chemical Feminization. <i>Environmental Science & Technology</i> , 2009, 43, 7895-7901.	4.6	34
381	Attenuation of the cortisol response to stress in female rainbow trout chronically exposed to dietary selenomethionine. <i>Aquatic Toxicology</i> , 2011, 105, 643-651.	1.9	34
382	Composition and effects of inhalable size fractions of atmospheric aerosols in the polluted atmosphere. Part II. In vitro biological potencies. <i>Environment International</i> , 2014, 63, 64-70.	4.8	34
383	Effects of Tris(1,3-dichloro-2-propyl) Phosphate (TDCPP) in <i>Tetrahymena Thermophila</i> : Targeting the Ribosome. <i>Scientific Reports</i> , 2015, 5, 10562.	1.6	34
384	A mixture of the novel brominated flame retardants TBPH and TBB affects fecundity and transcript profiles of the HPGL-axis in Japanese medaka. <i>Aquatic Toxicology</i> , 2015, 158, 14-21.	1.9	34
385	Fluorescence regional integration and differential fluorescence spectroscopy for analysis of structural characteristics and proton binding properties of fulvic acid sub-fractions. <i>Journal of Environmental Sciences</i> , 2018, 74, 116-125.	3.2	34
386	Spatial and temporal trends in poly- and per-fluorinated compounds in the Laurentian Great Lakes Erie, Ontario and St. Clair. <i>Environmental Pollution</i> , 2018, 237, 396-405.	3.7	34
387	PM2.5 bound phthalates in four metropolitan cities of China: Concentration, seasonal pattern and health risk via inhalation. <i>Science of the Total Environment</i> , 2019, 696, 133982.	3.9	34
388	Antioxidant and Antimicrobial Activities of Chemically-Characterized Essential Oil from <i>Artemisia aragonensis</i> Lam. against Drug-Resistant Microbes. <i>Molecules</i> , 2022, 27, 1136.	1.7	34
389	Bicarbonate as a Potential Confounding Factor in Cladoceran Toxicity Assessments of Pore Water from Contaminated Sediments. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1992, 49, 1633-1640.	0.7	33
390	Induction of erod activity in HEPA mouse hepatoma cells and estrogenicity in mcf7 human breast cancer cells by extracts of pulp mill effluents, sludge, and sediment exposed to effluents. <i>Environmental Toxicology and Chemistry</i> , 1998, 17, 1499-1507.	2.2	33
391	SCRAM: A Scoring and Ranking System for Persistent, Bioaccumulative, and Toxic Substances for the North American Great Lakes-Resulting Chemical Scores and Rankings. <i>Human and Ecological Risk Assessment (HERA)</i> , 2002, 8, 537-557.	1.7	33
392	Characterization of Estrogenic Activity of Riverine Sediments from the Czech Republic. <i>Archives of Environmental Contamination and Toxicology</i> , 2002, 43, 175-185.	2.1	33
393	Assessment of laryngeal muscle and testicular cell types in <i>Xenopus laevis</i> (Anura Pipidae) inhabiting maize and non-maize growing areas of South Africa. <i>African Journal of Herpetology</i> , 2005, 54, 69-76.	0.3	33
394	Modulation of steroidogenic gene expression and hormone synthesis in H295R cells exposed to PCP and TCP. <i>Toxicology</i> , 2011, 282, 146-153.	2.0	33
395	Predicting criteria continuous concentrations of 34 metals or metalloids by use of quantitative ion character-activity relationships species sensitivity distributions (QICAR-SSD) model. <i>Environmental Pollution</i> , 2014, 188, 50-55.	3.7	33
396	End-of-life (EoL) mobile phone management in Hong Kong households. <i>Journal of Environmental Management</i> , 2017, 200, 22-28.	3.8	33

#	ARTICLE	IF	CITATIONS
397	Mechanisms of oxidative stress caused by CuO nanoparticles to membranes of the bacterium <i>Streptomyces coelicolor</i> M145. <i>Ecotoxicology and Environmental Safety</i> , 2018, 158, 123-130.	2.9	33
398	Exposure to organophosphate esters in elderly people: Relationships of OPE body burdens with indoor air and dust concentrations and food consumption. <i>Environment International</i> , 2021, 157, 106803.	4.8	33
399	UO ₂ ²⁺ -humate interactions in soft, acid, humate-rich waters. <i>Journal of Environmental Radioactivity</i> , 1986, 4, 39-64.	0.9	32
400	Butyltin Compounds in River Otters (<i>Lutra canadensis</i>) from the Northwestern United States. <i>Archives of Environmental Contamination and Toxicology</i> , 1999, 36, 462-468.	2.1	32
401	Endocrine effects of methoxylated brominated diphenyl ethers in three in vitro models. <i>Marine Pollution Bulletin</i> , 2011, 62, 2356-2361.	2.3	32
402	Differences in Activation of Aryl Hydrocarbon Receptors of White Sturgeon Relative to Lake Sturgeon Are Predicted by Identities of Key Amino Acids in the Ligand Binding Domain. <i>Environmental Science & Technology</i> , 2015, 49, 4681-4689.	4.6	32
403	Linking Oxidative Stress and Magnitude of Compensatory Responses with Life-Stage Specific Differences in Sensitivity of White Sturgeon (<i>Acipenser transmontanus</i>) to Copper or Cadmium. <i>Environmental Science & Technology</i> , 2016, 50, 9717-9726.	4.6	32
404	Model for Predicting Toxicities of Metals and Metalloids in Coastal Marine Environments Worldwide. <i>Environmental Science & Technology</i> , 2018, 52, 4199-4206.	4.6	32
405	Amendment of water quality standards in China: viewpoint on strategic considerations. <i>Environmental Science and Pollution Research</i> , 2018, 25, 3078-3092.	2.7	32
406	Anthracene bioconcentration and biotransformation in chironomids: Effects of temperature and concentration. <i>Environmental Pollution Series A, Ecological and Biological</i> , 1983, 30, 175-188.	0.8	31
407	Extractable Organohalogen (EOX) in Sediment and Biota Collected at an Estuarine Marsh near a Former Chloralkali Facility. <i>Environmental Science & Technology</i> , 1999, 33, 1004-1008.	4.6	31
408	Associations between regional differences in polychlorinated biphenyls and dichlorodiphenyldichloroethylene in blood of nestling bald eagles and reproductive productivity. <i>Environmental Toxicology and Chemistry</i> , 2003, 22, 371-376.	2.2	31
409	Leptin-Mediated Modulation of Steroidogenic Gene Expression in Hypoxic Zebrafish Embryos: Implications for the Disruption of Sex Steroids. <i>Environmental Science & Technology</i> , 2012, 46, 9112-9119.	4.6	31
410	Changes in concentrations of hydrophilic organic contaminants and of endocrine-disrupting potential downstream of small communities located adjacent to headwaters. <i>Environment International</i> , 2012, 45, 22-31.	4.8	31
411	Effect-directed analysis: Current status and future challenges. <i>Ocean Science Journal</i> , 2016, 51, 413-433.	0.6	31
412	Inhibition of ABC transport proteins by oil sands process affected water. <i>Aquatic Toxicology</i> , 2016, 170, 81-88.	1.9	31
413	Immunotoxicity of aflatoxin M ₁ : as a potent suppressor of innate and acquired immune systems in a subacute study. <i>Journal of the Science of Food and Agriculture</i> , 2018, 98, 5884-5892.	1.7	31
414	Multiple Bioassays and Targeted and Nontargeted Analyses to Characterize Potential Toxicological Effects Associated with Sediments of Masan Bay: Focusing on AhR-Mediated Potency. <i>Environmental Science & Technology</i> , 2020, 54, 4443-4454.	4.6	31

#	ARTICLE	IF	CITATIONS
415	Dietary exposure of mink to carp from Saginaw Bay, Michigan: 2. Hematology and liver pathology. Archives of Environmental Contamination and Toxicology, 1995, 29, 411-417.	2.1	30
416	Population structure of the African Clawed Frog (<i>Xenopus laevis</i>) in maize-growing areas with atrazine application versus non-maize-growing areas in South Africa. African Journal of Herpetology, 2005, 54, 61-68.	0.3	30
417	AhR-active compounds in sediments of the Haihe and Dagu Rivers, China. Chemosphere, 2006, 63, 1222-1230.	4.2	30
418	Measuring and monitoring persistent organic pollutants in the context of risk assessment. Marine Pollution Bulletin, 2008, 57, 236-244.	2.3	30
419	Endocrine effects of contaminated sediments on the freshwater snail <i>Potamopyrgus antipodarum</i> in vivo and in the cell bioassays in vitro. Aquatic Toxicology, 2008, 89, 172-179.	1.9	30
420	In vitro modulation of intracellular receptor signaling and cytotoxicity induced by extracts of cyanobacteria, complex water blooms and their fractions. Aquatic Toxicology, 2011, 105, 497-507.	1.9	30
421	Contribution of Priority PAHs and POPs to Ah Receptor-Mediated Activities in Sediment Samples from the River Elbe Estuary, Germany. PLoS ONE, 2013, 8, e75596.	1.1	30
422	Seasonal concentrations of lead in outdoor and indoor dust and blood of children in Riyadh, Saudi Arabia. Environmental Geochemistry and Health, 2014, 36, 583-593.	1.8	30
423	Isolation and Characterization of Chinese Standard Fulvic Acid Sub-fractions Separated from Forest Soil by Stepwise Elution with Pyrophosphate Buffer. Scientific Reports, 2015, 5, 8723.	1.6	30
424	Untargeted Screening and Distribution of Organo-Iodine Compounds in Sediments from Lake Michigan and the Arctic Ocean. Environmental Science & Technology, 2016, 50, 10097-10105.	4.6	30
425	Polycyclic aromatic hydrocarbons in infant formulae, follow-on formulae, and baby foods in Iran: An assessment of risk. Food and Chemical Toxicology, 2019, 131, 110640.	1.8	30
426	Short-chain chlorinated paraffins (SCCPs) disrupt hepatic fatty acid metabolism in liver of male rat via interacting with peroxisome proliferator-activated receptor α (PPAR α). Ecotoxicology and Environmental Safety, 2019, 181, 164-171.	2.9	30
427	Using the carotenoid biosynthesis inhibiting herbicide, Fluridone, to investigate the ability of carotenoid pigments to protect algae from the photo-induced toxicity of anthracene. Aquatic Toxicology, 1993, 27, 61-70.	1.9	29
428	TREE SWALLOW (TACHYGINETA BICOLOR) EXPOSURE TO POLYCHLORINATED BIPHENYLS AT THE KALAMAZOO RIVER SUPERFUND SITE, MICHIGAN, USA. Environmental Toxicology and Chemistry, 2006, 25, 428.	2.2	29
429	Regulation of CYP11B1 and CYP11B2 steroidogenic genes by hypoxia-inducible miR-10b in H295R cells. Marine Pollution Bulletin, 2014, 85, 344-351.	2.3	29
430	Acute toxicity of copper, lead, cadmium, and zinc to early life stages of white sturgeon (<i>Acipenser</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Research, 2014, 21, 8176-8187.	2.7	29
431	Identification and expression of aryl hydrocarbon receptors (AhR1 and AhR2) provide insight in an evolutionary context regarding sensitivity of white sturgeon (<i>Acipenser transmontanus</i>) to dioxin-like compounds. Aquatic Toxicology, 2014, 150, 27-35.	1.9	29
432	In Vitro Assessment of Endocrine Disrupting Potential of Naphthenic Acid Fractions Derived from Oil Sands-Influenced Water. Environmental Science & Technology, 2015, 49, 5743-5752.	4.6	29

#	ARTICLE	IF	CITATIONS
433	Maternal transfer, distribution, and metabolism of BDE-47 and its related hydroxylated, methoxylated analogs in zebrafish (<i>Danio rerio</i>). <i>Chemosphere</i> , 2015, 120, 31-36.	4.2	29
434	Hazard posed by metals and As in PM _{2.5} in air of five megacities in the Beijing-Tianjin-Hebei region of China during APEC. <i>Environmental Science and Pollution Research</i> , 2016, 23, 17603-17612.	2.7	29
435	Chemical-, site-, and taxa-dependent benthic community health in coastal areas of the Bohai Sea and northern Yellow Sea: A sediment quality triad approach. <i>Science of the Total Environment</i> , 2018, 645, 743-752.	3.9	29
436	Application of the Target Lipid Model and Passive Samplers to Characterize the Toxicity of Bioavailable Organics in Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2018, 52, 8039-8049.	4.6	29
437	Effects of acute exposure to microcystins on hypothalamic-pituitary-adrenal (HPA), -gonad (HPG) and -thyroid (HPT) axes of female rats. <i>Science of the Total Environment</i> , 2021, 778, 145196.	3.9	29
438	TOXIC RESPONSES OF MEDAKA, d-rR STRAIN, TO POLYCHLORINATED NAPHTHALENE MIXTURES AFTER EMBRYONIC EXPOSURE BY IN OVO NANOINJECTION: A PARTIAL LIFE-CYCLE ASSESSMENT. <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 432.	2.2	29
439	A Comparison of Water Quality Criteria for the Great Lakes Based on Human and Wildlife Health. <i>Journal of Great Lakes Research</i> , 1993, 19, 789-807.	0.8	28
440	Accumulation of 2,3,7,8-tetrachlorodibenzo-p-dioxin by rainbow trout (<i>Onchorhynchus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 3 <i>Chemistry</i> , 2001, 20, 344-350.	2.2	28
441	Toxaphene and other persistent organochlorine pesticides in three species of albatrosses from the north and south Pacific Ocean. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 413-423.	2.2	28
442	Effects of bisphenol A-related diphenylalkanes on vitellogenin production in male carp (<i>Cyprinus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 3 <i>Toxicology and Applied Pharmacology</i> , 2005, 209, 95-104.	1.3	28
443	Persistent halogenated compounds in aquaculture environments of South China: Implications for global consumers' health risk via fish consumption. <i>Environment International</i> , 2011, 37, 1190-1195.	4.8	28
444	Measured and predicted affinities of binding and relative potencies to activate the AhR of PAHs and their alkylated analogues. <i>Chemosphere</i> , 2015, 139, 23-29.	4.2	28
445	Probabilistic ecological risk assessment of copper in Chinese offshore marine environments from 2005 to 2012. <i>Marine Pollution Bulletin</i> , 2015, 94, 96-102.	2.3	28
446	Beekeeping and the Need for Pollination from an Agricultural Perspective in Egypt. <i>Bee World</i> , 2018, 95, 107-112.	0.3	28
447	<i>Lavandula dentata</i> L.: Phytochemical Analysis, Antioxidant, Antifungal and Insecticidal Activities of Its Essential Oil. <i>Plants</i> , 2022, 11, 311.	1.6	28
448	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 52-61.	2.7	27
449	Effects of Primary Exposure to Environmental and Natural Estrogens on Vitellogenin Production in Carp (<i>Cyprinus carpio</i>) Hepatocytes. <i>Toxicological Sciences</i> , 2002, 67, 75-80.	1.4	27
450	Organochlorines and dioxin-like compounds in green-lipped mussels <i>Perna viridis</i> from Hong Kong mariculture zones. <i>Marine Pollution Bulletin</i> , 2005, 51, 677-687.	2.3	27

#	ARTICLE	IF	CITATIONS
451	Receptor-mediated in vitro bioassay for characterization of Ah-R-active compounds and activities in sediment from Korea. <i>Chemosphere</i> , 2006, 62, 1261-1271.	4.2	27
452	Organochlorine pesticides in soils around Guanting Reservoir, China. <i>Environmental Geochemistry and Health</i> , 2007, 29, 491-501.	1.8	27
453	Polybrominated diphenyl ethers and their methoxylated metabolites in anchovy (<i>Coilia sp.</i>) from the Yangtze River Delta, China. <i>Environmental Science and Pollution Research</i> , 2010, 17, 634-642.	2.7	27
454	PBDEs and methoxylated analogues in sediment cores from two Michigan, USA, inland lakes. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 1236-1242.	2.2	27
455	A comparison of statistical methods for deriving freshwater quality criteria for the protection of aquatic organisms. <i>Environmental Science and Pollution Research</i> , 2014, 21, 159-167.	2.7	27
456	Can zero-valent iron nanoparticles remove waterborne estrogens?. <i>Journal of Environmental Management</i> , 2015, 150, 387-392.	3.8	27
457	Spatial and temporal distribution and sources of polycyclic aromatic hydrocarbons in sediments of Taihu Lake, eastern China. <i>Environmental Science and Pollution Research</i> , 2015, 22, 5350-5358.	2.7	27
458	Identification of Chemicals that Cause Oxidative Stress in Oil Sands Process-Affected Water. <i>Environmental Science & Technology</i> , 2017, 51, 8773-8781.	4.6	27
459	Exposure to Al ₂ O ₃ nanoparticles facilitates conjugative transfer of antibiotic resistance genes from <i>Escherichia coli</i> to <i>Streptomyces</i> . <i>Nanotoxicology</i> , 2019, 13, 1422-1436.	1.6	27
460	Metals and PFAS in stormwater and surface runoff in a semi-arid Canadian city subject to large variations in temperature among seasons. <i>Environmental Science and Pollution Research</i> , 2020, 27, 18232-18241.	2.7	27
461	Particulate formation due to freezing humic waters. <i>Water Resources Research</i> , 1978, 14, 542-544.	1.7	26
462	In vitro mitogenesis of peripheral blood lymphocytes from rainbow trout (<i>Salmo gairdneri</i>). <i>Comparative Biochemistry and Physiology A, Comparative Physiology</i> , 1988, 89, 25-35.	0.7	26
463	Persistent organochlorine pollutants in eggs of colonial waterbirds from Galveston Bay and East Texas, USA. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 608-617.	2.2	26
464	Plasma steroid hormone concentrations, aromatase activities and GSI in ranid frogs collected from agricultural and non-agricultural sites in Michigan (USA). <i>Aquatic Toxicology</i> , 2006, 77, 153-166.	1.9	26
465	Fluorescence in situ hybridization techniques (FISH) to detect changes in CYP19a gene expression of Japanese medaka (<i>Oryzias latipes</i>). <i>Toxicology and Applied Pharmacology</i> , 2008, 232, 226-235.	1.3	26
466	Effects of subchronic exposure of early life stages of white sturgeon (<i>Acipenser</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 Td (transn 2497-2505.	2.2	26
467	Polycyclic aromatic hydrocarbons in soils along the coastal and estuarine areas of the northern Bohai and Yellow Seas, China. <i>Environmental Monitoring and Assessment</i> , 2013, 185, 8185-8195.	1.3	26
468	Effects of dechlorane plus on the hepatic proteome of juvenile Chinese sturgeon (<i>Acipenser sinensis</i>). <i>Aquatic Toxicology</i> , 2014, 148, 83-91.	1.9	26

#	ARTICLE	IF	CITATIONS
469	Comparison on the molecular response profiles between nano zinc oxide (ZnO) particles and free zinc ion using a genome-wide toxicogenomics approach. <i>Environmental Science and Pollution Research</i> , 2015, 22, 17434-17442.	2.7	26
470	Long-term spatial trends in sedimentary algal pigments in a narrow river-valley reservoir, Lake Diefenbaker, Canada. <i>Journal of Great Lakes Research</i> , 2015, 41, 56-66.	0.8	26
471	Effects of environmentally-relevant mixtures of four common organophosphorus insecticides on the honey bee (<i>Apis mellifera</i> L.). <i>Journal of Insect Physiology</i> , 2015, 82, 85-91.	0.9	26
472	Effect of Lipid Partitioning on Predictions of Acute Toxicity of Oil Sands Process Affected Water to Embryos of Fathead Minnow (<i>Pimephales promelas</i>). <i>Environmental Science & Technology</i> , 2016, 50, 8858-8866.	4.6	26
473	Bioconcentration of Dissolved Organic Compounds from Oil Sands Process-Affected Water by Medaka (<i>Oryzias latipes</i>): Importance of Partitioning to Phospholipids. <i>Environmental Science & Technology</i> , 2016, 50, 6574-6582.	4.6	26
474	Bioaccumulation of Polycyclic Aromatic Hydrocarbons (PAHs) by the Marine Clam, <i>Macraa veneriformis</i> , Chronically Exposed to Oil-Suspended Particulate Matter Aggregates. <i>Environmental Science & Technology</i> , 2018, 52, 7910-7920.	4.6	26
475	Extraction efficiency of anthracene from sediments. <i>Analytical Chemistry</i> , 1983, 55, 1197-1200.	3.2	25
476	Accumulation of 2,3,7,8-Tetrachlorodibenzo-p-dioxin Equivalents by Double-Crested Cormorant (<i>Phalacrocorax auritus</i> , Pelicaniformes) Chicks in the North American Great Lakes. <i>Ecotoxicology and Environmental Safety</i> , 1994, 27, 192-209.	2.9	25
477	Rainbow trout cell bioassay-derived relative potencies for halogenated aromatic hydrocarbons: Comparison and sensitivity analysis. <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 879-888.	2.2	25
478	Effects of 4-nonylphenol on fecundity and biomarkers of estrogenicity in fathead minnows (<i>Pimephales promelas</i>). <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 1368-1377.	2.2	25
479	Docking and CoMSIA studies on steroids and non-steroidal chemicals as androgen receptor ligands. <i>Ecotoxicology and Environmental Safety</i> , 2013, 89, 143-149.	2.9	25
480	Microalga <i>Euglena</i> as a bioindicator for testing genotoxic potentials of organic pollutants in Taihu Lake, China. <i>Ecotoxicology</i> , 2014, 23, 633-640.	1.1	25
481	Effects of the brominated flame retardant TBCO on fecundity and profiles of transcripts of the HPGL-axis in Japanese medaka. <i>Aquatic Toxicology</i> , 2015, 160, 180-187.	1.9	25
482	Glucuronide and Sulfate Conjugates of Bisphenol A: Chemical Synthesis and Correlation Between Their Urinary Levels and Plasma Bisphenol A Content in Voluntary Human Donors. <i>Archives of Environmental Contamination and Toxicology</i> , 2017, 73, 410-420.	2.1	25
483	Accumulation rates, focusing factors, and chronologies from depth profiles of ²¹⁰ Pb and ¹³⁷ Cs in sediments of the Laurentian Great Lakes. <i>Journal of Great Lakes Research</i> , 2018, 44, 693-704.	0.8	25
484	Three decades of changes in water environment of a large freshwater Lake and its relationship with socio-economic indicators. <i>Journal of Environmental Sciences</i> , 2019, 77, 156-166.	3.2	25
485	Major AhR-active chemicals in sediments of Lake Sihwa, South Korea: Application of effect-directed analysis combined with full-scan screening analysis. <i>Environment International</i> , 2019, 133, 105199.	4.8	25
486	Consequences of a short-term exposure to a sub lethal concentration of CdO nanoparticles on key life history traits in the fruit fly (<i>Drosophila melanogaster</i>). <i>Journal of Hazardous Materials</i> , 2021, 410, 124671.	6.5	25

#	ARTICLE	IF	CITATIONS
487	Polycyclic aromatic hydrocarbons, pesticides, and metals in olive: analysis and probabilistic risk assessment. <i>Environmental Science and Pollution Research</i> , 2021, 28, 39723-39741.	2.7	25
488	Curcumin protects against tartrazine-mediated oxidative stress and hepatotoxicity in male rats. <i>European Review for Medical and Pharmacological Sciences</i> , 2017, 21, 635-645.	0.5	25
489	Effects of chronic cadmium exposure on crayfish survival, growth, and tolerance to elevated temperatures. <i>Archives of Environmental Contamination and Toxicology</i> , 1979, 8, 449-456.	2.1	24
490	The histological and biochemical effects of cadmium exposure in the bluegill sunfish (<i>Lepomis</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622	2.9	24
491	Toward Identifying the Next Generation of Superfund and Hazardous Waste Site Contaminants. <i>Environmental Health Perspectives</i> , 2011, 119, 6-10.	2.8	24
492	Equol Induces Gonadal Intersex in Japanese Medaka (<i>Oryzias latipes</i>) at Environmentally Relevant Concentrations: Comparison with 17 β -Estradiol. <i>Environmental Science & Technology</i> , 2016, 50, 7852-7860.	4.6	24
493	Pesticides, metals, and polycyclic aromatic hydrocarbons in date fruits: A probabilistic assessment of risk to health of Iranian consumers. <i>Journal of Food Composition and Analysis</i> , 2021, 98, 103815.	1.9	24
494	Occurrence of Butyltin Compounds in Tissues of Water Birds and Seaducks from the United States and Canada. <i>Archives of Environmental Contamination and Toxicology</i> , 1998, 35, 64-69.	2.1	23
495	Chapter 2 Emission, Contamination and Exposure, Fate and Transport, and National Management Strategy of Persistent Organic Pollutants in South Korea. <i>Developments in Environmental Science</i> , 2007, 7, 31-157.	0.5	23
496	Do water quality criteria based on nonnative species provide appropriate protection for native species?. <i>Environmental Toxicology and Chemistry</i> , 2015, 34, 1793-1798.	2.2	23
497	Chemical characterization and antioxidant properties of Canadian propolis. <i>Journal of Apicultural Research</i> , 2016, 55, 305-314.	0.7	23
498	Peroxisome Proliferator-Activated Receptor β 3 is a Sensitive Target for Oil Sands Process-Affected Water: Effects on Adipogenesis and Identification of Ligands. <i>Environmental Science & Technology</i> , 2016, 50, 7816-7824.	4.6	23
499	Effect of oil sands process-affected water on toxicity of retene to early life-stages of Japanese medaka (<i>Oryzias latipes</i>). <i>Aquatic Toxicology</i> , 2016, 176, 1-9.	1.9	23
500	Legacy polychlorinated organic pollutants in the sediment of the Great Lakes. <i>Journal of Great Lakes Research</i> , 2018, 44, 682-692.	0.8	23
501	Current understanding of potential ecological risks of retinoic acids and their metabolites in aquatic environments. <i>Environment International</i> , 2020, 136, 105464.	4.8	23
502	STIMULATION OF GROWTH IN SCENEDESMUS OBLIQUUS (CHLOROPHYCEAE) BY HUMIC ACIDS UNDER IRON LIMITED CONDITIONS ^{1,2} . <i>Journal of Phycology</i> , 1976, 12, 172-179.	1.0	23
503	EXAMINATION OF REPRODUCTIVE ENDPOINTS IN GOLDFISH (<i>CARASSIUS AURATUS</i>) EXPOSED IN SITU TO MUNICIPAL SEWAGE TREATMENT PLANT EFFLUENT DISCHARGES IN MICHIGAN, USA. <i>Environmental Toxicology and Chemistry</i> , 2003, 22, 2416.	2.2	22
504	SQUAMOUS EPITHELIAL LESION OF THE MANDIBLES AND MAXILLAE OF WILD MINK (<i>MUSTELA VISON</i>) NATURALLY EXPOSED TO POLYCHLORINATED BIPHENYLS. <i>Environmental Toxicology and Chemistry</i> , 2005, 24, 674.	2.2	22

#	ARTICLE	IF	CITATIONS
505	Human Exposure to Dioxin-Like Compounds in Fish and Shellfish Consumed in South Korea. Human and Ecological Risk Assessment (HERA), 2007, 13, 223-235.	1.7	22
506	Enhancement of AhR-mediated activity of selected pollutants and their mixtures after interaction with dissolved organic matter. Environment International, 2011, 37, 960-964.	4.8	22
507	Protective effects of eicosapentaenoic acid on genotoxicity and oxidative stress of cyclophosphamide in mice. Environmental Toxicology, 2011, 26, 217-223.	2.1	22
508	Environmental concentrations and bioaccumulations of cadmium and zinc in coastal watersheds along the Chinese Northern Bohai and Yellow Seas. Environmental Toxicology and Chemistry, 2013, 32, 831-840.	2.2	22
509	AhR-mediated activities of polycyclic aromatic compound (PAC) mixtures are predictable by the concept of concentration addition. Environment International, 2014, 73, 94-103.	4.8	22
510	Functionality of Aryl Hydrocarbon Receptors (AhR1 and AhR2) of White Sturgeon (<i>Acipenser</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 5 Environmental Science & Technology, 2014, 48, 8219-8226.	4.6	22
511	Bioaccumulation and molecular effects of sediment-bound metals in zebrafish embryos. Environmental Science and Pollution Research, 2015, 22, 16290-16304.	2.7	22
512	Causes of endocrine disrupting potencies in surface water in East China. Chemosphere, 2016, 144, 1435-1442.	4.2	22
513	Products of biotransformation of polycyclic aromatic hydrocarbons in fishes of the Athabasca/Slave river system, Canada. Environmental Geochemistry and Health, 2016, 38, 577-591.	1.8	22
514	Elucidating mechanisms of toxic action of dissolved organic chemicals in oil sands process-affected water (OSPW). Chemosphere, 2017, 186, 893-900.	4.2	22
515	Combined effects of degradable film fragments and micro/nanoplastics on growth of wheat seedling and rhizosphere microbes. Environmental Pollution, 2022, 294, 118516.	3.7	22
516	Concentrations of Dissolved and Particulate Polychlorinated Biphenyls in Water from the Saginaw River, Michigan. Journal of Great Lakes Research, 1995, 21, 219-233.	0.8	21
517	Assessment of Sediment Quality in Dredged and Undredged Areas of the Trenton Channel of the Detroit River, Michigan USA, using the Sediment Quality Triad. Journal of Great Lakes Research, 1996, 22, 683-696.	0.8	21
518	PCBs in the Detroit River Water Column. Journal of Great Lakes Research, 1997, 23, 440-449.	0.8	21
519	Atmospheric Deposition and Fluxes of Organochlorine Pesticides and Coplanar Polychlorinated Biphenyls in Aquatic Environments of Hong Kong, China. Environmental Science & Technology, 2004, 38, 6513-6521.	4.6	21
520	Toxicity reference values for mink exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) equivalents (TEQs). Ecotoxicology and Environmental Safety, 2008, 69, 325-349.	2.9	21
521	Risk to humans of consuming metals in anchovy (<i>Coilia</i> sp.) from the Yangtze River Delta. Environmental Geochemistry and Health, 2009, 31, 727-740.	1.8	21
522	Synthesis and Characterization of Bromophenol Glucuronide and Sulfate Conjugates for Their Direct LC-MS/MS Quantification in Human Urine as Potential Exposure Markers for Polybrominated Diphenyl Ethers. Analytical Chemistry, 2012, 84, 9881-9888.	3.2	21

#	ARTICLE	IF	CITATIONS
523	Relationship between mercury and organic carbon in sediment cores from Lakes Qinghai and Chenghai, China. <i>Journal of Soils and Sediments</i> , 2013, 13, 1084-1092.	1.5	21
524	Differential modulation of expression of nuclear receptor mediated genes by tris(2-butoxyethyl) phosphate (TBOEP) on early life stages of zebrafish (<i>Danio rerio</i>). <i>Aquatic Toxicology</i> , 2015, 169, 196-203.	1.9	21
525	Evidence for MicroRNA-Mediated Regulation of Steroidogenesis by Hypoxia. <i>Environmental Science & Technology</i> , 2015, 49, 1138-1147.	4.6	21
526	Life cycle analysis of perfluorooctanoic acid (PFOA) and its salts in China. <i>Environmental Science and Pollution Research</i> , 2017, 24, 11254-11264.	2.7	21
527	Spatial distribution and hazard of halogenated flame retardants and polychlorinated biphenyls to common kingfisher (<i>Alcedo atthis</i>) from a region of South China affected by electronic waste recycling. <i>Environment International</i> , 2019, 130, 104952.	4.8	21
528	Tissue distribution, bioaccumulation, and carcinogenic risk of polycyclic aromatic hydrocarbons in aquatic organisms from Lake Chaohu, China. <i>Science of the Total Environment</i> , 2020, 749, 141577.	3.9	21
529	Effects of Naturally Occurring Aquatic Organic Fractions on 241 Am Uptake by <i>Scenedesmus obliquus</i> (Chlorophyceae) and <i>Aeromonas hydrophila</i> (Pseudomonadaceae). <i>Applied and Environmental Microbiology</i> , 1977, 33, 89-96.	1.4	21
530	Organochlorine Contaminants in Double-Crested Cormorants from Green Bay, Wisconsin: II. Effects of an Extract Derived from Cormorant Eggs on the Chicken Embryo. <i>Archives of Environmental Contamination and Toxicology</i> , 1997, 32, 316-322.	2.1	20
531	Comparison of Risk Assessment Methodologies for Exposure of Mink to PCBs on the Kalamazoo River, Michigan. <i>Environmental Science & Technology</i> , 2004, 38, 6451-6459.	4.6	20
532	Distribution and sources of mercury in soils from former industrialized urban areas of Beijing, China. <i>Environmental Monitoring and Assessment</i> , 2009, 158, 507-517.	1.3	20
533	Dioxin-Like and Endocrine Disruptive Activity of Traffic-Contaminated Soil Samples. <i>Archives of Environmental Contamination and Toxicology</i> , 2009, 57, 639-650.	2.1	20
534	Effects of in ovo exposure of white leghorn chicken, common pheasant, and Japanese quail to 2,3,7,8-tetrachlorodibenzo-p-dioxin and two chlorinated dibenzofurans on CYP1A induction. <i>Environmental Toxicology and Chemistry</i> , 2010, 29, 1490-1502.	2.2	20
535	PCB concentrations in walleyes and their prey from the Saginaw River, Lake Huron: A comparison between 1990 and 2007. <i>Journal of Great Lakes Research</i> , 2010, 36, 267-276.	0.8	20
536	Status and fuzzy comprehensive assessment of metals and arsenic contamination in farmland soils along the Yanghe River, China. <i>Chemistry and Ecology</i> , 2011, 27, 415-426.	0.6	20
537	Characterization of a bystander effect induced by the endocrine-disrupting chemical 6-propyl-2-thiouracil in zebrafish embryos. <i>Aquatic Toxicology</i> , 2012, 118-119, 108-115.	1.9	20
538	Effects of multigenerational exposures of <i>D. magna</i> to environmentally relevant concentrations of pentachlorophenol. <i>Environmental Science and Pollution Research</i> , 2014, 21, 234-243.	2.7	20
539	Derivation of marine water quality criteria for metals based on a novel QICAR-SSD model. <i>Environmental Science and Pollution Research</i> , 2015, 22, 4297-4304.	2.7	20
540	Urinary bromophenol glucuronide and sulfate conjugates: Potential human exposure molecular markers for polybrominated diphenyl ethers. <i>Chemosphere</i> , 2015, 133, 6-12.	4.2	20

#	ARTICLE	IF	CITATIONS
541	Organobromine compound profiling in human adipose: Assessment of sources of bromophenol. <i>Environmental Pollution</i> , 2015, 204, 81-89.	3.7	20
542	Oil sands process-affected water impairs feeding by <i>Daphnia magna</i> . <i>Chemosphere</i> , 2017, 175, 465-472.	4.2	20
543	Assessment of potential biological activities and distributions of endocrine-disrupting chemicals in sediments of the west coast of South Korea. <i>Chemosphere</i> , 2017, 168, 441-449.	4.2	20
544	Maternal transfer of bioactive polychlorinated aromatic hydrocarbons in spawning chinook salmon (<i>Oncorhynchus tshawytscha</i>). <i>Marine Environmental Research</i> , 1989, 28, 231-234.	1.1	19
545	Polychlorinated biphenyls and 2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents in eggs of double-crested cormorants from a colony near Green Bay, Wisconsin, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 1995, 29, 327-333.	2.1	19
546	Effects of copper-contaminated sediments on <i>Hyalella azteca</i> , <i>Daphnia magna</i> , and <i>Ceriodaphnia dubia</i> : Survival, growth, and enzyme inhibition. <i>Archives of Environmental Contamination and Toxicology</i> , 1995, 29, 97-103.	2.1	19
547	In vitro assessment of potential mechanism-specific effects of polybrominated diphenyl ethers. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 2431-2433.	2.2	19
548	Quaternary benzo[c]phenanthridine alkaloids sanguinarine and chelerythrine do not affect transcriptional activity of aryl hydrocarbon receptor: Analyses in rat hepatoma cell line H4IIE.luc. <i>Food and Chemical Toxicology</i> , 2006, 44, 1466-1473.	1.8	19
549	RISK ASSESSMENT OF GREAT HORNED OWLS (<i>BUBO VIRGINIANUS</i>) EXPOSED TO POLYCHLORINATED BIPHENYLS AND DDT ALONG THE KALAMAZOO RIVER, MICHIGAN, USA. <i>Environmental Toxicology and Chemistry</i> , 2007, 26, 1386.	2.2	19
550	Effects of energy conservation in major energy-intensive industrial sectors on emissions of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in China. <i>Energy Policy</i> , 2010, 38, 2346-2356.	4.2	19
551	Endocrine disruption effects of 2,2,4,4,6-pentabromodiphenylether (BDE100) in reporter gene assays. <i>Journal of Environmental Monitoring</i> , 2011, 13, 850.	2.1	19
552	Detection, Identification, and Quantification of Hydroxylated Bis(2-ethylhexyl)-Tetrabromophthalate Isomers in House Dust. <i>Environmental Science & Technology</i> , 2015, 49, 2999-3006.	4.6	19
553	Sunlight Irradiation of Highly Brominated Polyphenyl Ethers Generates Polybenzofuran Products That Alter Dioxin-responsive mRNA Expression in Chicken Hepatocytes. <i>Environmental Science & Technology</i> , 2016, 50, 2318-2327.	4.6	19
554	Using dual isotopes and a Bayesian isotope mixing model to evaluate sources of nitrate of Tai Lake, China. <i>Environmental Science and Pollution Research</i> , 2018, 25, 32631-32639.	2.7	19
555	In vitro assessment of endocrine disrupting potential of organic fractions extracted from hydraulic fracturing flowback and produced water (HF-FPW). <i>Environment International</i> , 2018, 121, 824-831.	4.8	19
556	Molecular Initiating Events of Bisphenols on Androgen Receptor-Mediated Pathways Provide Guidelines for <i>in Silico</i> Screening and Design of Substitute Compounds. <i>Environmental Science and Technology Letters</i> , 2019, 6, 205-210.	3.9	19
557	Polychlorinated naphthalenes, biphenyls, dibenzo-p-dioxins, and dibenzofurans as well as polycyclic aromatic hydrocarbons and alkylphenols in sediment from the Detroit and Rouge Rivers, Michigan, USA. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 1878-89.	2.2	19
558	The effect of season and location on phosphoadenylate concentrations and adenylate energy charge in two species of freshwater clams. <i>Oecologia</i> , 1981, 49, 1-7.	0.9	18

#	ARTICLE	IF	CITATIONS
559	Sewage effluent biomonitoring. <i>Ecotoxicology and Environmental Safety</i> , 1985, 10, 22-39.	2.9	18
560	Influences on copper bioaccumulation, growth, and survival of the midge, <i>Chironomus tentans</i> , in metal-contaminated sediments. <i>Journal of Aquatic Ecosystem Health</i> , 1995, 4, 157-168.	0.4	18
561	Polychlorinated Dibenzo-p-Dioxins (PCDDs) and Dibenzofurans (PCDFs) in Muscle and Eggs of Salmonid Fishes from the Great Lakes. <i>Archives of Environmental Contamination and Toxicology</i> , 1999, 36, 432-446.	2.1	18
562	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 176-184.	2.7	18
563	Passerine Exposure to Primarily PCDFs and PCDDs in the River Floodplains Near Midland, Michigan, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 2010, 58, 1048-1064.	2.1	18
564	2,3,4,7,8-pentachlorodibenzofuran is a more potent cytochrome P4501A inducer than 2,3,7,8-tetrachlorodibenzo-p-dioxin in herring gull hepatocyte cultures. <i>Environmental Toxicology and Chemistry</i> , 2010, 29, 2088-2095.	2.2	18
565	Probabilistic ecological risk assessment for three chlorophenols in surface waters of China. <i>Journal of Environmental Sciences</i> , 2012, 24, 329-334.	3.2	18
566	Lead in drinking water and human blood in Riyadh City, Saudi Arabia. <i>Arabian Journal of Geosciences</i> , 2013, 6, 3103-3109.	0.6	18
567	Effect-based assessment of passive air samples from four countries in Eastern Europe. <i>Environmental Monitoring and Assessment</i> , 2014, 186, 3905-3916.	1.3	18
568	Instrumental and bioanalytical measures of dioxin-like compounds and activities in sediments of the Pohang Area, Korea. <i>Science of the Total Environment</i> , 2014, 470-471, 1517-1525.	3.9	18
569	Reduction of dioxin-like toxicity in effluents by additional wastewater treatment and related effects in fish. <i>Ecotoxicology and Environmental Safety</i> , 2016, 132, 47-58.	2.9	18
570	Activation of AhR-mediated toxicity pathway by emerging pollutants polychlorinated diphenyl sulfides. <i>Chemosphere</i> , 2016, 144, 1754-1762.	4.2	18
571	A high-throughput, computational system to predict if environmental contaminants can bind to human nuclear receptors. <i>Science of the Total Environment</i> , 2017, 576, 609-616.	3.9	18
572	Neonicotinoid insecticides in pollen, honey and adult bees in colonies of the European honey bee (<i>Apis mellifera</i>). <i>Overlock 10</i> , 2017, 1.1, 18.	1.1	18
573	Residues levels of pesticides in walnuts of Iran and associated health risks. <i>Human and Ecological Risk Assessment (HERA)</i> , 2021, 27, 191-204.	1.7	18
574	Organophosphate esters in agro-foods: Occurrence, sources and emerging challenges. <i>Science of the Total Environment</i> , 2022, 827, 154271.	3.9	18
575	Fate of anthracene in an artificial stream: A case study. <i>Ecotoxicology and Environmental Safety</i> , 1984, 8, 183-201.	2.9	17
576	Toxicity of Vertical Sediments in the Trenton Channel, Detroit River, Michigan, to <i>Chironomus Tentans</i> (Insecta: Chironomidae). <i>Journal of Great Lakes Research</i> , 1989, 15, 570-580.	0.8	17

#	ARTICLE	IF	CITATIONS
577	Immunoassay monitoring of polychlorinated biphenyls (PCBs) in the Great Lakes. <i>Environmental Science and Pollution Research</i> , 1994, 1, 69-74.	2.7	17
578	Sources and distribution of polychlorinated dibenzo-p-dioxins and dibenzofurans in sediments from Masan Bay, Korea. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 245-252.	2.2	17
579	EVALUATION OF THE METHOXYTRIAZINE HERBICIDE PROMETON USING A SHORT-TERM FATHEAD MINNOW REPRODUCTION TEST AND A SUITE OF IN VITRO BIOASSAYS. <i>Environmental Toxicology and Chemistry</i> , 2006, 25, 2143.	2.2	17
580	Endocrine-disrupting equivalents in industrial effluents discharged into Yangtze River. <i>Ecotoxicology</i> , 2009, 18, 685-692.	1.1	17
581	Perfluorinated Compounds in Aquatic Products from Bohai Bay, Tianjin, China. <i>Human and Ecological Risk Assessment (HERA)</i> , 2011, 17, 1279-1291.	1.7	17
582	In vitro effects of pollutants from particulate and volatile fractions of air samples day and night variability. <i>Environmental Science and Pollution Research</i> , 2013, 20, 6620-6627.	2.7	17
583	Occurrence and Potential Causes of Androgenic Activities in Source and Drinking Water in China. <i>Environmental Science & Technology</i> , 2013, 47, 130828135947000.	4.6	17
584	Perfluoroalkyl substances and organochlorine pesticides in sediments from Huaihe watershed in China. <i>Journal of Environmental Sciences</i> , 2014, 26, 2198-2206.	3.2	17
585	Dioxins and dioxin-like compounds in composts and digestates from European countries as determined by the in vitro bioassay and chemical analysis. <i>Chemosphere</i> , 2015, 122, 168-175.	4.2	17
586	Exposure of honeybees (<i>Apis mellifera</i>) in Saskatchewan, Canada to organophosphorus insecticides. <i>Apidologie</i> , 2015, 46, 667-678.	0.9	17
587	Identification and response to metals of metallothionein in two ancient fishes: White sturgeon (<i>Acipenser transmontanus</i>) and lake sturgeon (<i>Acipenser fulvescens</i>). <i>Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology</i> , 2015, 171, 41-48.	1.3	17
588	Impairment of reproduction of adult zebrafish (<i>Danio rerio</i>) by binary mixtures of environmentally relevant concentrations of triclocarban and inorganic mercury. <i>Ecotoxicology and Environmental Safety</i> , 2016, 134, 124-132.	2.9	17
589	Generalized concentration addition accurately predicts estrogenic potentials of mixtures and environmental samples containing partial agonists. <i>Toxicology in Vitro</i> , 2018, 46, 294-303.	1.1	17
590	Ractopamine and Other Growth-Promoting Compounds in Beef Cattle Operations: Fate and Transport in Feedlot Pens and Adjacent Environments. <i>Environmental Science & Technology</i> , 2021, 55, 1730-1739.	4.6	17
591	Antioxidant, Antimicrobial, and Insecticidal Properties of a Chemically Characterized Essential Oil from the Leaves of <i>Dittrichia viscosa</i> L.. <i>Molecules</i> , 2022, 27, 2282.	1.7	17
592	The effect of short-term exposure to pentachlorophenol and osmotic stress on the free amino acid pool of the freshwater amphipod <i>Gammarus pseudolimnaeus</i> Bousfield. <i>Archives of Environmental Contamination and Toxicology</i> , 1987, 16, 167-176.	2.1	16
593	Polychlorinated Dibenzo-p-Dioxins (PCDDs), Dibenzofurans (PCDFs), Biphenyls (PCBs), and Organochlorine Pesticides in Yellow-Blotched Map Turtle from the Pascagoula River Basin, Mississippi, USA. <i>Archives of Environmental Contamination and Toxicology</i> , 2000, 38, 362-370.	2.1	16
594	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 220-224.	2.7	16

#	ARTICLE	IF	CITATIONS
595	Predicted Distribution and Ecological Risk Assessment of a "Segregated" Hydrofluoroether in the Japanese Environment. <i>Environmental Science & Technology</i> , 2002, 36, 4761-4769.	4.6	16
596	Dioxin-like and non-dioxin like effects of polychlorinated biphenyls: Implications for risk assessment. <i>Lakes and Reservoirs: Research and Management</i> , 2002, 7, 139-181.	0.6	16
597	Cytotoxicity of HC Orange NO. 1 to L929 fibroblast cells. <i>Environmental Toxicology and Pharmacology</i> , 2008, 26, 309-314.	2.0	16
598	Toxicokinetics Of 2,3,7,8-TCDF and 2,3,4,7,8-PeCDF in Mink (<i>Mustela vison</i>) at Ecologically Relevant Exposures. <i>Toxicological Sciences</i> , 2008, 105, 33-43.	1.4	16
599	Mechanisms of toxicity of triphenyltin chloride (TPTC) determined by a live cell reporter array. <i>Environmental Science and Pollution Research</i> , 2013, 20, 803-811.	2.7	16
600	Development of aquatic life criteria in China: viewpoint on the challenge. <i>Environmental Science and Pollution Research</i> , 2014, 21, 61-66.	2.7	16
601	Dioxin-like activity in sediments from Tai Lake, China determined by use of the H4IIIE-luc bioassay and quantification of individual AhR agonists. <i>Environmental Science and Pollution Research</i> , 2014, 21, 1480-1488.	2.7	16
602	Toxicokinetics and toxicodynamics of chlorpyrifos is altered in embryos of Japanese medaka exposed to oil sands process-affected water: evidence for inhibition of P-glycoprotein. <i>Journal of Applied Toxicology</i> , 2017, 37, 591-601.	1.4	16
603	Extended Virtual Screening Strategies To Link Antiandrogenic Activities and Detected Organic Contaminants in Soils. <i>Environmental Science & Technology</i> , 2017, 51, 12528-12536.	4.6	16
604	Assessment of tools for protection of quality of water: Uncontrollable discharges of pollutants. <i>Ecotoxicology and Environmental Safety</i> , 2018, 161, 190-197.	2.9	16
605	Underlying mechanisms of reproductive toxicity caused by multigenerational exposure of 2, bromo-4, 6-dinitroaniline (BDNA) to Zebrafish (<i>Danio rerio</i>) at environmental relevant levels. <i>Aquatic Toxicology</i> , 2019, 216, 105285.	1.9	16
606	Adsorption, aggregation and sedimentation of titanium dioxide nanoparticles and nanotubes in the presence of different sources of humic acids. <i>Science of the Total Environment</i> , 2019, 692, 660-668.	3.9	16
607	Effects of the husky oil spill on gut microbiota of native fishes in the North Saskatchewan River, Canada. <i>Aquatic Toxicology</i> , 2020, 229, 105658.	1.9	16
608	Fates of cadmium introduced into channels microcosm. <i>Environment International</i> , 1981, 5, 159-175.	4.8	15
609	A thermal effluent as a sporadic cornucopia: effects on fish and zooplankton. <i>Environmental Biology of Fishes</i> , 1984, 11, 191-203.	0.4	15
610	Lysosomal enzyme release in the bluegill sunfish (<i>Lepomis macrochirus Rafinesque</i>) exposed to cadmium. <i>Archives of Environmental Contamination and Toxicology</i> , 1985, 14, 631-640.	2.1	15
611	Planar chlorinated hydrocarbons (PCHs) in colonial fish-eating waterbird eggs from the Great Lakes. <i>Marine Environmental Research</i> , 1989, 28, 505-508.	1.1	15
612	Spatial and Temporal Trends of Mercury Loadings to Michigan Inland Lakes. <i>Environmental Science & Technology</i> , 2007, 41, 5634-5640.	4.6	15

#	ARTICLE	IF	CITATIONS
613	In situ effects of urban river pollution on the mudsnail <i>Potamopyrgus antipodarum</i> as part of an integrated assessment. <i>Aquatic Toxicology</i> , 2014, 150, 83-92.	1.9	15
614	Non-parametric kernel density estimation of species sensitivity distributions in developing water quality criteria of metals. <i>Environmental Science and Pollution Research</i> , 2015, 22, 13980-13989.	2.7	15
615	Classification and toxicity mechanisms of novel flame retardants (NFRs) based on whole genome expression profiling. <i>Chemosphere</i> , 2016, 144, 2150-2157.	4.2	15
616	Phthalate Esters on Hands of Office Workers: Estimating the Influence of Touching Surfaces. <i>Environmental Science and Technology Letters</i> , 2017, 4, 1-5.	3.9	15
617	Transcriptomic responses of <i>Artemia salina</i> exposed to an environmentally relevant dose of <i>Alexandrium minutum</i> cells or Gonyautoxin2/3. <i>Chemosphere</i> , 2020, 238, 124661.	4.2	15
618	Effects of tris (2-chloroethyl) phosphate (TCEP) on growth, reproduction and gene transcription in the protozoan <i>Tetrahymena thermophila</i> . <i>Aquatic Toxicology</i> , 2020, 222, 105477.	1.9	15
619	Difference in performance and mechanism for methylene blue when TiO ₂ nanoparticles are converted to nanotubes. <i>Journal of Cleaner Production</i> , 2021, 297, 126498.	4.6	15
620	Novel polar AhR-active chemicals detected in sediments of an industrial area using effect-directed analysis based on in vitro bioassays with full-scan high resolution mass spectrometric screening. <i>Science of the Total Environment</i> , 2021, 779, 146566.	3.9	15
621	Arsenic concentrations in water and fish from Chautauqua Lake, New York. <i>Environmental Biology of Fishes</i> , 1978, 3, 361-367.	0.4	14
622	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 115-115.	2.7	14
623	Assessing environmental change through chemical-sediment chronologies from inland lakes. <i>Lakes and Reservoirs: Research and Management</i> , 2002, 7, 217-230.	0.6	14
624	Chronic toxicity of contaminated sediments on reproduction and histopathology of the crustacean <i>Gammarus fossarum</i> and relationship with the chemical contamination and in vitro effects. <i>Journal of Soils and Sediments</i> , 2010, 10, 423-433.	1.5	14
625	Organochlorine pesticides (HCHs and DDTs) in soils along the north coastal areas of the Bohai Sea, China. <i>Chemistry and Ecology</i> , 2010, 26, 339-352.	0.6	14
626	Effects of treatments with Apivar [®] and Thymovar [®] on <i>V. destructor</i> populations, virus infections and indoor winter survival of Canadian honey bee (<i>Apis mellifera</i>)	Tj ETQq0 0 0 BT /Overlock 10 Tf	14
627	In vitro dioxin-like potencies of HO- and MeO-PBDEs and inter-species sensitivity variation in birds. <i>Ecotoxicology and Environmental Safety</i> , 2016, 126, 202-210.	2.9	14
628	Predicting criteria continuous concentrations of metals or metalloids for protecting marine life by use of quantitative ion characteristic activity relationships species sensitivity distributions (QICAR-SSD). <i>Marine Pollution Bulletin</i> , 2017, 124, 639-644.	2.3	14
629	Spatial and temporal ecological risk assessment of unionized ammonia nitrogen in Tai Lake, China (2004-2015). <i>Ecotoxicology and Environmental Safety</i> , 2017, 140, 249-255.	2.9	14
630	Temporal and spatial differences in deposition of organic matter and black carbon in Lake Michigan sediments over the period 1850-2010. <i>Journal of Great Lakes Research</i> , 2018, 44, 705-715.	0.8	14

#	ARTICLE	IF	CITATIONS
631	Integrated in silico and in vivo approaches to investigate effects of BDEâ€™99 mediated by the nuclear receptors on developing zebrafish. <i>Environmental Toxicology and Chemistry</i> , 2018, 37, 780-787.	2.2	14
632	Identification of potential toxicants in sediments from an industrialized area in Pohang, South Korea: Application of a cell viability assay of microalgae using flow cytometry. <i>Journal of Hazardous Materials</i> , 2021, 405, 124230.	6.5	14
633	Reproductive toxicity and metabolic perturbations in male rats exposed to boron. <i>Science of the Total Environment</i> , 2021, 785, 147370.	3.9	14
634	Organophosphate esters cause thyroid dysfunction via multiple signaling pathways in zebrafish brain. <i>Environmental Science and Ecotechnology</i> , 2022, 12, 100198.	6.7	14
635	Metal binding capacity of northern European surface waters for Cd, Cu, and Pb. <i>Organic Geochemistry</i> , 1980, 2, 57-67.	0.9	13
636	Classification of Chemicals Based on Concentration-Dependent Toxicological Data Using ToxClust. <i>Environmental Science & Technology</i> , 2009, 43, 3926-3932.	4.6	13
637	Daily intake of selenium and concentrations in blood of residents of Riyadh City, Saudi Arabia. <i>Environmental Geochemistry and Health</i> , 2012, 34, 417-431.	1.8	13
638	Effects of pig manure containing copper and zinc on microbial community assessed via phospholipids in soils. <i>Environmental Monitoring and Assessment</i> , 2014, 186, 5297-5306.	1.3	13
639	A national pilot scheme for monitoring and assessment of ecological integrity of surface waters in China. <i>Environmental Development</i> , 2014, 10, 104-107.	1.8	13
640	Improvement on species sensitivity distribution methods for deriving site-specific water quality criteria. <i>Environmental Science and Pollution Research</i> , 2015, 22, 5271-5282.	2.7	13
641	Effects of fulvic acid on aggregation, sedimentation, and adsorption of Fe ₃ O ₄ magnetic nanoparticles. <i>Environmental Science and Pollution Research</i> , 2019, 26, 21463-21474.	2.7	13
642	Halogenated flame retardants in sediments from the Upper Laurentian Great Lakes: Implications to long-range transport and evidence of long-term transformation. <i>Journal of Hazardous Materials</i> , 2020, 384, 121346.	6.5	13
643	Sediment pore water toxicity identification in the lower fox river and Green Bay, Wisconsin, using the microtox assay. <i>Ecotoxicology and Environmental Safety</i> , 1992, 23, 343-354.	2.9	12
644	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the north american great lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 51-51.	2.7	12
645	Organochlorine Insecticides in Mudflats of Hong Kong, China. <i>Archives of Environmental Contamination and Toxicology</i> , 2006, 50, 153-165.	2.1	12
646	Reproductive success of passerines exposed to polychlorinated biphenyls through the terrestrial food web of the Kalamazoo River. <i>Ecotoxicology and Environmental Safety</i> , 2007, 66, 107-118.	2.9	12
647	Ethoxyresorufin O-deethylase induction by TCDD, PeCDF and TCDF in ring-necked pheasant and Japanese quail hepatocytes: Time-dependent effects on concentrationâ€™response curves. <i>Toxicology in Vitro</i> , 2010, 24, 1301-1305.	1.1	12
648	Evaluation and Spatial Diffusion of Health Risk of Persistent Organic Pollutants (POPs) in Soils Surrounding Chemical Industrial Parks in China. <i>Human and Ecological Risk Assessment (HERA)</i> , 2010, 16, 989-1006.	1.7	12

#	ARTICLE	IF	CITATIONS
649	Effects on tree swallows exposed to dioxin-like compounds associated with the Tittabawassee River and floodplain near Midland, Michigan, USA. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 1354-1365.	2.2	12
650	Developmental and posthatch effects of in ovo exposure to 2,3,7,8-TCDD, 2,3,4,7,8-PECDF, and 2,3,7,8-TCDF in Japanese quail (<i>Coturnix japonica</i>), common pheasant (<i>Phasianus colchicus</i>), and white leghorn chicken (<i>Gallus gallus domesticus</i>) embryos. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 1659-1668.	2.2	12
651	Effects of a non-steroidal aromatase inhibitor on ovarian function in cattle. <i>Reproduction, Fertility and Development</i> , 2012, 24, 631.	0.1	12
652	Spatio-temporal effects of fertilization in Anhui Province, China. <i>Environment, Development and Sustainability</i> , 2015, 17, 1197-1207.	2.7	12
653	Linking genomic responses of gonads with reproductive impairment in marine medaka (<i>Oryzias latipes</i>) exposed to 10 ng BT (Overlock (DIM)). <i>Aquatic Toxicology</i> , 2017, 183, 135-143.	1.9	12
654	Using solid ¹³ C NMR coupled with solution ³¹ P NMR spectroscopy to investigate molecular species and lability of organic carbon and phosphorus from aquatic plants in Tai Lake, China. <i>Environmental Science and Pollution Research</i> , 2017, 24, 1880-1889.	2.7	12
655	Aryl hydrocarbon receptor-mediated potencies in field-deployed plastics vary by type of polymer. <i>Environmental Science and Pollution Research</i> , 2019, 26, 9079-9088.	2.7	12
656	Exposure to short-chain chlorinated paraffins inhibited PPAR α -mediated fatty acid oxidation and stimulated aerobic glycolysis in vitro in human cells. <i>Science of the Total Environment</i> , 2021, 772, 144957.	3.9	12
657	Essential Oils from Leaves of <i>Juniperus thurifera</i> L., Exhibiting Antioxidant, Antifungal and Antibacterial Activities against Antibiotic-Resistant Microbes. <i>Horticulturae</i> , 2022, 8, 321.	1.2	12
658	Comparison of primary and secondary sludge carbon sources derived from hydrolysis or acidogenesis for nitrate reduction and denitrification kinetics: Organics utilization and microbial community shift. <i>Environmental Research</i> , 2022, 212, 113403.	3.7	12
659	Assessment of sediment contamination at Great Lakes Areas of Concern: the ARCS Program Toxicity-Chemistry Work Group strategy. <i>Journal of Aquatic Ecosystem Health</i> , 1992, 1, 193-200.	0.4	11
660	Polychlorinated biphenyls and 2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents in eggs of red-breasted mergansers near Green Bay, Wisconsin, USA, in 1977-78 and 1990. <i>Archives of Environmental Contamination and Toxicology</i> , 1995, 29, 52-60.	2.1	11
661	Assessment of potential exposure to agent orange and its associated TCDD. <i>Environmental Science and Pollution Research</i> , 2004, 11, 347-348.	2.7	11
662	Estrogenic and Dioxin-like Activities and Cytotoxicity of Sediments and Biota from Hong Kong Mudflats. <i>Archives of Environmental Contamination and Toxicology</i> , 2005, 48, 575-586.	2.1	11
663	Spatial variability and temporal trends of HCH and DDT in soils around Beijing Guanting Reservoir, China. <i>Environmental Geochemistry and Health</i> , 2010, 32, 441-449.	1.8	11
664	Effect-Directed Analysis of Ah-Receptor Mediated Toxicants, Mutagens, and Endocrine Disruptors in Sediments and Biota. <i>Handbook of Environmental Chemistry</i> , 2011, , 285-313.	0.2	11
665	Effects of Exposure to 17 β -Ethinylestradiol during Sexual Differentiation on the Transcriptome of the African Clawed Frog (<i>Xenopus laevis</i>). <i>Environmental Science & Technology</i> , 2013, 47, 4822-4828.	4.6	11
666	Weighted species sensitivity distribution method to derive site-specific quality criteria for copper in Tai Lake, China. <i>Environmental Science and Pollution Research</i> , 2014, 21, 12968-12978.	2.7	11

#	ARTICLE	IF	CITATIONS
667	Longer-term and short-term variability in pollution of fluvial sediments by dioxin-like and endocrine disruptive compounds. <i>Environmental Science and Pollution Research</i> , 2014, 21, 5007-5022.	2.7	11
668	Distribution and bioaccumulation of lead in the coastal watersheds of the Northern Bohai and Yellow Seas in China. <i>Environmental Geochemistry and Health</i> , 2015, 37, 491-506.	1.8	11
669	Distribution Characteristics and Risk Assessments of PAHs in Fish from Lake Taihu, China. <i>Human and Ecological Risk Assessment (HERA)</i> , 2015, 21, 1753-1765.	1.7	11
670	Bioanalytical and instrumental screening of the uptake of sediment-borne, dioxin-like compounds in roach (<i>Rutilus rutilus</i>). <i>Environmental Science and Pollution Research</i> , 2016, 23, 12060-12074.	2.7	11
671	Environmental DNA of preservative ethanol performed better than water samples in detecting macroinvertebrate diversity using metabarcoding. <i>Diversity and Distributions</i> , 2021, 27, 1989-2002.	1.9	11
672	USE OF LINEAR ORTHOGONAL CONTRASTS IN ANALYSIS OF ENVIRONMENTAL DATA. <i>Environmental Toxicology and Chemistry</i> , 1990, 9, 815.	2.2	11
673	Utility of the QSAR modeling system for predicting the toxicity of substances on the European inventory of existing commercial chemicals. <i>Toxicological and Environmental Chemistry</i> , 1990, 28, 167-188.	0.6	10
674	A risk-based protocol to develop acceptable concentrations of bioaccumulative organic chemicals in sediments for the protection of piscivorous wildlife. <i>Toxicological and Environmental Chemistry</i> , 1996, 54, 243-259.	0.6	10
675	Instrumental and Bioanalytical Measures of Endocrine Disruptors in Water. <i>ACS Symposium Series</i> , 1999, , 73-95.	0.5	10
676	Concentrations and profiles of polychlorinated biphenyls, -dibenzo-p-dioxins and -dibenzofurans in livers of mink from South Carolina and Louisiana, U.S.A. <i>Environmental Monitoring and Assessment</i> , 2003, 83, 17-33.	1.3	10
677	Oxidative Stress in Laboratory-Incubated Double-Crested Cormorant Eggs Collected from the Great Lakes. <i>Archives of Environmental Contamination and Toxicology</i> , 2003, 45, 533-546.	2.1	10
678	Comparison of gene expression methods to identify genes responsive to perfluorooctane sulfonic acid. <i>Environmental Toxicology and Pharmacology</i> , 2005, 19, 153-160.	2.0	10
679	Exposure and Multiple Lines of Evidence Assessment of Risk for PCBs Found in the Diets of Passerine Birds at the Kalamazoo River Superfund Site, Michigan. <i>Human and Ecological Risk Assessment (HERA)</i> , 2006, 12, 924-946.	1.7	10
680	In situ hybridization to detect spatial gene expression in medaka. <i>Ecotoxicology and Environmental Safety</i> , 2009, 72, 1257-1264.	2.9	10
681	Genotoxicity of crude extracts of cyanobacteria from Taihu Lake on carp (<i>Cyprinus carpio</i>). <i>Ecotoxicology</i> , 2011, 20, 1010-1017.	1.1	10
682	Multiple Lines of Evidence Risk Assessment of Terrestrial Passerines Exposed to PCDFs and PCDDs in the Tittabawassee River Floodplain, Midland, Michigan, USA. <i>Human and Ecological Risk Assessment (HERA)</i> , 2011, 17, 159-186.	1.7	10
683	Environmental and health challenges of the global growth of electronic waste. <i>Environmental Science and Pollution Research</i> , 2012, 19, 2460-2462.	2.7	10
684	Effects of Columbia River water on early life-stages of white sturgeon (<i>Acipenser transmontanus</i>). <i>Ecotoxicology and Environmental Safety</i> , 2014, 101, 23-30.	2.9	10

#	ARTICLE	IF	CITATIONS
685	Characterization of endocrine disruption potentials of coastal sediments of Taean, Korea employing H295R and MVLN assays—Reconnaissance at 5 years after Hebei Spirit oil spill. <i>Marine Pollution Bulletin</i> , 2018, 127, 264-272.	2.3	10
686	Characterization and sources of dissolved and particulate phosphorus in 10 freshwater lakes with different trophic statuses in China by solution ³¹ P nuclear magnetic resonance spectroscopy. <i>Ecological Research</i> , 2019, 34, 106-118.	0.7	10
687	Light, but Not Nutrients, Drives Seasonal Congruence of Taxonomic and Functional Diversity of Phytoplankton in a Eutrophic Highland Lake in China. <i>Frontiers in Plant Science</i> , 2020, 11, 179.	1.7	10
688	Long-term trends of persistent toxic substances and potential toxicities in sediments along the west coast of South Korea. <i>Marine Pollution Bulletin</i> , 2020, 151, 110821.	2.3	10
689	Molecular mechanisms of zooplanktonic toxicity in the okadaic acid-producing dinoflagellate <i>Prorocentrum lima</i> . <i>Environmental Pollution</i> , 2021, 279, 116942.	3.7	10
690	Effect-directed identification of novel aryl hydrocarbon receptor-active aromatic compounds in coastal sediments collected from a highly industrialized area. <i>Science of the Total Environment</i> , 2022, 803, 149969.	3.9	10
691	Identification of Thyroid Receptor Ant/Agonists in Water Sources Using Mass Balance Analysis and Monte Carlo Simulation. <i>PLoS ONE</i> , 2013, 8, e73883.	1.1	10
692	Effect of 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) on the Epidermal Growth Factor Receptor in Hepatic Plasma Membranes of Rainbow Trout. <i>Toxicology and Applied Pharmacology</i> , 1993, 118, 119-130.	1.3	9
693	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 116-121.	2.7	9
694	Hormesis — does it have relevance at the population, community or ecosystem levels of organization?. <i>Human and Experimental Toxicology</i> , 2001, 20, 517-520.	1.1	9
695	Distribution of PCDDs and PCDFs in Soils Collected from the Denver Front Range - Principal Components Analysis of Diffuse Dioxin Sources (10 pp). <i>Environmental Science and Pollution Research</i> , 2005, 12, 189-198.	2.7	9
696	Changes of AhR-mediated activity of humic substances after irradiation. <i>Environment International</i> , 2007, 33, 812-816.	4.8	9
697	Expression profile of oestrogen receptors and oestrogen-related receptors is organ specific and sex dependent: the Japanese medaka <i>Oryzias latipes</i> model. <i>Journal of Fish Biology</i> , 2013, 83, 295-310.	0.7	9
698	Comparison of waterborne and in ovo nanoinjection exposures to assess effects of PFOS on zebrafish embryos. <i>Environmental Science and Pollution Research</i> , 2015, 22, 2303-2310.	2.7	9
699	Microbial Biomass and Community Composition Involved in Cycling of Organic Phosphorus in Sediments of Lake Dianchi, Southwest China. <i>Geomicrobiology Journal</i> , 2017, 34, 249-260.	1.0	9
700	Determination of water environment standards based on water quality criteria in China: Limitations and feasibilities. <i>Journal of Environmental Sciences</i> , 2017, 57, 127-136.	3.2	9
701	Hydroxylated 2-Ethylhexyl tetrabromobenzoate isomers in house dust and their agonistic potencies with several nuclear receptors. <i>Environmental Pollution</i> , 2017, 227, 578-586.	3.7	9
702	Biological toxicity estimates show involvement of a wider range of toxic compounds in sediments from Durban, South Africa than indicated from instrumental analyses. <i>Marine Pollution Bulletin</i> , 2019, 138, 49-57.	2.3	9

#	ARTICLE	IF	CITATIONS
703	Exposure of zebrafish to environmentally relevant concentrations of mercury during early life stages impairs subsequent reproduction in adults but can be recovered in offspring. <i>Aquatic Toxicology</i> , 2020, 229, 105655.	1.9	9
704	Bioaccumulation profiles of polychlorinated biphenyl congeners and organochlorine pesticides in Ganges river dolphins. , 1999, 18, 1511.		9
705	EFFECTS OF NONYLPHENOL ETHOXYLATE EXPOSURE ON REPRODUCTIVE OUTPUT AND BIOINDICATORS OF ENVIRONMENTAL ESTROGEN EXPOSURE IN FATHEAD MINNOWS, PIMEPHALES PROMELAS. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 510.	2.2	9
706	Frequency Distributions of the Concentrations of Essential and Nonessential Elements in Largemouth Bass, <i>Micropterus Salmoides</i> . <i>Ecology</i> , 1981, 62, 456-468.	1.5	8
707	Changes in cytochrome P4501A activity during development in common tern chicks fed polychlorinated biphenyls, as measured by the caffeine breath test. <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 712-718.	2.2	8
708	2,3,7,8-tetrachlorodibenzo-p-dioxin equivalents in tissue samples from three species in the Denver, Colorado, USA, metropolitan area. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 2433-2442.	2.2	8
709	Sediment TCDD-EQs and EROD and MROD Activities in Ranid Frogs from Agricultural and Nonagricultural Sites in Michigan (USA). <i>Archives of Environmental Contamination and Toxicology</i> , 2006, 51, 467-477.	2.1	8
710	Sequencing and characterization of mixed function monooxygenase genes CYP1A1 and CYP1A2 of Mink (<i>Mustela vison</i>) to facilitate study of dioxin-like compounds. <i>Toxicology and Applied Pharmacology</i> , 2009, 234, 306-313.	1.3	8
711	Standard purity and response factors of perfluorinated compounds. <i>Toxicological and Environmental Chemistry</i> , 2010, 92, 1219-1232.	0.6	8
712	Dietary exposure of three passerine species to PCDD/DFs from the Chippewa, Tittabawassee, and Saginaw River floodplains, Midland, Michigan, USA. <i>Environmental Monitoring and Assessment</i> , 2011, 172, 91-112.	1.3	8
713	A novel chemical additive for in-situ recovery of heavy oil using waterflooding process. <i>Journal of Petroleum Science and Engineering</i> , 2015, 135, 484-497.	2.1	8
714	Comparison of the Effects of Extraction Techniques on Mass Spectrometry Profiles of Dissolved Organic Compounds in Oil Sand Process-Affected Water. <i>Energy & Fuels</i> , 2019, 33, 7001-7008.	2.5	8
715	In vitro-in vivo and cross-life stage extrapolation of uptake and biotransformation of benzo[a]pyrene in the fathead minnow (<i>Pimephales promelas</i>). <i>Aquatic Toxicology</i> , 2020, 228, 105616.	1.9	8
716	Mechanisms of pH-Dependent Uptake of Ionizable Organic Chemicals by Fish from Oil Sands Process-Affected Water (OSPW). <i>Environmental Science & Technology</i> , 2020, 54, 9547-9555.	4.6	8
717	Ecotoxicological risk assessment of metal cocktails based on maximum cumulative ratio during multi-generational exposures. <i>Water Research</i> , 2021, 200, 117274.	5.3	8
718	Derivation and application of relative potency estimates based on in vitro bioassay results. , 2000, 19, 2835.		8
719	Effects of novel brominated flame retardants on steroidogenesis in primary porcine testicular cells. <i>Toxicology Letters</i> , 2014, 224, 141-6.	0.4	8
720	Sewage effluent biomonitoring. <i>Ecotoxicology and Environmental Safety</i> , 1985, 10, 40-52.	2.9	7

#	ARTICLE	IF	CITATIONS
721	Retinoids in eggs and embryos of birds fed fish from the Great Lakes. <i>Environmental Toxicology and Pharmacology</i> , 1997, 3, 277-288.	2.0	7
722	SCRAM: A scoring and ranking system for persistent, bioaccumulative, and toxic substances for the North American Great Lakes. <i>Environmental Science and Pollution Research</i> , 2000, 7, 219-219.	2.7	7
723	Laboratory analyses of the potential toxicity of sediment-associated polydimethylsiloxane to benthic macroinvertebrates. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 2611-2616.	2.2	7
724	Distribution of Copper, Cadmium, and Lead in Soils from Former Industrialized Urban Areas of Beijing, China. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2009, 82, 378-383.	1.3	7
725	Biochemical responses and DNA damage in red sea bream from coastal Fujian Province, China. <i>Ecotoxicology and Environmental Safety</i> , 2011, 74, 1526-1535.	2.9	7
726	Experience in South Africa of combining bioanalysis and instrumental analysis of PCDD/Fs. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 46, 189-197.	5.8	7
727	PHOTO-ENHANCED TOXICITY: SERENDIPITY OF A PREPARED MIND AND FLEXIBLE PROGRAM MANAGEMENT. <i>Environmental Toxicology and Chemistry</i> , 2013, 32, 969-971.	2.2	7
728	Species-specific relative AHR1 binding affinities of 2,3,4,7,8-pentachlorodibenzofuran explain avian species differences in its relative potency. <i>Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology</i> , 2014, 161, 21-25.	1.3	7
729	Responses of the zebrafish hypothalamic-pituitary-gonadal-liver axis PCR array to prochloraz are dependent on timing of sampling. <i>Aquatic Toxicology</i> , 2016, 175, 154-159.	1.9	7
730	Historical record of effects of human activities on absolute and relative concentrations of Polycyclic aromatic hydrocarbons (PAHs) in Lake Chao, China. <i>Journal of Environmental Sciences</i> , 2016, 46, 1-4.	3.2	7
731	Human dietary intake and hazard characterization for residues of neonicotinoid and organophosphorus pesticides in Egyptian honey. <i>Toxicological and Environmental Chemistry</i> , 2017, 99, 1397-1408.	0.6	7
732	In vitro tools for the toxicological evaluation of sediments and dredged materials: intra- and inter-laboratory comparisons of chemical and bioanalytical methods. <i>Environmental Science and Pollution Research</i> , 2018, 25, 4037-4050.	2.7	7
733	Down-Regulation of <i>hspb9</i> and <i>hspb11</i> Contributes to Wavy Notochord in Zebrafish Embryos Following Exposure to Polychlorinated Diphenylsulfides. <i>Environmental Science & Technology</i> , 2018, 52, 12829-12840.	4.6	7
734	Sublethal effects of chronic exposure to chlorpyrifos or imidacloprid insecticides or their binary mixtures on <i>Culex pipiens</i> mosquitoes. <i>Physiological Entomology</i> , 2019, 44, 123-132.	0.6	7
735	Effects of chemical fractions from an oil sands end-pit lake on reproduction of fathead minnows. <i>Chemosphere</i> , 2020, 249, 126073.	4.2	7
736	miR-155 influences cell-mediated immunity in Balb/c mice treated with aflatoxin M ₁ . <i>Drug and Chemical Toxicology</i> , 2021, 44, 39-46.	1.2	7
737	BIOCHEMICAL AND DEVELOPMENTAL EFFECTS OF DIETARY EXPOSURE TO POLYCHLORINATED BIPHENYLS 126 AND 153 IN COMMON TERN CHICKS (<i>STERNA HIRUNDO</i>). <i>Environmental Toxicology and Chemistry</i> , 2000, 19, 719.	2.2	7
738	FLOW CYTOMETRIC DETERMINATION OF THE PHOTOINDUCED TOXICITY OF ANTHRACENE TO THE GREEN ALGA <i>SELENASTRUM CAPRICORNUTUM</i> . <i>Environmental Toxicology and Chemistry</i> , 1994, 13, 831.	2.2	7

#	ARTICLE	IF	CITATIONS
739	A novel passive sampling and sequential extraction approach to investigate desorption kinetics of emerging organic contaminants at the sediment-water interface. <i>Water Research</i> , 2022, 217, 118455.	5.3	7
740	Development of a caffeine breath test to measure cytochrome P450-1A activity in birds. <i>Environmental Toxicology and Pharmacology</i> , 1996, 1, 51-61.	2.0	6
741	Relationship between polychlorinated biphenyl 126 treatment and cytochrome p4501a activity in chickens, as measured by in vivo caffeine and ex vivo ethoxyresorufin metabolism. <i>Environmental Toxicology and Chemistry</i> , 1999, 18, 2013-2022.	2.2	6
742	Hepatic P450 Enzyme Activity, Tissue Morphology and Histology of Mink (<i>Mustela vison</i>) Exposed to Polychlorinated Dibenzofurans. <i>Archives of Environmental Contamination and Toxicology</i> , 2009, 57, 416-425.	2.1	6
743	Ecological risk assessment of great horned owls (<i>Bubo virginianus</i>) exposed to PCDD/DF in the Tittabawassee River floodplain in Midland, Michigan, USA. <i>Environmental Toxicology and Chemistry</i> , 2010, 29, 2341-2349.	2.2	6
744	Dietary and tissue based exposure of belted kingfisher to PCDFs and PCDDs in the Tittabawassee River floodplain, Midland, MI, USA. <i>Environmental Toxicology and Chemistry</i> , 2012, 31, 1158-1168.	2.2	6
745	AhR-mediated activities and compounds in sediments of Meiliang Bay, Taihu Lake, China determined by in vitro bioassay and instrumental analysis. <i>RSC Advances</i> , 2015, 5, 55746-55755.	1.7	6
746	Identification of polycyclic aromatic hydrocarbons in soils in Taizhou, East China. <i>Environmental Geochemistry and Health</i> , 2015, 37, 429-439.	1.8	6
747	Optimization of QuEChERS extraction of steroid hormones from infant formulae for mass spectrometric analysis. <i>Toxicological and Environmental Chemistry</i> , 2021, 103, 1-17.	0.6	6
748	Transmission of SARS-CoV-2 virus and ambient temperature: a critical review. <i>Environmental Science and Pollution Research</i> , 2021, 28, 37051-37059.	2.7	6
749	Life Cycle Exposure to Environmentally Relevant Concentrations of Diphenyl Phosphate (DPhP) Inhibits Growth and Energy Metabolism of Zebrafish in a Sex-Specific Manner. <i>Environmental Science & Technology</i> , 2021, 55, 13122-13131.	4.6	6
750	Response to Comment on "Gonadal Development of Larval Male <i>Xenopus laevis</i> Exposed to Atrazine in Outdoor Microcosms". <i>Environmental Science & Technology</i> , 2005, 39, 7759-7760.	4.6	5
751	Polycyclic aromatic hydrocarbons in soils of an industrial area of China: multivariate analyses and geostatistics. <i>Chemistry and Ecology</i> , 2010, 26, 35-48.	0.6	5
752	Modulation of estrogen synthesis through activation of protein kinase A in H295R cells by extracts of estuary sediments. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 2793-2801.	2.2	5
753	Effects of dietary exposure of mink (<i>Mustela vison</i>) to 2,3,7,8-tetrachlorodibenzo-p-dioxin, 2,3,4,7,8-pentachlorodibenzofuran, and 2,3,7,8-tetrachlorodibenzofuran on reproduction and offspring viability and growth. <i>Environmental Toxicology and Chemistry</i> , 2012, 31, 360-369.	2.2	5
754	Vertical distributions of bound saturated fatty acids and compound-specific stable carbon isotope compositions in sediments of two lakes in China: implication for the influence of eutrophication. <i>Environmental Science and Pollution Research</i> , 2014, 21, 13138-13147.	2.7	5
755	The case for establishing a board of review for resolving environmental issues: The science court in Canada. <i>Integrated Environmental Assessment and Management</i> , 2016, 12, 572-579.	1.6	5
756	Integrated assessment of west coast of South Korea by use of benthic bacterial community structure as determined by eDNA, concentrations of contaminants, and in vitro bioassays. <i>Environment International</i> , 2020, 137, 105569.	4.8	5

#	ARTICLE	IF	CITATIONS
757	Toxicokinetic Models for Bioconcentration of Organic Contaminants in Two Life Stages of White Sturgeon (<i>Acipenser transmontanus</i>). <i>Environmental Science & Technology</i> , 2021, 55, 11590-11600.	4.6	5
758	Identification of novel polar aryl hydrocarbon receptor agonists accumulated in liver of black-tailed gulls in Korea using advanced effect-directed analysis. <i>Journal of Hazardous Materials</i> , 2022, 429, 128305.	6.5	5
759	Identification and quantitation of nonylphenol ethoxylates and nonylphenol in fish tissues from Michigan. <i>Environmental Science & Technology</i> , 2001, 35, 10-3.	4.6	5
760	Analysis of trace organic contaminants in sediment, pore water, and water samples from Onsan Bay, Korea: instrumental analysis and in vitro gene expression assay. <i>Environmental Toxicology and Chemistry</i> , 2002, 21, 1796-803.	2.2	5
761	Effects of in situ experimental selenium exposure on finescale dace (<i>Phoxinus neogaeus</i>) gut microbiome. <i>Environmental Research</i> , 2022, 212, 113151.	3.7	5
762	Errors in Determining Elemental Concentrations and the Structure of Interelement Correlation Matrices. <i>Ecology</i> , 1981, 62, 483-485.	1.5	4
763	Characterization of epidermal growth factor binding to hepatic plasma membranes of rainbow trout (<i>Oncorhynchus mykiss</i>). <i>General and Comparative Endocrinology</i> , 1991, 83, 345-353.	0.8	4
764	RAPID COMMUNICATION: BACKGROUND CONCENTRATIONS OF DIOXINS, FURANS, AND PCBs IN SPRAGUE-DAWLEY RATS AND JUVENILE SWINE. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2004, 67, 845-850.	1.1	4
765	Site-Specific Assessments of Environmental Risk and Natural Resource Damage Based on Great Horned Owls. <i>Human and Ecological Risk Assessment (HERA)</i> , 2007, 13, 966-985.	1.7	4
766	Tissue-based risk assessment of great blue heron (<i>Ardea herodias</i>) exposed to PCDD/DF in the Tittabawassee River floodplain, Michigan, USA. <i>Environmental Toxicology and Chemistry</i> , 2010, 29, 2544-2558.	2.2	4
767	Inferring sources for mercury to inland lakes using sediment chronologies of polycyclic aromatic hydrocarbons. <i>Environmental Sciences: Processes and Impacts</i> , 2014, 16, 2108-2116.	1.7	4
768	Human health risk assessment of soil dioxin/furans contamination and dioxin-like activity determined by ethoxyresorufin-O-deethylase bioassay. <i>Environmental Science and Pollution Research</i> , 2015, 22, 5218-5227.	2.7	4
769	Water-effect ratio of copper and its application on setting site-specific water quality criteria for protecting marine ecosystems of Hong Kong. <i>Environmental Science and Pollution Research</i> , 2018, 25, 3170-3182.	2.7	4
770	Tissue-based assessment of hazard posed by mercury and selenium to wild fishes in two shallow Chinese lakes. <i>Environmental Science and Pollution Research</i> , 2019, 26, 15989-15999.	2.7	4
771	Concentrations of Metals in Fishes from the Athabasca and Slave Rivers of Northern Canada. <i>Environmental Toxicology and Chemistry</i> , 2020, 39, 2180-2195.	2.2	4
772	Accumulation of 2,3,7,8-tetrachlorodibenzo-p-dioxin by rainbow trout (<i>Onchorhynchus mykiss</i>) at environmentally relevant dietary concentrations. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 344-50.	2.2	4
773	Polychlorinated-naphthalenes, -biphenyls, -dibenzo-p-dioxins, -dibenzofurans and p,p'-DDE in bluefin tuna, swordfish, cormorants and barn swallows from Italy. <i>Ambio</i> , 2002, 31, 207-11.	2.8	4
774	Identification of AhR agonists in sediments of the Bohai and Yellow Seas using advanced effect-directed analysis and in silico prediction. <i>Journal of Hazardous Materials</i> , 2022, 435, 128908.	6.5	4

#	ARTICLE	IF	CITATIONS
775	Free Amino Acid Pools of Five Species of Freshwater Oligochaetes. Canadian Journal of Fisheries and Aquatic Sciences, 1986, 43, 600-607.	0.7	3
776	Characterization studies of a semi-automated separation method for analysis of non-ortho-substituted polychlorinated biphenyl (PCB) congeners in environmental samples. Toxicological and Environmental Chemistry, 1995, 51, 229-241.	0.6	3
777	Risk Assessment of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin Equivalents in Tissue Samples from Three Species in the Denver Metropolitan Area. Human and Ecological Risk Assessment (HERA), 2000, 6, 1087-1099.	1.7	3
778	Response to Comment on "Occurrence of Butyltin Compounds in Human Blood". Environmental Science & Technology, 2000, 34, 1879-1880.	4.6	3
779	ECOLOGICAL RISK ASSESSMENT OF PESTICIDES. Human and Ecological Risk Assessment (HERA), 2001, 7, 493-495.	1.7	3
780	An automated enantioselective isolation system for the study of estrogenic potencies: Study of the estrogenic activity of \pm -hexachlorocyclohexane. Journal of Separation Science, 2003, 26, 903-907.	1.3	3
781	Nondestructive Scat Sampling in Assessment of Mink (<i>Mustela vison</i>) Exposed to Polychlorinated Dibenzofurans (PCDFs). Archives of Environmental Contamination and Toxicology, 2008, 55, 529-537.	2.1	3
782	Great horned owl (<i>Bubo virginianus</i>) dietary exposure to PCDD/DF in the Tittabawassee River floodplain in Midland, Michigan, USA. Environmental Toxicology and Chemistry, 2010, 29, 2350-2362.	2.2	3
783	Incidence of jaw lesions and activity and gene expression of hepatic P4501A enzymes in mink (<i>Mustela vison</i>) exposed to dietary 2,3,7,8-tetrachlorodibenzo-p-dioxin, 2,3,7,8-tetrachlorodibenzofuran, and 2,3,4,7,8-pentachlorodibenzofuran. Environmental Toxicology and Chemistry, 2012, 31, 2545-2556.	2.2	3
784	Ecotoxicology of organochlorine chemicals in birds of the great lakes. Environmental Toxicology and Chemistry, 2013, 32, 490-492.	2.2	3
785	Tissue Residue Guideline for \sim DDT for Protection of Aquatic Birds in China. Human and Ecological Risk Assessment (HERA), 2014, 20, 1629-1642.	1.7	3
786	A Reagent-Free Screening Assay for Evaluation of the Effects of Chemicals on the Proliferation and Morphology of HeLa-GFP Cells. Environmental Science and Technology Letters, 2016, 3, 322-326.	3.9	3
787	Response to Comment on "Mutagenic Azo Dyes, Rather than Flame Retardants, are the Predominant Brominated Compounds in House Dust". Environmental Science & Technology, 2017, 51, 3591-3592.	4.6	3
788	Receptor-mediated potencies of polycyclic aromatic hydrocarbons in urban sediments: comparisons of toxic equivalency risk assessment. International Journal of Environmental Science and Technology, 2019, 16, 6405-6418.	1.8	3
789	Remodeling of Arctic char (<i>Salvelinus alpinus</i>) lipidome under a stimulated scenario of Arctic warming. Global Change Biology, 2021, 27, 3282-3298.	4.2	3
790	Health status of fathead minnow (<i>Pimephales promelas</i>) populations in a municipal wastewater effluent-dominated stream in the Canadian prairies, Wascana Creek, Saskatchewan. Aquatic Toxicology, 2021, 238, 105933.	1.9	3
791	Occurrence, toxicity and ecological risk of larvicidal oil in the coastal marine ecosystem of Hong Kong. Marine Pollution Bulletin, 2020, 156, 111178.	2.3	3
792	Insights into the Influence of Natural Retinoic Acids on Imposex Induction in Female Marine Gastropods in the Coastal Environment. Environmental Science and Technology Letters, 2021, 8, 1002-1008.	3.9	3

#	ARTICLE	IF	CITATIONS
793	Role of endocrine disruption in toxicity of 6-benzylaminopurine (6-BA) to early-life stages of Zebrafish. <i>Ecotoxicology and Environmental Safety</i> , 2022, 232, 113287.	2.9	3
794	Effects induced by feeding organochlorine-contaminated carp from Saginaw Bay, Lake Huron, to laying White Leghorn hens. II. Embryotoxic and teratogenic effects. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 1996, 49, 409-38.	1.1	3
795	Dioxin-like and non-dioxin-like toxic effects of polychlorinated biphenyls (PCBs): implications for risk assessment. <i>Central European Journal of Public Health</i> , 2000, 8 Suppl, 43-5.	0.4	3
796	Ecotoxicological Risk Assessment of Atrazine in Amphibians. <i>ACS Symposium Series</i> , 2005, , 124-137.	0.5	2
797	Determinations of dioxinlike activity in selected mollusks from the coast of the Bohai Sea, China, using the H4IIE-luc bioassay. <i>Ecotoxicology and Environmental Safety</i> , 2007, 67, 157-162.	2.9	2
798	Risk assessment methodologies for exposure of great horned owls (<i>Bubo virginianus</i>) to PCBs on the Kalamazoo river, Michigan. <i>Integrated Environmental Assessment and Management</i> , 2008, 4, 24-40.	1.6	2
799	Dietary exposure of great blue heron (<i>Ardea herodias</i>) to PCDD/DFs in the Tittabawassee River floodplain, MI, USA. <i>Ecotoxicology and Environmental Safety</i> , 2011, 74, 494-503.	2.9	2
800	Reproductive success of three passerine species exposed to dioxin-like compounds near Midland, Michigan, USA. <i>Ecotoxicology</i> , 2012, 21, 1145-1154.	1.1	2
801	Establishment of a three-step method to evaluate effects of chemicals on development of zebrafish embryo/larvae. <i>Chemosphere</i> , 2017, 186, 209-217.	4.2	2
802	Correlations between slow pyrolysis characteristics and organic carbon structure of aquatic plant biomass. <i>Environmental Science and Pollution Research</i> , 2019, 26, 17555-17566.	2.7	2
803	Using zooplankton metabarcoding to assess the efficacy of different techniques to clean-up an oil-spill in a boreal lake. <i>Aquatic Toxicology</i> , 2021, 236, 105847.	1.9	2
804	Prefertilization Exposure of Rainbow Trout Eggs to Perâ€and Polyfluoroalkyl Substances to Simulate Accumulation During Oogenesis. <i>Environmental Toxicology and Chemistry</i> , 2021, 40, 3159-3165.	2.2	2
805	HYDROXYLATED AND METHYLSULFONYL POLYCHLORINATED BIPHENYL METABOLITES IN ALBATROSSES FROM MIDWAY ATOLL, NORTH PACIFIC OCEAN. <i>Environmental Toxicology and Chemistry</i> , 1998, 17, 1620.	2.2	2
806	SETAC: PART OF THE SOLUTION OR PART OF THE PROBLEM?. <i>Environmental Toxicology and Chemistry</i> , 1990, 9, 1327.	2.2	2
807	RNA metabarcoding helps reveal zooplankton community response to environmental stressors. <i>Environmental Pollution</i> , 2022, 292, 118446.	3.7	2
808	Assessment of combined exposures to multiple chemicals: pesticides, metals, and polycyclic aromatic hydrocarbons levels in fig fruits. <i>International Journal of Environmental Analytical Chemistry</i> , 2024, 104, 827-846.	1.8	2
809	Laboratory analyses of the potential toxicity of sediment-associated polydimethylsiloxane to benthic macroinvertebrates. <i>Environmental Toxicology and Chemistry</i> , 2001, 20, 2611-6.	2.2	2
810	Temporal Patterns of Bacterial and Viral Communities during Algae Blooms of a Reservoir in Macau. <i>Toxins</i> , 2021, 13, 894.	1.5	2

#	ARTICLE	IF	CITATIONS
811	Best available technique for the recovery of marine benthic communities in a gravel shore after the oil spill: A mesocosm-based sediment triad assessment. <i>Journal of Hazardous Materials</i> , 2022, 435, 128945.	6.5	2
812	Studies of Adenine Nucleotide Biochemistry in the Chediak-Higashi Syndrome. <i>Experimental and Molecular Pathology</i> , 1993, 58, 40-52.	0.9	1
813	Transcriptional changes in African clawed frogs (<i>Xenopus laevis</i>) exposed to 17 β -ethynylestradiol during early development. <i>Ecotoxicology</i> , 2015, 24, 321-329.	1.1	1
814	Relative sensitivities among avian species to individual and mixtures of aryl hydrocarbon receptor-â€œactive compounds. <i>Environmental Toxicology and Chemistry</i> , 2016, 35, 1239-1246.	2.2	1
815	Investigation of eluted characteristics of fulvic acids using differential spectroscopy combined with Gaussian deconvolution and spectral indices. <i>Environmental Science and Pollution Research</i> , 2020, 27, 11000-11011.	2.7	1
816	Changes in cytochrome P4501A activity during development in common tern chicks fed polychlorinated biphenyls, as measured by the caffeine breath test. , 2000, 19, 712.		1
817	In vitro assessment of potential mechanism-specific effects of polybrominated diphenyl ethers. , 2002, 21, 2431.		1
818	Influence of Environmental Variables on Benthic Macroinvertebrate Communities in a Shallow Eutrophic Lowland Lake (Ge Lake, China). <i>Tecnologia Y Ciencias Del Agua</i> , 2019, 10, 88-119.	0.1	1
819	A framework for assessing freshwater vulnerability along China's Belt and Road Initiative: An exposure, sensitivity and adaptive capacity approach. <i>Environmental Science and Policy</i> , 2022, 132, 247-261.	2.4	1
820	Physicochemical Characterization and Assessment of Magnitude of Pollution to Contribute to Water Sustainability. <i>Sustainability</i> , 2022, 14, 6689.	1.6	1
821	Science-Based decision making to reduce risks from persistent organic pollutants (POPS) joint workshop of the chinese academy of sciences, SCOPE-China and the US National Academies Beijing, Peoples Republic of China, June 7â€œ10. <i>Environmental Science and Pollution Research</i> , 2004, 11, 378-378.	2.7	0
822	Monitoring of Exposure to and Potential Effects of Contaminants in the Environment. <i>Environmental Bioindicators</i> , 2007, 2, 129-130.	0.4	0
823	SETAC: Part of the solution or part of the problem?. <i>Environmental Toxicology and Chemistry</i> , 2010, 9, 1327-1330.	2.2	0
824	Multiple lines of evidence risk assessment of american robins exposed to polychlorinated dibenzofurans (PCDFS) and polychlorinated dibenzoâ€œdioxins (PCDDS) in the tittabawassee river floodplain, Midland, Michigan, Usa. <i>Environmental Toxicology and Chemistry</i> , 2013, 32, 1304-1316.	2.2	0
825	The effect of IPC formulation on bitumen properties â€œ An experimental study. <i>Journal of Petroleum Science and Engineering</i> , 2018, 170, 525-540.	2.1	0
826	A novel Mg(OH) ₂ binding layer-based DGT technique for measuring phosphorus in water and sediment. <i>Environmental Sciences: Processes and Impacts</i> , 2020, 22, 340-349.	1.7	0
827	In memory of Dr. Doris Au (29 April 1965â€œ7 February 2020). <i>Marine Pollution Bulletin</i> , 2021, 167, 112278.	2.3	0
828	Does the African potato (<i>Hypoxis hemerocallidea</i>) activate the aryl hydrocarbon receptor in H4IIE-luc cells?. , 0, , .		0

#	ARTICLE	IF	CITATIONS
829	2,3,7,8-Tetrachlorodibenzo-p-dioxin equivalents in tissue samples from three species in the Denver, Colorado, USA, metropolitan area. Environmental Toxicology and Chemistry, 2001, 20, 2433-42.	2.2	0