## **Denis Feliers**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8115661/publications.pdf Version: 2024-02-01



NENIS FELIEDS

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Proximal tubular epithelial insulin receptor mediates high-fat diet–induced kidney injury. JCI Insight,<br>2021, 6, .   | 5.0 | 8         |
| 2  | Chloride channel accessory 1 integrates chloride channel activity and mTORC1 in agingâ€related kidney injury. Aging Cell, 2021, 20, e13407.   | 6.7 | 11        |
| 3  | Interplay between RNA-binding protein HuR and Nox4 as a novel therapeutic target in diabetic kidney<br>disease. Molecular Metabolism, 2020, 36, 100968.   | 6.5 | 35        |
| 4  | Marmoset as a Model to Study Kidney Changes Associated With Aging. Journals of Gerontology - Series<br>A Biological Sciences and Medical Sciences, 2019, 74, 315-324.   | 3.6 | 19        |
| 5  | A Multi-Parameter Analysis of Cellular Coordination of Major Transcriptome Regulation Mechanisms.<br>Scientific Reports, 2018, 8, 5742.   | 3.3 | 7         |
| 6  | Hydrogen sulfide ameliorates aging-associated changes in the kidney. GeroScience, 2018, 40, 163-176.  | 4.6 | 49        |
| 7  | Neutrophil peptidyl arginine deiminase-4 has a pivotal role in ischemia/reperfusion-induced<br>acuteÂkidney injury. Kidney International, 2018, 93, 365-374.  | 5.2 | 116       |
| 8  | Hydrogen sulfide as a regulatory factor in kidney health and disease. Biochemical Pharmacology, 2018,<br>149, 29-41.  | 4.4 | 34        |
| 9  | An electrochemically deposited collagen wound matrix combined with adipose-derived stem cells<br>improves cutaneous wound healing in a mouse model of type 2 diabetes. Journal of Biomaterials<br>Applications, 2018, 33, 553-565.              | 2.4 | 13        |
| 10 | Hydrogen sulfide inhibits high glucose-induced NADPH oxidase 4 expression and matrix increase by recruiting inducible nitric oxide synthase in kidney proximal tubular epithelial cells. Journal of Biological Chemistry, 2017, 292, 5665-5675. | 3.4 | 40        |
| 11 | Spleen contributes significantly to increased circulating levels of fibroblast growth factor 23 in response to lipopolysaccharide-induced inflammation. Nephrology Dialysis Transplantation, 2017, 32, 960-968.                                 | 0.7 | 25        |
| 12 | Hydrogen Sulfide in Renal Physiology and Disease. Antioxidants and Redox Signaling, 2016, 25, 720-731.  | 5.4 | 82        |
| 13 | Rapamycin Increases Mortality in <i>db/db</i> Mice, a Mouse Model of Type 2 Diabetes. Journals of<br>Gerontology - Series A Biological Sciences and Medical Sciences, 2016, 71, 850-857.  | 3.6 | 57        |
| 14 | Metformin prevents renal interstitial fibrosis in mice with unilateral ureteral obstruction.<br>Molecular and Cellular Endocrinology, 2015, 412, 116-122.   | 3.2 | 59        |
| 15 | Tadalafil Integrates Nitric Oxide-Hydrogen Sulfide Signaling to Inhibit High Glucose-induced Matrix<br>Protein Synthesis in Podocytes. Journal of Biological Chemistry, 2015, 290, 12014-12026.   | 3.4 | 38        |
| 16 | Epigenetic control of podocyte differentiation: a new target of the renin–angiotensin system in kidney disease. Kidney International, 2015, 88, 668-670.  | 5.2 | 8         |
| 17 | Symmetric dimethylarginine alters endothelial nitric oxide activity in glomerular endothelial cells.<br>Cellular Signalling, 2015, 27, 1-5.   | 3.6 | 28        |
| 18 | Stabilization of HIF-2α through redox regulation of mTORC2 activation and initiation of mRNA translation. Oncogene, 2013, 32, 3147-3155.  | 5.9 | 47        |

**DENIS FELIERS** 

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Apelin retards the progression of diabetic nephropathy. American Journal of Physiology - Renal<br>Physiology, 2013, 304, F788-F800.  | 2.7 | 86        |
| 20 | Hydrogen Sulfide Inhibits High Glucose-induced Matrix Protein Synthesis by Activating AMP-activated<br>Protein Kinase in Renal Epithelial Cells. Journal of Biological Chemistry, 2012, 287, 4451-4461.                    | 3.4 | 108       |
| 21 | Molecular events in matrix protein metabolism in the aging kidney. Aging Cell, 2012, 11, 1065-1073.  | 6.7 | 38        |
| 22 | Erk in Kidney Diseases. Journal of Signal Transduction, 2011, 2011, 1-8.   | 2.0 | 37        |
| 23 | The complex world of kidney microRNAs. Kidney International, 2011, 80, 334-337.  | 5.2 | 31        |
| 24 | Resveratrol ameliorates high glucose-induced protein synthesis in glomerular epithelial cells.<br>Cellular Signalling, 2010, 22, 65-70.  | 3.6 | 42        |
| 25 | Acute hyperglycemia rapidly stimulates VEGF mRNA translation in the kidney. Role of angiotensin type 2 receptor (AT2). Cellular Signalling, 2010, 22, 1849-1857.   | 3.6 | 23        |
| 26 | Mechanism of VEGF expression by high glucose in proximal tubule epithelial cells. Molecular and<br>Cellular Endocrinology, 2010, 314, 136-142.   | 3.2 | 27        |
| 27 | Vascular endothelial growth factor as a prognostic marker of lupus nephritis. Kidney International,<br>2009, 75, 1251-1253.  | 5.2 | 6         |
| 28 | Regulation of mRNA translation in renal physiology and disease. American Journal of Physiology -<br>Renal Physiology, 2009, 297, F1153-F1165.  | 2.7 | 52        |
| 29 | Novel mechanisms of protein synthesis in diabetic nephropathy—role of mRNA translation. Reviews in<br>Endocrine and Metabolic Disorders, 2008, 9, 255-266.   | 5.7 | 18        |
| 30 | Raptor-rictor axis in TGFÎ <sup>2</sup> -induced protein synthesis. Cellular Signalling, 2008, 20, 409-423.  | 3.6 | 60        |
| 31 | PKCδ regulates the stimulation of vascular endothelial factor mRNA translation by angiotensin II<br>through hnRNP K. Cellular Signalling, 2008, 20, 969-977.   | 3.6 | 31        |
| 32 | High Glucose, High Insulin, and Their Combination Rapidly Induce Laminin-β1 Synthesis by Regulation of mRNA Translation in Renal Epithelial Cells. Diabetes, 2007, 56, 476-485.  | 0.6 | 71        |
| 33 | A role for AMP-activated protein kinase in diabetes-induced renal hypertrophy. American Journal of<br>Physiology - Renal Physiology, 2007, 292, F617-F627.   | 2.7 | 253       |
| 34 | Heterogeneous nuclear ribonucleoprotein K contributes to angiotensin II stimulation of vascular<br>endothelial growth factor mRNA translation. American Journal of Physiology - Renal Physiology,<br>2007, 293, F607-F615. | 2.7 | 30        |
| 35 | Regulation of Elongation Phase of mRNA Translation in Diabetic Nephropathy. American Journal of<br>Pathology, 2007, 171, 1733-1742.  | 3.8 | 114       |
| 36 | Angiotensin II stimulation of VEGF mRNA translation requires production of reactive oxygen species.<br>American Journal of Physiology - Renal Physiology, 2006, 290, F927-F936.  | 2.7 | 68        |

Denis Feliers

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | mRNA Translation. Journal of the American Society of Nephrology: JASN, 2006, 17, 3281-3292.   | 6.1 | 56        |
| 38 | mRNA Translation in Diabetic Nephropathy. , 2006, , 97-116.   |     | 3         |
| 39 | VEGF regulation of endothelial nitric oxide synthase in glomerular endothelial cells. Kidney<br>International, 2005, 68, 1648-1659.   | 5.2 | 118       |
| 40 | Translational regulation of vascular endothelial growth factor expression in renal epithelial cells<br>by angiotensin II. American Journal of Physiology - Renal Physiology, 2005, 288, F521-F529.  | 2.7 | 45        |
| 41 | Cloning of the 5'-flanking region of the murine bone morphogenetic protein-7 gene. Molecular and<br>Cellular Biochemistry, 2002, 233, 31-37.  | 3.1 | 7         |
| 42 | Activation of Cyclin D1-Cdk4 and Cdk4-Directed Phosphorylation of RB Protein in Diabetic Mesangial<br>Hypertrophy. Diabetes, 2002, 51, 3290-3299.   | 0.6 | 37        |
| 43 | Insulin regulation of protein translation repressor 4E-BP1, an eIF4E-binding protein, in renal epithelial cells. Kidney International, 2001, 59, 866-875.   | 5.2 | 73        |
| 44 | Activation of renal signaling pathways in db/db mice with type 2 diabetes. Kidney International, 2001, 60, 495-504.   | 5.2 | 94        |
| 45 | Angiotensin II activates Akt/protein kinase B by an arachidonic acid/redoxâ€dependent pathway and<br>independent of phosphoinositide 3â€kinase. FASEB Journal, 2001, 15, 1909-1920.   | 0.5 | 99        |
| 46 | Insulin regulation of protein translation repressor 4E-BP1, an eIF4E-binding protein, in renal epithelial cells. Kidney International, 2001, 59, 866-875.   | 5.2 | 7         |
| 47 | Potential role of insulin-like growth factor binding protein-4 in the uncoupling of bone turnover in multiple myeloma. British Journal of Haematology, 1999, 104, 715-722.  | 2.5 | 16        |
| 48 | Characterization and regulation of insulin-like growth factor binding proteins in human hepatic stellate cells. , 1998, 174, 240-250.   |     | 32        |
| 49 | Inhibitory Effect of Annexin V on Protein Kinase C Activity in Mesangial Cell Lysates. FEBS Journal, 1995,<br>232, 865-872.   | 0.2 | 14        |
| 50 | Inhibitory Effect of Annexin V on Protein Kinase C Activity in Mesangial Cell Lysates. FEBS Journal, 1995,<br>232, 865-872.   | 0.2 | 2         |
| 51 | Inhibitory Effect of Annexin V on Protein Kinase C Activity in Mesangial Cell Lysates. FEBS Journal, 1995,<br>232, 865-872.   | 0.2 | 25        |
| 52 | Species Differences of the Thyroid Protein Kinase C Heterogeneity. Thyroid, 1994, 4, 459-465.   | 4.5 | 5         |
| 53 | Forskolin mimics TSH action on the expression of protein kinase C isozymes in pig thyroid cell cultures. Cellular Signalling, 1994, 6, 513-522.   | 3.6 | 1         |
| 54 | Immunological identification of protein kinase C-α and protein kinase C-Î′ in cultured rat mesangial cells:<br>Diffferential sensitivity of the two isoforms towards the protein kinase inhibitor H7. Cellular<br>Signalling, 1992, 4, 559-569. | 3.6 | 13        |

**DENIS FELIERS** 

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Heterogeneity of protein kinase C in cultured rat mesangial cells. Cellular Signalling, 1992, 4, 179-188.              | 3.6 | 5         |
| 56 | Characterization and regulation of insulin-like growth factor binding proteins in human hepatic stellate cells. , 0, . |     | 1         |