

# Frauke Schultze-Lutter

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8117921/frauke-schultze-lutter-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

170  
papers

8,996  
citations

51  
h-index

92  
g-index

190  
ext. papers

10,739  
ext. citations

4.5  
avg, IF

6.26  
L-index

#	Paper	IF	Citations
170	Does childhood trauma predict schizotypal traits? A path modelling approach in a cohort of help-seeking subjects.. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2022</b> , 1	5.1	1
169	Relationships between global functioning and neuropsychological predictors in subjects at high risk of psychosis or with a recent onset of depression.. <i>World Journal of Biological Psychiatry</i> , <b>2022</b> , 1-9	3.8	0
168	Using combined environmental-clinical classification models to predict role functioning outcome in clinical high-risk states for psychosis and recent-onset depression.. <i>British Journal of Psychiatry</i> , <b>2022</b> , 1-17	5.4	0
167	Clinical high-risk criteria of psychosis in 8-17-year-old community subjects and inpatients not suspected of developing psychosis.. <i>World Journal of Psychiatry</i> , <b>2022</b> , 12, 425-449	3	0
166	Pattern of predictive features of continued cannabis use in patients with recent-onset psychosis and clinical high-risk for psychosis.. <i>NPJ Schizophrenia</i> , <b>2022</b> , 8, 19	5.5	
165	MR-Spectroscopy of GABA and Glutamate/Glutamine Concentrations in Auditory Cortex in Clinical High-Risk for Psychosis Individuals.. <i>Frontiers in Psychiatry</i> , <b>2022</b> , 13, 859322	5	
164	Psychometric properties of the Kessler psychological scales in a Swiss young-adult community sample indicate poor suitability for community screening for mental disorders.. <i>Microbial Biotechnology</i> , <b>2022</b> ,	3.3	1
163	(Attenuated) hallucinations join basic symptoms in a transdiagnostic network cluster analysis.. <i>Schizophrenia Research</i> , <b>2022</b> , 243, 43-54	3.6	2
162	Computerised cognitive training during early-stage psychosis improves cognitive deficits and gamma-band oscillations: A pilot study.. <i>Schizophrenia Research</i> , <b>2022</b> , 243, 217-219	3.6	
161	Prevalence and predictors of suicidality and non-suicidal self-harm among individuals at clinical high-risk for psychosis: Results from a community-recruited sample. <i>Microbial Biotechnology</i> , <b>2021</b> , 15, 1256-1265	3.3	2
160	Positive psychotic symptoms in childhood and adolescence.. <i>Current Opinion in Psychology</i> , <b>2021</b> , 45, 101287	6.2	1
159	Detailed clinical phenotyping and generalisability in prognostic models of functioning in at-risk populations.. <i>British Journal of Psychiatry</i> , <b>2021</b> , 1-4	5.4	
158	Basic symptoms and gray matter volumes of patients at clinical high risk for psychosis. <i>Psychological Medicine</i> , <b>2021</b> , 51, 2666-2674	6.9	1
157	Association between age of cannabis initiation and gray matter covariance networks in recent onset psychosis. <i>Neuropsychopharmacology</i> , <b>2021</b> , 46, 1484-1493	8.7	5
156	Cognitive subtypes in recent onset psychosis: distinct neurobiological fingerprints?. <i>Neuropsychopharmacology</i> , <b>2021</b> , 46, 1475-1483	8.7	2
155	Multimodal prognosis of negative symptom severity in individuals at increased risk of developing psychosis. <i>Translational Psychiatry</i> , <b>2021</b> , 11, 312	8.6	3
154	The relationship between cognitive deficits and impaired short-term functional outcome in clinical high-risk for psychosis participants: A machine learning and modelling approach. <i>Schizophrenia Research</i> , <b>2021</b> , 231, 24-31	3.6	3

153	The Important Role of Stereotypes in the relation between Mental Health Literacy and Stigmatization of Depression and Psychosis in the Community. <i>Community Mental Health Journal</i> , <b>2021</b> , 1	2.1	0
152	The Bern Early Recognition and Intervention Centre for mental crisis (FETZ Bern)-An 8-year evaluation. <i>Microbial Biotechnology</i> , <b>2021</b> ,	3.3	1
151	Is there a diagnosis-specific influence of childhood trauma on later educational attainment? A machine learning analysis in a large help-seeking sample. <i>Journal of Psychiatric Research</i> , <b>2021</b> , 138, 591-597	5.3	5
150	Schizotypie und Schizotype (Persönlichkeits-)Störung. <i>PTT Persönlichkeitsstörungen Theorie Und Therapie</i> , <b>2021</b> , 25, 96-109	0.1	
149	Towards clinical application of prediction models for transition to psychosis: A systematic review and external validation study in the PRONIA sample. <i>Neuroscience and Biobehavioral Reviews</i> , <b>2021</b> , 125, 478-492	9	13
148	Toward Generalizable and Transdiagnostic Tools for Psychosis Prediction: An Independent Validation and Improvement of the NAPLS-2 Risk Calculator in the Multisite PRONIA Cohort. <i>Biological Psychiatry</i> , <b>2021</b> , 90, 632-642	7.9	8
147	The Psychopathology and Neuroanatomical Markers of Depression in Early Psychosis. <i>Schizophrenia Bulletin</i> , <b>2021</b> , 47, 249-258	1.3	9
146	Clinical high-risk of and conversion to psychosis in the community: A 3-year follow-up of a cohort study. <i>Schizophrenia Research</i> , <b>2021</b> , 228, 616-618	3.6	3
145	Heterogeneity and Classification of Recent Onset Psychosis and Depression: A Multimodal Machine Learning Approach. <i>Schizophrenia Bulletin</i> , <b>2021</b> , 47, 1130-1140	1.3	3
144	Multimodal Machine Learning Workflows for Prediction of Psychosis in Patients With Clinical High-Risk Syndromes and Recent-Onset Depression. <i>JAMA Psychiatry</i> , <b>2021</b> , 78, 195-209	14.5	31
143	The non-specific nature of mental health and structural brain outcomes following childhood trauma. <i>Psychological Medicine</i> , <b>2021</b> , 1-10	6.9	1
142	Duration of basic and attenuated-psychotic symptoms in individuals at clinical high risk for psychosis: pattern of symptom onset and effects of duration on functioning and cognition. <i>BMC Psychiatry</i> , <b>2021</b> , 21, 339	4.2	1
141	Characterising cognitive heterogeneity in individuals at clinical high-risk for psychosis: a cluster analysis with clinical and functional outcome prediction. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2021</b> , 1	5.1	2
140	Predictors of help-seeking behaviour in people with mental health problems: a 3-year prospective community study. <i>BMC Psychiatry</i> , <b>2021</b> , 21, 432	4.2	2
139	Novel Gyrification Networks Reveal Links with Psychiatric Risk Factors in Early Illness. <i>Cerebral Cortex</i> , <b>2021</b> ,	5.1	1
138	40-Hz Auditory Steady-State Responses Characterize Circuit Dysfunctions and Predict Clinical Outcomes in Clinical High-Risk for Psychosis Participants: A Magnetoencephalography Study. <i>Biological Psychiatry</i> , <b>2021</b> , 90, 419-429	7.9	11
137	The clinical relevance of formal thought disorder in the early stages of psychosis: results from the PRONIA study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2021</b> , 1	5.1	0
136	Traces of Trauma: A Multivariate Pattern Analysis of Childhood Trauma, Brain Structure, and Clinical Phenotypes. <i>Biological Psychiatry</i> , <b>2020</b> , 88, 829-842	7.9	15

135 Subjective disturbances in emerging psychosis **2020**, 59-80

134	Main Symptomatic Treatment Targets in Suspected and Early Psychosis: New Insights From Network Analysis. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, 884-895	1.3	13
133	Association of Magnetoencephalographically Measured High-Frequency Oscillations in Visual Cortex With Circuit Dysfunctions in Local and Large-scale Networks During Emerging Psychosis. <i>JAMA Psychiatry</i> , <b>2020</b> , 77, 852-862	14.5	15
132	Basic Symptoms Are Associated With Age in Patients With a Clinical High-Risk State for Psychosis: Results From the PRONIA Study. <i>Frontiers in Psychiatry</i> , <b>2020</b> , 11, 552175	5	2
131	Effects of age and sex on clinical high-risk for psychosis in the community. <i>World Journal of Psychiatry</i> , <b>2020</b> , 10, 101-124	3	11
130	Validation of the Bullying Scale for Adults - Results of the PRONIA-study. <i>Journal of Psychiatric Research</i> , <b>2020</b> , 129, 88-97	5.2	4
129	General psychopathology links burden of recent life events and psychotic symptoms in a network approach. <i>NPJ Schizophrenia</i> , <b>2020</b> , 6, 40	5.5	8
128	Grey-matter abnormalities in clinical high-risk participants for psychosis. <i>Schizophrenia Research</i> , <b>2020</b> , 226, 120-128	3.6	1
127	T223. MULTIVARIATE PREDICTION OF FOLLOW UP SOCIAL AND OCCUPATIONAL OUTCOME IN CLINICAL HIGH-RISK INDIVIDUALS BASED ON GRAY MATTER VOLUMES AND HISTORY OF ENVIRONMENTAL ADVERSE EVENTS. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S317-S318	1.3	78
126	M1. INVESTIGATING THE RELATIONSHIP BETWEEN CHILDHOOD TRAUMA AND PSYCHIATRIC DISEASE USING MACHINE LEARNING TECHNIQUES. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S133-S133	1.3	78
125	S94. PREDICTION OF CANNABIS RELAPSE IN CLINICAL HIGH-RISK INDIVIDUALS AND RECENT ONSET PSYCHOSIS - PRELIMINARY RESULTS FROM THE PRONIA STUDY. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S69-S70	1.3	78
124	Trapped in a Glass Bell Jar: Neural Correlates of Depersonalization and Derealization in Subjects at Clinical High-Risk of Psychosis and Depersonalization-Derealization Disorder. <i>Frontiers in Psychiatry</i> , <b>2020</b> , 11, 535652	5	2
123	M121. CLINICAL PREDICTION MODELS FOR TRANSITION TO PSYCHOSIS: AN EXTERNAL VALIDATION STUDY IN THE PRONIA SAMPLE. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S181-S181	1.3	78
122	Altered Autonomic Function in Individuals at Clinical High Risk for Psychosis. <i>Frontiers in Psychiatry</i> , <b>2020</b> , 11, 580503	5	1
121	O6.6. MULTIMODAL PROGNOSIS OF NEGATIVE SYMPTOM SEVERITY IN INDIVIDUALS WITH INCREASED RISK OF DEVELOPING PSYCHOSIS. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S15-S16	1.3	78
120	O10.5. CLINICAL HIGH RISK (CHR) CRITERIA IN CHILDREN AND ADOLESCENTS NOT SUSPECTED TO DEVELOP PSYCHOSIS: NO TRANSDIAGNOSTIC SEVERITY MARKER OF MENTAL HEALTH PROBLEMS. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S26-S26	1.3	78
119	O6.4. ASSOCIATION BETWEEN CLUSTERS OF FORMAL THOUGHT DISORDERS SEVERITY AND NEUROCOGNITIVE AND FUNCTIONAL OUTCOME INDICES IN THE EARLY STAGES OF PSYCHOSIS □ RESULTS FROM THE PRONIA COHORT. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S14-S15	1.3	78
118	S219. SINGLE-SUBJECT PREDICTION OF FUNCTIONAL OUTCOMES ACROSS DIAGNOSTIC GROUPS USING CLINICAL DATA. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S122-S122	1.3	78

117	O8.5. SIGNS OF ADVERSITY - A NOVEL MACHINE LEARNING APPROACH TO CHILDHOOD TRAUMA, BRAIN STRUCTURE AND CLINICAL PROFILES. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S20-S20	1.3	78
116	S64. COGNITIVE IMPAIRMENTS AND PREDICTION OF FUNCTIONAL OUTCOME IN INDIVIDUALS AT CLINICAL HIGH-RISK FOR PSYCHOSIS. <i>Schizophrenia Bulletin</i> , <b>2020</b> , 46, S57-S58	1.3	78
115	Age effects on basic symptoms in the community: A route to gain new insight into the neurodevelopment of psychosis?. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2020</b> , 270, 311-324	5.1	11
114	Childhood physical abuse and emotional neglect are specifically associated with adult mental disorders. <i>Journal of Mental Health</i> , <b>2020</b> , 29, 376-384	2.7	18
113	Neuropsychological deficits in participants at clinical high risk for psychosis recruited from the community: relationships to functioning and clinical symptoms. <i>Psychological Medicine</i> , <b>2020</b> , 50, 77-85	6.9	8
112	Sex differences in symptomatology of psychosis-risk patients and in prediction of psychosis. <i>Archives of Womens Mental Health</i> , <b>2020</b> , 23, 339-349	5	5
111	Basic symptoms in offspring of parents with mood and psychotic disorders. <i>BJPsych Open</i> , <b>2019</b> , 5, e54	5	7
110	The interrelationship between schizotypy, clinical high risk for psychosis and related symptoms: Cognitive disturbances matter. <i>Schizophrenia Research</i> , <b>2019</b> , 210, 188-196	3.6	19
109	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , <b>2019</b> , 10, 345	5	18
108	Psychosis and Schizophrenia-Spectrum Personality Disorders Require Early Detection on Different Symptom Dimensions. <i>Frontiers in Psychiatry</i> , <b>2019</b> , 10, 476	5	27
107	Using Online Screening in the General Population to Detect Participants at Clinical High-Risk for Psychosis. <i>Schizophrenia Bulletin</i> , <b>2019</b> , 45, 600-609	1.3	32
106	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical high-risk patients of psychosis. Results of the EPOS project. <i>Microbial Biotechnology</i> , <b>2019</b> , 13, 935-942	3.3	7
105	Coping as a mediator between locus of control, competence beliefs, and mental health: A systematic review and structural equation modelling meta-analysis. <i>Behaviour Research and Therapy</i> , <b>2019</b> , 121, 103442	5.2	26
104	Associations of psychosis-risk symptoms with quality of life and self-rated health in the Community. <i>European Psychiatry</i> , <b>2019</b> , 62, 116-123	6	4
103	Functional and structural correlates of abnormal involuntary movements in psychosis risk and first episode psychosis. <i>Schizophrenia Research</i> , <b>2019</b> , 212, 196-203	3.6	9
102	Clinical High Risk for Psychosis Syndromes Among Swiss and German Youth and Young Adults: Early Identification and Intervention <b>2019</b> , 115-142		
101	Prevalence, course and psychosis-predictive value of negative symptoms in 22q11.2 deletion syndrome. <i>Schizophrenia Research</i> , <b>2019</b> , 206, 386-393	3.6	10
100	The trait-state distinction between schizotypy and clinical high risk: results from a one-year follow-up. <i>World Psychiatry</i> , <b>2019</b> , 18, 108-109	14.4	13

99	Influence of demographic characteristics on attenuated positive psychotic symptoms in a young, help-seeking, at-risk population. <i>Microbial Biotechnology</i> , <b>2019</b> , 13, 53-56	3.3	6
98	Expressed emotion as a predictor of the first psychotic episode - Results of the European prediction of psychosis study. <i>Schizophrenia Research</i> , <b>2018</b> , 199, 346-352	3.6	9
97	Psychosis-risk criteria in the general population: frequent misinterpretations and current evidence. <i>World Psychiatry</i> , <b>2018</b> , 17, 107-108	14.4	14
96	The Frankfurt Complaint Questionnaire for self-assessment of basic symptoms in the early detection of psychosis-Factor structure, reliability, and predictive validity. <i>International Journal of Methods in Psychiatric Research</i> , <b>2018</b> , 27, e1600	4.3	6
95	Demographic and clinical characteristics of diagnosed and non-diagnosed psychotic disorders in the community. <i>Microbial Biotechnology</i> , <b>2018</b> , 12, 87-90	3.3	5
94	Course of clinical high-risk states for psychosis beyond conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2018</b> , 268, 39-48	5.1	43
93	Increased Striatal and Reduced Prefrontal Cerebral Blood Flow in Clinical High Risk for Psychosis. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, 182-192	1.3	22
92	Prevalence and clinical relevance of interview-assessed psychosis-risk symptoms in the young adult community. <i>Psychological Medicine</i> , <b>2018</b> , 48, 1167-1178	6.9	32
91	Striatal cerebral blood flow, executive functioning, and fronto-striatal functional connectivity in clinical high risk for psychosis. <i>Schizophrenia Research</i> , <b>2018</b> , 201, 231-236	3.6	13
90	Functioning mediates help-seeking for mental problems in the general population. <i>European Psychiatry</i> , <b>2018</b> , 54, 1-9	6	11
89	Resting-state gamma-band power alterations in schizophrenia reveal E/I-balance abnormalities across illness-stages. <i>ELife</i> , <b>2018</b> , 7,	8.9	50
88	Author response: Resting-state gamma-band power alterations in schizophrenia reveal E/I-balance abnormalities across illness-stages <b>2018</b> ,		2
87	T239. SINGLE-SUBJECT PREDICTION OF FUNCTIONAL OUTCOMES IN CLINICAL HIGH RISK SUBJECTS USING CLINICAL DATA. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, S209-S210	1.3	78
86	O11.8. PREVALENCE AND PREDICTORS OF INTERVIEW-ASSESSED CLINICAL HIGH-RISK SYMPTOMS IN THE GENERAL POPULATION. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, S109-S109	1.3	78
85	F26. THE NATURE OF CLINICAL HIGH-RISK SYMPTOMS: NEW INSIGHTS GAINED FROM AGE EFFECTS. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, S228-S229	1.3	78
84	Prediction Models of Functional Outcomes for Individuals in the Clinical High-Risk State for Psychosis or With Recent-Onset Depression: A Multimodal, Multisite Machine Learning Analysis. <i>JAMA Psychiatry</i> , <b>2018</b> , 75, 1156-1172	14.5	156
83	S211. SCHIZOTYPY IN PATIENTS FROM A CLINICAL HIGH RISK SERVICE: TRAIT OR STATE?. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, S407-S408	1.3	2
82	Psychopathology-a Precision Tool in Need of Re-sharpening. <i>Frontiers in Psychiatry</i> , <b>2018</b> , 9, 446	5	25

81	Enhancing Psychosis-Spectrum Nosology Through an International Data Sharing Initiative. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, S460-S467	1.3	9
80	Convergent and concurrent validity of the Frankfurt Complaint Questionnaire as a screener for psychosis risk. <i>Journal of Risk Research</i> , <b>2017</b> , 20, 1480-1496	4.2	9
79	The concept of basic symptoms: its scientific and clinical relevance. <i>World Psychiatry</i> , <b>2017</b> , 16, 104-105	14.4	27
78	The Italian version of the 92-item Prodromal Questionnaire: Concurrent validity with the SIPS and factor analysis in a sample of 258 outpatients aged 11-36years. <i>Schizophrenia Research</i> , <b>2017</b> , 189, 50-56	3.6	11
77	Association between mental health-related stigma and active help-seeking: systematic review and meta-analysis. <i>British Journal of Psychiatry</i> , <b>2017</b> , 210, 261-268	5.4	205
76	Age effect on prevalence of ultra-high risk for psychosis symptoms: replication in a clinical sample of an early detection of psychosis service. <i>European Child and Adolescent Psychiatry</i> , <b>2017</b> , 26, 1401-1405	5.5	21
75	No age effect in the prevalence and clinical significance of ultra-high risk symptoms and criteria for psychosis in 22q11 deletion syndrome: Confirmation of the genetically driven risk for psychosis?. <i>PLoS ONE</i> , <b>2017</b> , 12, e0174797	3.7	10
74	AuthorsSreply. <i>British Journal of Psychiatry</i> , <b>2017</b> , 211, 182-183	5.4	1
73	Mediators Linking Childhood Adversities and Trauma to Suicidality in Individuals at Risk for Psychosis. <i>Frontiers in Psychiatry</i> , <b>2017</b> , 8, 242	5	10
72	Psychosis-predictive value of self-reported schizotypy in a clinical high-risk sample. <i>Journal of Abnormal Psychology</i> , <b>2016</b> , 125, 923-932	7	46
71	The Dark Side of the Moon: Meta-analytical Impact of Recruitment Strategies on Risk Enrichment in the Clinical High Risk State for Psychosis. <i>Schizophrenia Bulletin</i> , <b>2016</b> , 42, 732-43	1.3	144
70	Stimulant Medication and Psychotic Symptoms in Offspring of Parents With Mental Illness. <i>Pediatrics</i> , <b>2016</b> , 137,	7.4	16
69	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , <b>2016</b> , 51, 247-57	4.5	17
68	Intensive community outreach for those at ultra high risk of psychosis: dilution, not solution. <i>Lancet Psychiatry</i> , <b>2016</b> , 3, 18	23.3	24
67	Attenuated psychotic and basic symptom characteristics in adolescents with ultra-high risk criteria for psychosis, other non-psychotic psychiatric disorders and early-onset psychosis. <i>European Child and Adolescent Psychiatry</i> , <b>2016</b> , 25, 1091-102	5.5	24
66	Revisiting the Basic Symptom Concept: Toward Translating Risk Symptoms for Psychosis into Neurobiological Targets. <i>Frontiers in Psychiatry</i> , <b>2016</b> , 7, 9	5	40
65	Psychometric properties of the Trauma and Distress Scale, TADS, in an adult community sample in Finland. <i>Högre Utbildning</i> , <b>2016</b> , 7, 30062	5	24
64	Abnormal involuntary movements are linked to psychosis-risk in children and adolescents: Results of a population-based study. <i>Schizophrenia Research</i> , <b>2016</b> , 174, 58-64	3.6	28

63	Ultra high risk status and transition to psychosis in 22q11.2 deletion syndrome. <i>World Psychiatry</i> , <b>2016</b> , 15, 259-265	14.4	41
62	Developing psychosis and its risk states through the lens of schizotypy. <i>Schizophrenia Bulletin</i> , <b>2015</b> , 41 Suppl 2, S396-407	1.3	120
61	Personality dimensions in persons symptomatically at risk of psychosis: pronounced but lacking a characteristic profile. <i>Microbial Biotechnology</i> , <b>2015</b> , 9, 242-7	3.3	6
60	Duration of unspecific prodromal and clinical high risk states, and early help-seeking in first-admission psychosis patients. <i>Social Psychiatry and Psychiatric Epidemiology</i> , <b>2015</b> , 50, 1831-41	4.5	41
59	Age matters in the prevalence and clinical significance of ultra-high-risk for psychosis symptoms and criteria in the general population: Findings from the BEAR and BEARS-kid studies. <i>World Psychiatry</i> , <b>2015</b> , 14, 189-97	14.4	80
58	At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. <i>World Psychiatry</i> , <b>2015</b> , 14, 322-32	14.4	172
57	Twelve-month psychosis-predictive value of the ultra-high risk criteria in children and adolescents. <i>Schizophrenia Research</i> , <b>2015</b> , 169, 186-192	3.6	35
56	Prediction and prevention of psychosis: current progress and future tasks. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2014</b> , 264 Suppl 1, S9-16	5.1	16
55	A stratified model for psychosis prediction in clinical practice. <i>Schizophrenia Bulletin</i> , <b>2014</b> , 40, 1533-42	1.3	53
54	Prevalence and clinical significance of DSM-5-attenuated psychosis syndrome in adolescents and young adults in the general population: the Bern Epidemiological At-Risk (BEAR) study. <i>Schizophrenia Bulletin</i> , <b>2014</b> , 40, 1499-508	1.3	70
53	Improving the clinical prediction of psychosis by combining ultra-high risk criteria and cognitive basic symptoms. <i>Schizophrenia Research</i> , <b>2014</b> , 154, 100-6	3.6	94
52	Differences in coping, self-efficacy, and external control beliefs between patients at-risk for psychosis and patients with first-episode psychosis. <i>Psychiatry Research</i> , <b>2014</b> , 219, 95-102	9.9	33
51	Early Detection and Treatment of Psychosis: The Bern Child and Adolescent Psychiatric Perspective. <i>Advances in Psychiatry</i> , <b>2014</b> , 2014, 1-16		5
50	Self-reported psychotic-like experiences are a poor estimate of clinician-rated attenuated and frank delusions and hallucinations. <i>Psychopathology</i> , <b>2014</b> , 47, 194-201	3.4	47
49	Self-reported attenuated psychotic-like experiences in help-seeking adolescents and their association with age, functioning and psychopathology. <i>Schizophrenia Research</i> , <b>2014</b> , 160, 110-7	3.6	37
48	Can quantitative EEG measures predict clinical outcome in subjects at Clinical High Risk for psychosis? A prospective multicenter study. <i>Schizophrenia Research</i> , <b>2014</b> , 153, 42-7	3.6	38
47	Reliability of telephone assessments of at-risk criteria of psychosis: a comparison to face-to-face interviews. <i>Schizophrenia Research</i> , <b>2014</b> , 153, 251-3	3.6	15
46	Pathways to care in subjects at high risk for psychotic disorders - a European perspective. <i>Schizophrenia Research</i> , <b>2014</b> , 152, 400-7	3.6	21



45	Psychosocial outcome in patients at clinical high risk of psychosis: a prospective follow-up. <i>Social Psychiatry and Psychiatric Epidemiology</i> , <b>2013</b> , 48, 303-11	4.5	27
44	The Schizophrenia Proneness Instrument, Child and Youth version (SPI-CY): practicability and discriminative validity. <i>Schizophrenia Research</i> , <b>2013</b> , 146, 69-78	3.6	65
43	The psychosis high-risk state: a comprehensive state-of-the-art review. <i>JAMA Psychiatry</i> , <b>2013</b> , 70, 107-20	4.5	965
42	A rose is a rose is a rose, but at-risk criteria differ. <i>Psychopathology</i> , <b>2013</b> , 46, 75-87	3.4	51
41	The significance of at-risk symptoms for psychosis in children and adolescents. <i>Canadian Journal of Psychiatry</i> , <b>2013</b> , 58, 32-40	4.8	61
40	Personality disorders and accentuations in at-risk persons with and without conversion to first-episode psychosis. <i>Microbial Biotechnology</i> , <b>2012</b> , 6, 389-98	3.3	34
39	Pharmacological prevention and treatment in clinical at-risk states for psychosis. <i>Current Pharmaceutical Design</i> , <b>2012</b> , 18, 550-7	3.3	16
38	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , <b>2012</b> , 38, 1130-4	1.3	74
37	Axis I diagnoses and transition to psychosis in clinical high-risk patients EPOS project: prospective follow-up of 245 clinical high-risk outpatients in four countries. <i>Schizophrenia Research</i> , <b>2012</b> , 138, 192-7	3.6	85
36	Comparing the prodrome of schizophrenia-spectrum psychoses and affective disorders with and without psychotic features. <i>Schizophrenia Research</i> , <b>2012</b> , 138, 218-22	3.6	13
35	Early detection and intervention of psychosis in children and adolescents: urgent need for studies. <i>European Child and Adolescent Psychiatry</i> , <b>2012</b> , 21, 239-41	5.5	45
34	Basic symptoms and the prediction of first-episode psychosis. <i>Current Pharmaceutical Design</i> , <b>2012</b> , 18, 351-7	3.3	121
33	Prediction of psychosis by mismatch negativity. <i>Biological Psychiatry</i> , <b>2011</b> , 69, 959-66	7.9	219
32	What percentage of people in the general population satisfies the current clinical at-risk criteria of psychosis?. <i>Schizophrenia Research</i> , <b>2011</b> , 125, 99-100	3.6	41
31	Cannabis use disorder and age at onset of psychosis--a study in first-episode patients. <i>Schizophrenia Research</i> , <b>2011</b> , 129, 52-6	3.6	40
30	The near Babylonian speech confusion in early detection of psychosis. <i>Schizophrenia Bulletin</i> , <b>2011</b> , 37, 653-5	1.3	54
29	Prediction and prevention of schizophrenia: what has been achieved and where to go next?. <i>World Psychiatry</i> , <b>2011</b> , 10, 165-74	14.4	73
28	Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. <i>Schizophrenia Bulletin</i> , <b>2010</b> , 36, 182-91	1.3	152

27	Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. <i>Archives of General Psychiatry</i> , <b>2010</b> , 67, 241-51		491
26	Probably at-risk, but certainly ill—advocating the introduction of a psychosis spectrum disorder in DSM-V. <i>Schizophrenia Research</i> , <b>2010</b> , 120, 23-37	3.6	114
25	Disability in people clinically at high risk of psychosis. <i>British Journal of Psychiatry</i> , <b>2010</b> , 197, 278-84	5.4	101
24	Anandamide elevation in cerebrospinal fluid in initial prodromal states of psychosis. <i>British Journal of Psychiatry</i> , <b>2009</b> , 194, 371-2	5.4	140
23	Binocular depth inversion as a paradigm of reduced visual information processing in prodromal state, antipsychotic-naïve and treated schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2009</b> , 259, 195-202	5.1	74
22	Subjective symptoms of schizophrenia in research and the clinic: the basic symptom concept. <i>Schizophrenia Bulletin</i> , <b>2009</b> , 35, 5-8	1.3	221
21	Early detection of psychosis - establishing a service for persons at risk. <i>European Psychiatry</i> , <b>2009</b> , 24, 1-10	6	35
20	Intervention in the at-risk state to prevent transition to psychosis. <i>Current Opinion in Psychiatry</i> , <b>2009</b> , 22, 177-83	4.9	22
19	Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. <i>Biological Psychiatry</i> , <b>2008</b> , 64, 376-84	7.9	181
18	Kraepelin and psychotic prodromal conditions. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2008</b> , 258 Suppl 2, 74-84	5.1	37
17	Neurocognitive indicators for a conversion to psychosis: comparison of patients in a potentially initial prodromal state who did or did not convert to a psychosis. <i>Schizophrenia Research</i> , <b>2007</b> , 92, 116-25	2.6	161
16	Orienting of attention in unmedicated patients with schizophrenia, prodromal subjects and healthy relatives. <i>Schizophrenia Research</i> , <b>2007</b> , 97, 35-42	3.6	27
15	Relationship between subjective and objective cognitive function in the early and late prodrome. <i>British Journal of Psychiatry</i> , <b>2007</b> , 51, s43-51	5.4	36
14	Basic symptoms in early psychotic and depressive disorders. <i>British Journal of Psychiatry</i> , <b>2007</b> , 51, s31-7	5.4	59
13	The initial prodrome of schizophrenia: different duration, different underlying deficits?. <i>Comprehensive Psychiatry</i> , <b>2007</b> , 48, 479-88	7.3	31
12	CSF metabolic and proteomic profiles in patients prodromal for psychosis. <i>PLoS ONE</i> , <b>2007</b> , 2, e756	3.7	80
11	Neurocognitive functioning in subjects at risk for a first episode of psychosis compared with first- and multiple-episode schizophrenia. <i>Journal of Clinical and Experimental Neuropsychology</i> , <b>2006</b> , 28, 1388-407	2.1	108
10	Disturbances of visual information processing in early states of psychosis and experimental delta-9-tetrahydrocannabinol altered states of consciousness. <i>Schizophrenia Research</i> , <b>2006</b> , 88, 142-50	3.6	79

9	Impaired mismatch negativity generation in prodromal subjects and patients with schizophrenia. <i>Schizophrenia Research</i> , <b>2005</b> , 73, 297-310	3.6	130
8	Subjective quality of life in subjects at risk for a first episode of psychosis: a comparison with first episode schizophrenia patients and healthy controls. <i>Schizophrenia Research</i> , <b>2005</b> , 79, 137-43	3.6	81
7	The European Prediction of Psychosis Study (EPOS): integrating early recognition and intervention in Europe. <i>World Psychiatry</i> , <b>2005</b> , 4, 161-7	14.4	97
6	Modeling the early course of schizophrenia. <i>Schizophrenia Bulletin</i> , <b>2003</b> , 29, 325-40	1.3	119
5	Self-experienced vulnerability, prodromal symptoms and coping strategies preceding schizophrenic and depressive relapses. <i>European Psychiatry</i> , <b>2002</b> , 17, 384-93	6	34
4	Diagnosing schizophrenia in the initial prodromal phase. <i>Archives of General Psychiatry</i> , <b>2001</b> , 58, 158-64		785
3	Facet theoretical analysis of the similarity relationships of personality factors and personality disorders. <i>European Psychiatry</i> , <b>1998</b> , 13, 193s-193s	6	
2	Can self-experienced neuropsychological deficits indicate propensity to schizophrenic psychosis? Results of an 8-year prospective follow-up study. <i>International Clinical Psychopharmacology</i> , <b>1998</b> , 13, S75-S80	2.2	2
1	Diagnostic validity of basic symptoms. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>1996</b> , 246, 147-54	5.1	101