

# Silvia Parolini

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8123549/publications.pdf>

Version: 2024-02-01

64  
papers

6,423  
citations

109264

35  
h-index

114418

63  
g-index

65  
all docs

65  
docs citations

65  
times ranked

7332  
citing authors

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Identification and Molecular Characterization of Nkp30, a Novel Triggering Receptor Involved in Natural Cytotoxicity Mediated by Human Natural Killer Cells. <i>Journal of Experimental Medicine</i> , 1999, 190, 1505-1516.                                   | 4.2  | 664       |
| 2  | X-Linked Lymphoproliferative Disease. <i>Journal of Experimental Medicine</i> , 2000, 192, 337-346.  | 4.2  | 438       |
| 3  | NKp44, A Triggering Receptor Involved in Tumor Cell Lysis by Activated Human Natural Killer Cells, Is a Novel Member of the Immunoglobulin Superfamily. <i>Journal of Experimental Medicine</i> , 1999, 189, 787-796.  | 4.2  | 396       |
| 4  | Identification of a subset of human natural killer cells expressing high levels of programmed death 1: phenotypic and functional characterization. <i>Journal of Allergy and Clinical Immunology</i> , 2017, 139, 335-346.e3.                                  | 1.5  | 379       |
| 5  | The role of chemerin in the colocalization of NK and dendritic cell subsets into inflamed tissues. <i>Blood</i> , 2007, 109, 3625-3632.  | 0.6  | 336       |
| 6  | NK cells at the interface between innate and adaptive immunity. <i>Cell Death and Differentiation</i> , 2008, 15, 226-233.   | 5.0  | 291       |
| 7  | Gntb-A, a Novel Sh2d1a-Associated Surface Molecule Contributing to the Inability of Natural Killer Cells to Kill Epstein-Barr Virus-Infected B Cells in X-Linked Lymphoproliferative Disease. <i>Journal of Experimental Medicine</i> , 2001, 194, 235-246.    | 4.2  | 287       |
| 8  | Human natural killer cell receptors and co-receptors. <i>Immunological Reviews</i> , 2001, 181, 203-214.   | 2.8  | 273       |
| 9  | 2B4 functions as a co-receptor in human NK cell activation. <i>European Journal of Immunology</i> , 2000, 30, 787-793.   | 1.6  | 202       |
| 10 | Identification of NKp80, a novel triggering molecule expressed by human NK cells. <i>European Journal of Immunology</i> , 2001, 31, 233-242.   | 1.6  | 185       |
| 11 | Early expression of triggering receptors and regulatory role of 2B4 in human natural killer cell precursors undergoing in vitro differentiation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 4526-4531. | 3.3  | 174       |
| 12 | Inherited DOCK2 Deficiency in Patients with Early-Onset Invasive Infections. <i>New England Journal of Medicine</i> , 2015, 372, 2409-2422.  | 13.9 | 169       |
| 13 | IL-21 induces both rapid maturation of human CD34+ cell precursors towards NK cells and acquisition of surface killer Ig-like receptors. <i>European Journal of Immunology</i> , 2003, 33, 3439-3447.  | 1.6  | 166       |
| 14 | A novel primary human immunodeficiency due to deficiency in the WASP-interacting protein WIP. <i>Journal of Experimental Medicine</i> , 2012, 209, 29-34.  | 4.2  | 158       |
| 15 | B7-H6-mediated downregulation of NKp30 in NK cells contributes to ovarian carcinoma immune escape. <i>Oncology</i> , 2015, 4, e1001224.  | 2.1  | 137       |
| 16 | Innate immunity defects in Hermansky-Pudlak type 2 syndrome. <i>Blood</i> , 2006, 107, 4857-4864.  | 0.6  | 136       |
| 17 | CD94 functions as a natural killer cell inhibitory receptor for different HLA class I alleles: identification of the inhibitory form of CD94 by the use of novel monoclonal antibodies. <i>European Journal of Immunology</i> , 1996, 26, 2487-2492.           | 1.6  | 130       |
| 18 | Impaired natural and CD16-mediated NK cell cytotoxicity in patients with WAS and XLT: ability of IL-2 to correct NK cell functional defect. <i>Blood</i> , 2004, 104, 436-443.   | 0.6  | 130       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | IL-12 or IL-4 Prime Human NK Cells to Mediate Functionally Divergent Interactions with Dendritic Cells or Tumors. <i>Journal of Immunology</i> , 2005, 174, 3992-3998.  | 0.4 | 117       |
| 20 | Basic Fibroblast Growth Factor-Induced Angiogenic Phenotype in Mouse Endothelium. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1997, 17, 454-464.  | 1.1 | 108       |
| 21 | Involvement of natural cytotoxicity receptors in human natural killer cell-mediated lysis of neuroblastoma and glioblastoma cell lines. <i>Journal of Neuroimmunology</i> , 2000, 107, 220-225.   | 1.1 | 103       |
| 22 | The leukocyte Ig-like receptor (LIR)-1 for the cytomegalovirus UL18 protein displays a broad specificity for different HLA class I alleles: analysis of LIR-1+ NK cell clones. <i>International Immunology</i> , 1999, 11, 29-35.   | 1.8 | 98        |
| 23 | Severe impairment of IFN- $\gamma$ and IFN- $\alpha$ responses in cells of a patient with a novel STAT1 splicing mutation. <i>Blood</i> , 2011, 118, 1806-1817.   | 0.6 | 84        |
| 24 | Selective cross-talk among natural cytotoxicity receptors in human natural killer cells. <i>European Journal of Immunology</i> , 2003, 33, 1235-1241.   | 1.6 | 77        |
| 25 | CD59 is physically and functionally associated with natural cytotoxicity receptors and activates human NK cell-mediated cytotoxicity. <i>European Journal of Immunology</i> , 2003, 33, 3367-3376.  | 1.6 | 77        |
| 26 | Exome sequencing reveals a pallidin mutation in a Hermansky-Pudlak-like primary immunodeficiency syndrome. <i>Blood</i> , 2012, 119, 3185-3187.   | 0.6 | 76        |
| 27 | Triggering receptors involved in natural killer cell-mediated cytotoxicity against choriocarcinoma cell lines. <i>Human Immunology</i> , 2000, 61, 1055-1058.   | 1.2 | 71        |
| 28 | Clinical, laboratory and molecular signs of immunodeficiency in patients with partial oculo-cutaneous albinism. <i>Orphanet Journal of Rare Diseases</i> , 2013, 8, 168.  | 1.2 | 70        |
| 29 | Defective natural killer cell cytotoxic activity in NFKB2-mutated CVID-like disease. <i>Journal of Allergy and Clinical Immunology</i> , 2015, 135, 1641-1643.e3.   | 1.5 | 68        |
| 30 | Reduced thymic output, increased spontaneous apoptosis and oligoclonal B cells in polyethylene glycol-adenosine deaminase-treated patients. <i>European Journal of Immunology</i> , 2005, 35, 3376-3386.  | 1.6 | 59        |
| 31 | Impaired natural killer cell functions in patients with signal transducer and activator of transcription 1 (STAT1) gain-of-function mutations. <i>Journal of Allergy and Clinical Immunology</i> , 2017, 140, 553-564.e4.   | 1.5 | 58        |
| 32 | Novel insights from adaptor protein 3 complex deficiency. <i>Journal of Allergy and Clinical Immunology</i> , 2007, 120, 735-741.   | 1.5 | 51        |
| 33 | Killer cell immunoglobulin-like receptor expression delineates in situ Sezary syndrome lymphocytes. <i>Journal of Pathology</i> , 2003, 199, 77-83.   | 2.1 | 47        |
| 34 | Linker for Activation of T Cells (LAT), a Novel Immunohistochemical Marker for T Cells, NK Cells, Mast Cells, and Megakaryocytes. <i>American Journal of Pathology</i> , 1999, 154, 1037-1046.  | 1.9 | 46        |
| 35 | Natural Killer Cells from Patients with Recombinase-Activating Gene and Non-Homologous End Joining Gene Defects Comprise a Higher Frequency of CD56bright NKG2A+++ Cells, and Yet Display Increased Degranulation and Higher Perforin Content. <i>Frontiers in Immunology</i> , 2017, 8, 798. | 2.2 | 41        |
| 36 | NFKB1 regulates human NK cell maturation and effector functions. <i>Clinical Immunology</i> , 2017, 175, 99-108.  | 1.4 | 38        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | A monoallelic activating mutation in RAC2 resulting in a combined immunodeficiency. <i>Journal of Allergy and Clinical Immunology</i> , 2019, 143, 1649-1653.e3.   | 1.5 | 37        |
| 38 | Strengthening the AntiTumor NK Cell Function for the Treatment of Ovarian Cancer. <i>International Journal of Molecular Sciences</i> , 2019, 20, 890.  | 1.8 | 34        |
| 39 | Occurrence of Nodular Lymphocyte-Predominant Hodgkin Lymphoma in Hermansky-Pudlak Type 2 Syndrome Is Associated to Natural Killer and Natural Killer T Cell Defects. <i>PLoS ONE</i> , 2013, 8, e80131.        | 1.1 | 34        |
| 40 | GPR56 as a novel marker identifying the CD56dull CD16+ NK cell subset both in blood stream and in inflamed peripheral tissues. <i>International Immunology</i> , 2010, 22, 91-100.                             | 1.8 | 33        |
| 41 | CTLA-4 regulates human Natural Killer cell effector functions. <i>Clinical Immunology</i> , 2018, 194, 43-45.  | 1.4 | 30        |
| 42 | Functional characterization of natural killer cells in type I leukocyte adhesion deficiency. <i>Blood</i> , 2007, 109, 4873-4881.  | 0.6 | 29        |
| 43 | Activin A as a Mediator of NK-Dendritic Cell Functional Interactions. <i>Journal of Immunology</i> , 2014, 192, 1241-1248.   | 0.4 | 27        |
| 44 | Distinctive Lack of CD48 Expression in Subsets of Human Dendritic Cells Tunes NK Cell Activation. <i>Journal of Immunology</i> , 2005, 175, 3690-3697.   | 0.4 | 26        |
| 45 | Pseudorabies Virus US3 Protein Kinase Protects Infected Cells from NK Cell-Mediated Lysis via Increased Binding of the Inhibitory NK Cell Receptor CD300a. <i>Journal of Virology</i> , 2016, 90, 1522-1533.   | 1.5 | 26        |
| 46 | NK cells and their receptors during viral infections. <i>Immunotherapy</i> , 2011, 3, 1075-1086.   | 1.0 | 25        |
| 47 | Effects of opioid therapy on human natural killer cells. <i>International Immunopharmacology</i> , 2014, 18, 169-174.  | 1.7 | 24        |
| 48 | In vitro treatment with concentrated growth factors (CGF) and sodium orthosilicate positively affects cell renewal in three different human cell lines. <i>Cell Biology International</i> , 2018, 42, 353-364. | 1.4 | 22        |
| 49 | XLP1 inhibitory effect by B220 does not affect DNAM1 and NKG2D activating pathways in NK cells. <i>European Journal of Immunology</i> , 2014, 44, 1526-1534.   | 1.6 | 20        |
| 50 | Up-regulation of urokinase-type plasminogen activator in squamous cell carcinoma of human larynx. <i>British Journal of Cancer</i> , 1996, 74, 1168-1174.  | 2.9 | 18        |
| 51 | Diagnosing XLP1 in patients with hemophagocytic lymphohistiocytosis. <i>Journal of Allergy and Clinical Immunology</i> , 2014, 134, 1381-1387.e7.  | 1.5 | 14        |
| 52 | Primitive Neuroectodermal Tumor in an Ovarian Cystic Teratoma: Natural Killer and Neuroblastoma Cell Analysis. <i>Case Reports in Oncology</i> , 2014, 7, 70-78.   | 0.3 | 12        |
| 53 | Natural killer cell hyporesponsiveness and impaired development in a CD247-deficient patient. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 137, 942-945.e4.                                       | 1.5 | 12        |
| 54 | The RAC2-PI3K axis regulates human NK cell maturation and function. <i>Clinical Immunology</i> , 2019, 208, 108257.  | 1.4 | 11        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | FUNCTION AND SPECIFICITY OF HUMAN NATURAL KILLER CELL RECEPTORS. <i>International Journal of Immunogenetics</i> , 1997, 24, 455-468.   | 1.2 | 7         |
| 56 | Cellular and molecular pathogenesis of X-linked lymphoproliferative disease. <i>Current Opinion in Allergy and Clinical Immunology</i> , 2001, 1, 513-517.   | 1.1 | 7         |
| 57 | p85 $\hat{1}$ is an intrinsic regulator of human natural killer cell effector functions. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 138, 605-608.e3.  | 1.5 | 7         |
| 58 | Combined immunodeficiency with autoimmunity caused by a homozygous missense mutation in inhibitor of nuclear factor $\gamma$ B kinase alpha (IKK $\hat{1}$ ). <i>Science Immunology</i> , 2021, 6, eabf6723. | 5.6 | 6         |
| 59 | Natural killer cell impairment in ovarian clear cell carcinoma. <i>Journal of Leukocyte Biology</i> , 2020, 108, 1425-1434.  | 1.5 | 3         |
| 60 | X-linked lymphoproliferative disease: the dark side of 2b4 function. <i>Advances in Experimental Medicine and Biology</i> , 2001, 495, 63-67.  | 0.8 | 3         |
| 61 | From Natural Killer Cell Receptor Discovery to Characterization of Natural Killer Cell Defects in Primary Immunodeficiencies. <i>Frontiers in Immunology</i> , 2019, 10, 1757.                               | 2.2 | 2         |
| 62 | Lack of DOCK8 impairs the primary biologic functions of human NK cells and abrogates CCR7 surface expression in a WASP-independent manner. <i>Clinical Immunology</i> , 2022, 237, 108974.                   | 1.4 | 2         |
| 63 | Response to the Letter to the Editor Regarding "Functional evaluation of natural killer cell cytotoxic activity in NFKB-2 mutated patients". <i>Immunology Letters</i> , 2018, 200, 16-17.                   | 1.1 | 0         |
| 64 | A novel primary human immunodeficiency due to deficiency in the WASP-interacting protein WIP. <i>Journal of Cell Biology</i> , 2012, 196, i1-i1.   | 2.3 | 0         |