

Federica Genovese

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8130909/publications.pdf>

Version: 2024-02-01

23
papers

917
citations

516215

16
h-index

642321

23
g-index

23
all docs

23
docs citations

23
times ranked

1260
citing authors

#	ARTICLE	IF	CITATIONS
1	The extracellular matrix in the kidney: a source of novel non-invasive biomarkers of kidney fibrosis?. <i>Fibrogenesis and Tissue Repair</i> , 2014, 7, 4.	3.4	266
2	Suitability Of Nitisinone In Alkaptonuria 1 (SONIA 1): an international, multicentre, randomised, open-label, no-treatment controlled, parallel-group, dose-response study to investigate the effect of once daily nitisinone on 24-h urinary homogentisic acid excretion in patients with alkaptonuria after 4â€¦weeks of treatment. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 362-367.	0.5	123
3	Serum endotrophin, a type VI collagen cleavage product, is associated with increased mortality in chronic kidney disease. <i>PLoS ONE</i> , 2017, 12, e0175200.	1.1	56
4	Serum and urine markers of collagen degradation reflect renal fibrosis in experimental kidney diseases. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 1112-1121.	0.4	53
5	Urinary endotrophin predicts disease progression in patients with chronic kidney disease. <i>Scientific Reports</i> , 2017, 7, 17328.	1.6	52
6	Higher Collagen VI Formation Is Associated With All-Cause Mortality in Patients With Type 2 Diabetes and Microalbuminuria. <i>Diabetes Care</i> , 2018, 41, 1493-1500.	4.3	51
7	Turnover of type III collagen reflects disease severity and is associated with progression and microinflammation in patients with IgA nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, 472-479.	0.4	45
8	Markers of Collagen Formation and Degradation Reflect Renal Function and Predict Adverse Outcomes in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2019, 42, 1760-1768.	4.3	30
9	Protein degradation fragments as diagnostic and prognostic biomarkers of connective tissue diseases: understanding the extracellular matrix message and implication for current and future serological biomarkers. <i>Expert Review of Proteomics</i> , 2016, 13, 213-225.	1.3	28
10	Non-invasive quantification of collagen turnover in renal transplant recipients. <i>PLoS ONE</i> , 2017, 12, e0175898.	1.1	27
11	Collagen turnover profiles in chronic kidney disease. <i>Scientific Reports</i> , 2019, 9, 16062.	1.6	27
12	Atrial fibrillation and cardiac fibrosis: A review on the potential of extracellular matrix proteins as biomarkers. <i>Matrix Biology</i> , 2020, 91-92, 188-203.	1.5	20
13	Accelerated collagen turnover in women with angina pectoris without obstructive coronary artery disease: An iPOWER substudy. <i>European Journal of Preventive Cardiology</i> , 2018, 25, 719-727.	0.8	19
14	Clinical evaluation of a matrix metalloproteinase-12 cleaved fragment of titin as a cardiovascular serological biomarker. <i>Journal of Translational Medicine</i> , 2012, 10, 140.	1.8	18
15	Homogentisic acid is not only eliminated by glomerular filtration and tubular secretion but also produced in the kidney in alkaptonuria. <i>Journal of Inherited Metabolic Disease</i> , 2020, 43, 737-747.	1.7	18
16	Imbalanced turnover of collagen type III is associated with disease progression and mortality in high-risk chronic kidney disease patients. <i>CKJ: Clinical Kidney Journal</i> , 2021, 14, 593-601.	1.4	18
17	<p>Cross-Linked Multimeric Pro-Peptides of Type III Collagen (PC3X) in Hepatocellular Carcinoma â€“ A Biomarker That Provides Additional Prognostic Value in AFP Positive Patients<p>. <i>Journal of Hepatocellular Carcinoma</i> , 2020, Volume 7, 301-313.	1.8	17
18	Subclinical ochronosis features in alkaptonuria: a cross-sectional study. <i>BMJ Innovations</i> , 2019, 5, 82-91.	1.0	15

#	ARTICLE	IF	CITATIONS
19	Collagen Type III and VI Remodeling Biomarkers Are Associated with Kidney Fibrosis in Lupus Nephritis. <i>Kidney360</i> , 2021, 2, 1473-1481.	0.9	12
20	The proteoglycan mimecan is associated with carotid plaque vulnerability and increased risk of future cardiovascular death. <i>Atherosclerosis</i> , 2020, 313, 88-95.	0.4	10
21	Evaluation of a novel biomarker of type XXVIII collagen formation, PRO-C28, in samples from cancer and heart failure with preserved ejection fraction patients. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 204, 114272.	1.4	6
22	Circulating Levels of Endotrophin Are Prognostic for Long-Term Mortality after AKI. <i>Kidney360</i> , 2022, 3, 809-817.	0.9	4
23	Increased proteolytic cleavage of osteoglycin is associated with a stable plaque phenotype and lower risk of cardiovascular events. <i>Atherosclerosis</i> , 2022, 355, 8-14.	0.4	2