

Federica Genovese

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8130909/federica-genovese-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

21
papers

597
citations

12
h-index

23
g-index

23
ext. papers

755
ext. citations

4.9
avg, IF

3.68
L-index

#	Paper	IF	Citations
21	The extracellular matrix in the kidney: a source of novel non-invasive biomarkers of kidney fibrosis?. <i>Fibrogenesis and Tissue Repair</i> , 2014 , 7, 4		200
20	Suitability Of Nitisinone In Alkaptonuria 1 (SONIA 1): an international, multicentre, randomised, open-label, no-treatment controlled, parallel-group, dose-response study to investigate the effect of once daily nitisinone on 24-h urinary homogentisic acid excretion in patients with alkaptonuria after 4 weeks of treatment. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 362-7	2.4	92
19	Serum and urine markers of collagen degradation reflect renal fibrosis in experimental kidney diseases. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, 1112-21	4.3	45
18	Turnover of type III collagen reflects disease severity and is associated with progression and microinflammation in patients with IgA nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2016 , 31, 472-9	4.3	38
17	Serum endotrophin, a type VI collagen cleavage product, is associated with increased mortality in chronic kidney disease. <i>PLoS ONE</i> , 2017 , 12, e0175200	3.7	31
16	Higher Collagen VI Formation Is Associated With All-Cause Mortality in Patients With Type 2 Diabetes and Microalbuminuria. <i>Diabetes Care</i> , 2018 , 41, 1493-1500	14.6	30
15	Urinary endotrophin predicts disease progression in patients with chronic kidney disease. <i>Scientific Reports</i> , 2017 , 7, 17328	4.9	27
14	Non-invasive quantification of collagen turnover in renal transplant recipients. <i>PLoS ONE</i> , 2017 , 12, e0175898	3.7	18
13	Protein degradation fragments as diagnostic and prognostic biomarkers of connective tissue diseases: understanding the extracellular matrix message and implication for current and future serological biomarkers. <i>Expert Review of Proteomics</i> , 2016 , 13, 213-25	4.2	18
12	Markers of Collagen Formation and Degradation Reflect Renal Function and Predict Adverse Outcomes in Patients With Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 1760-1768	14.6	15
11	Clinical evaluation of a matrix metalloproteinase-12 cleaved fragment of titin as a cardiovascular serological biomarker. <i>Journal of Translational Medicine</i> , 2012 , 10, 140	8.5	15
10	Collagen turnover profiles in chronic kidney disease. <i>Scientific Reports</i> , 2019 , 9, 16062	4.9	14
9	Accelerated collagen turnover in women with angina pectoris without obstructive coronary artery disease: An iPOWER substudy. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 719-727	3.9	10
8	Atrial fibrillation and cardiac fibrosis: A review on the potential of extracellular matrix proteins as biomarkers. <i>Matrix Biology</i> , 2020 , 91-92, 188-203	11.4	9
7	Subclinical ochronosis features in alkaptonuria: a cross-sectional study. <i>BMJ Innovations</i> , 2019 , 5, 82-91	1.8	8
6	Cross-Linked Multimeric Pro-Peptides of Type III Collagen (PC3X) in Hepatocellular Carcinoma - A Biomarker That Provides Additional Prognostic Value in AFP Positive Patients. <i>Journal of Hepatocellular Carcinoma</i> , 2020 , 7, 301-313	5.3	8
5	Imbalanced turnover of collagen type III is associated with disease progression and mortality in high-risk chronic kidney disease patients. <i>CKJ: Clinical Kidney Journal</i> , 2021 , 14, 593-601	4.5	8

4	Homogentisic acid is not only eliminated by glomerular filtration and tubular secretion but also produced in the kidney in alkaptonuria. <i>Journal of Inherited Metabolic Disease</i> , 2020 , 43, 737-747	5.4	7
3	The proteoglycan mimecan is associated with carotid plaque vulnerability and increased risk of future cardiovascular death. <i>Atherosclerosis</i> , 2020 , 313, 88-95	3.1	2
2	Evaluation of a novel biomarker of type XXVIII collagen formation, PRO-C28, in samples from cancer and heart failure with preserved ejection fraction patients. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021 , 204, 114272	3.5	2
1	Collagen Type III and VI Remodeling Biomarkers Are Associated with Kidney Fibrosis in Lupus Nephritis.. <i>Kidney360</i> , 2021 , 2, 1473-1481	1.8	0