

# Joanna Mika

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8132137/publications.pdf>

Version: 2024-02-01

122  
papers

5,604  
citations

66315

42  
h-index

88593

70  
g-index

123  
all docs

123  
docs citations

123  
times ranked

5814  
citing authors

#	ARTICLE	IF	CITATIONS
1	Mirogabalin Decreases Pain-like Behaviours and Improves Opioid and Ketamine Antinociception in a Mouse Model of Neuropathic Pain. <i>Pharmaceuticals</i> , 2022, 15, 88.	1.7	3
2	Peripheral Mechanisms of Neuropathic Pain—The Role of Neuronal and Non-Neuronal Interactions and Their Implications for Topical Treatment of Neuropathic Pain. <i>Pharmaceuticals</i> , 2021, 14, 77.	1.7	26
3	Mirogabalin—A Novel Selective Ligand for the $\alpha_2\delta_1$ Calcium Channel Subunit. <i>Pharmaceuticals</i> , 2021, 14, 112.	1.7	20
4	Topical Treatments and Their Molecular/Cellular Mechanisms in Patients with Peripheral Neuropathic Pain—Narrative Review. <i>Pharmaceutics</i> , 2021, 13, 450.	2.0	27
5	Long pentraxin PTX3 is upregulated systemically and centrally after experimental neurotrauma, but its depletion leaves unaltered sensorimotor deficits or histopathology. <i>Scientific Reports</i> , 2021, 11, 9616.	1.6	12
6	Nitric oxide modulates tapentadol antinociceptive tolerance and physical dependence. <i>European Journal of Pharmacology</i> , 2021, 907, 174245.	1.7	5
7	Novel bifunctional hybrid compounds designed to enhance the effects of opioids and antagonize the pronociceptive effects of nonopioid peptides as potent analgesics in a rat model of neuropathic pain. <i>Pain</i> , 2021, 162, 432-445.	2.0	9
8	Initiators of Classical and Lectin Complement Pathways Are Differently Engaged after Traumatic Brain Injury—Time-Dependent Changes in the Cortex, Striatum, Thalamus and Hippocampus in a Mouse Model. <i>International Journal of Molecular Sciences</i> , 2021, 22, 45.	1.8	8
9	The Kynurenine Pathway as a Potential Target for Neuropathic Pain Therapy Design: From Basic Research to Clinical Perspectives. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11055.	1.8	22
10	Comparison of the Effects of Chemokine Receptors CXCR2 and CXCR3 Pharmacological Modulation in Neuropathic Pain Model—In Vivo and In Vitro Study. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11074.	1.8	13
11	Blockade of CC Chemokine Receptor Type 3 Diminishes Pain and Enhances Opioid Analgesic Potency in a Model of Neuropathic Pain. <i>Frontiers in Immunology</i> , 2021, 12, 781310.	2.2	15
12	The blockade of CC chemokine receptor type 1 influences the level of nociceptive factors and enhances opioid analgesic potency in a rat model of neuropathic pain. <i>Immunology</i> , 2020, 159, 413-428.	2.0	28
13	CCR4 Antagonist (C021) Administration Diminishes Hypersensitivity and Enhances the Analgesic Potency of Morphine and Buprenorphine in a Mouse Model of Neuropathic Pain. <i>Frontiers in Immunology</i> , 2020, 11, 1241.	2.2	16
14	Metamizole relieves pain by influencing cytokine levels in dorsal root ganglia in a rat model of neuropathic pain. <i>Pharmacological Reports</i> , 2020, 72, 1310-1322.	1.5	8
15	Traumatic brain injury in mice induces changes in the expression of the XCL1/XCR1 and XCL1/ITGA9 axes. <i>Pharmacological Reports</i> , 2020, 72, 1579-1592.	1.5	7
16	Novel hybrid compounds, opioid agonist+melanocortin 4 receptor antagonist, as efficient analgesics in mouse chronic constriction injury model of neuropathic pain. <i>Neuropharmacology</i> , 2020, 178, 108232.	2.0	14
17	Changes in macrophage inflammatory protein-1 (MIP-1) family members expression induced by traumatic brain injury in mice. <i>Immunobiology</i> , 2020, 225, 151911.	0.8	22
18	Bidirectional Action of Cenriciviroc, a CCR2/CCR5 Antagonist, Results in Alleviation of Pain-Related Behaviors and Potentiation of Opioid Analgesia in Rats With Peripheral Neuropathy. <i>Frontiers in Immunology</i> , 2020, 11, 615327.	2.2	17

#	ARTICLE	IF	CITATIONS
19	The CCL2/CCL7/CCL12/CCR2 pathway is substantially and persistently upregulated in mice after traumatic brain injury, and CCL2 modulates the complement system in microglia. <i>Molecular and Cellular Probes</i> , 2020, 54, 101671.	0.9	26
20	CCR4 antagonist (C021) influences the level of nociceptive factors and enhances the analgesic potency of morphine in a rat model of neuropathic pain. <i>European Journal of Pharmacology</i> , 2020, 880, 173166.	1.7	16
21	Comparison of the beneficial effects of RS504393, maraviroc and cenicriviroc on neuropathic pain-related symptoms in rodents: behavioral and biochemical analyses. <i>International Immunopharmacology</i> , 2020, 84, 106540.	1.7	16
22	Blockade of CCR4 Diminishes Hypersensitivity and Enhances Opioid Analgesia – Evidence from a Mouse Model of Diabetic Neuropathy. <i>Neuroscience</i> , 2020, 441, 77-92.	1.1	10
23	Neuropathic Pain Dysregulates Gene Expression of the Forebrain Opioid and Dopamine Systems. <i>Neurotoxicity Research</i> , 2020, 37, 800-814.	1.3	29
24	Pharmacological Blockade of Spinal CXCL3/CXCR2 Signaling by NVP CXCR2 20, a Selective CXCR2 Antagonist, Reduces Neuropathic Pain Following Peripheral Nerve Injury. <i>Frontiers in Immunology</i> , 2019, 10, 2198.	2.2	27
25	Chemokines CCL2 and CCL7, but not CCL12, play a significant role in the development of pain-related behavior and opioid-induced analgesia. <i>Cytokine</i> , 2019, 119, 202-213.	1.4	46
26	Mechanisms of Chemotherapy-Induced Peripheral Neuropathy. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1451.	1.8	414
27	Fluorinated indole-imidazole conjugates: Selective orally bioavailable 5-HT7 receptor low-basicity agonists, potential neuropathic painkillers. <i>European Journal of Medicinal Chemistry</i> , 2019, 170, 261-275.	2.6	22
28	Kynurenic acid and zaprinast diminished CXCL17-evoked pain-related behaviour and enhanced morphine analgesia in a mouse neuropathic pain model. <i>Pharmacological Reports</i> , 2019, 71, 139-148.	1.5	20
29	Apelin as a nociceptive processes regulator. <i>BMC</i> , 2019, 19, 1-9.	0.1	0
30	Antinociceptive effects of novel histamine H <sub>3</sub> and H <sub>4</sub> receptor antagonists and their influence on morphine analgesia of neuropathic pain in the mouse. <i>British Journal of Pharmacology</i> , 2018, 175, 2897-2910.	2.7	36
31	Lipopolysaccharide from <i>Rhodobacter sphaeroides</i> (TLR4 antagonist) attenuates hypersensitivity and modulates nociceptive factors. <i>Pharmaceutical Biology</i> , 2018, 56, 275-286.	1.3	18
32	Tapentadol – A representative of a new class of MOR-NRI analgesics. <i>Pharmacological Reports</i> , 2018, 70, 812-820.	1.5	12
33	The importance of chemokines in neuropathic pain development and opioid analgesic potency. <i>Pharmacological Reports</i> , 2018, 70, 821-830.	1.5	42
34	Dataset of (±)-NBI-74330 (CXCR3 antagonist) influence on chemokines under neuropathic pain. <i>Data in Brief</i> , 2018, 21, 1145-1150.	0.5	2
35	Involvement of microglial cells in the antinociceptive effects of metamizol in a mouse model of neuropathic pain. <i>Pharmacology Biochemistry and Behavior</i> , 2018, 175, 77-88.	1.3	11
36	Perioperative Immunosuppression and Risk of Cancer Progression: The Impact of Opioids on Pain Management. <i>Pain Research and Management</i> , 2018, 2018, 1-8.	0.7	20

#	ARTICLE	IF	CITATIONS
37	Zaprinast diminished pain and enhanced opioid analgesia in a rat neuropathic pain model. <i>European Journal of Pharmacology</i> , 2018, 839, 21-32.	1.7	12
38	Alterations in the Activity of Spinal and Thalamic Opioid Systems in a Mice Neuropathic Pain Model. <i>Neuroscience</i> , 2018, 390, 293-302.	1.1	12
39	Pharmacological blockade of CXCR3 by ( $\hat{A}$ $\pm$ )-NBI-74330 reduces neuropathic pain and enhances opioid effectiveness - Evidence from in vivo and in vitro studies. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 3418-3437.	1.8	37
40	Pharmacological Inhibition of Indoleamine 2,3-Dioxygenase-2 and Kynurenine 3-Monooxygenase, Enzymes of the Kynurenine Pathway, Significantly Diminishes Neuropathic Pain in a Rat Model. <i>Frontiers in Pharmacology</i> , 2018, 9, 724.	1.6	41
41	Involvement of Macrophage Inflammatory Protein-1 Family Members in the Development of Diabetic Neuropathy and Their Contribution to Effectiveness of Morphine. <i>Frontiers in Immunology</i> , 2018, 9, 494.	2.2	48
42	Targeting the NLRP3 Inflammasome-Related Pathways via Tianeptine Treatment-Suppressed Microglia Polarization to the M1 Phenotype in Lipopolysaccharide-Stimulated Cultures. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1965.	1.8	84
43	Botulinum Toxin Type A A Modulator of Spinal Neuron-Glia Interactions under Neuropathic Pain Conditions. <i>Toxins</i> , 2018, 10, 145.	1.5	35
44	Non-invasive transcutaneous Supraorbital Neurostimulation (tSNS) using Cefaly $\hat{A}$ device in prevention of primary headaches. <i>Neurologia I Neurochirurgia Polska</i> , 2017, 51, 127-134.	0.6	19
45	Transcranial direct current stimulation (tDCS) and its influence on analgesics effectiveness in patients suffering from migraine headache. <i>Pharmacological Reports</i> , 2017, 69, 714-721.	1.5	35
46	The RS504393 Influences the Level of Nociceptive Factors and Enhances Opioid Analgesic Potency in Neuropathic Rats. <i>Journal of NeuroImmune Pharmacology</i> , 2017, 12, 402-419.	2.1	52
47	Bifunctional opioid/nociceptin hybrid KGNOP1 effectively attenuates pain-related behaviour in a rat model of neuropathy. <i>European Journal of Pharmaceutical Sciences</i> , 2017, 104, 221-229.	1.9	11
48	Spinal CCL1/CCR8 signaling interplay as a potential therapeutic target Evidence from a mouse diabetic neuropathy model. <i>International Immunopharmacology</i> , 2017, 52, 261-271.	1.7	31
49	Analgesic Properties of Opioid/NK1 Multitarget Ligands with Distinct in Vitro Profiles in Naive and Chronic Constriction Injury Mice. <i>ACS Chemical Neuroscience</i> , 2017, 8, 2315-2324.	1.7	30
50	Blockade of P2X4 Receptors Inhibits Neuropathic Pain-Related Behavior by Preventing MMP-9 Activation and, Consequently, Pronociceptive Interleukin Release in a Rat Model. <i>Frontiers in Pharmacology</i> , 2017, 8, 48.	1.6	69
51	Comparison of the Expression Changes after Botulinum Toxin Type A and Minocycline Administration in Lipopolysaccharide-Stimulated Rat Microglial and Astroglial Cultures. <i>Frontiers in Cellular and Infection Microbiology</i> , 2017, 7, 141.	1.8	44
52	Biphalin, a Dimeric Enkephalin, Alleviates LPS-Induced Activation in Rat Primary Microglial Cultures in Opioid Receptor-Dependent and Receptor-Independent Manners. <i>Neural Plasticity</i> , 2017, 2017, 1-19.	1.0	24
53	Blockade of Toll-Like Receptors (TLR2, TLR4) Attenuates Pain and Potentiates Buprenorphine Analgesia in a Rat Neuropathic Pain Model. <i>Neural Plasticity</i> , 2016, 2016, 1-12.	1.0	77
54	Targeting the Microglial Signaling Pathways: New Insights in the Modulation of Neuropathic Pain. <i>Current Medicinal Chemistry</i> , 2016, 23, 2908-2928.	1.2	143

#	ARTICLE	IF	CITATIONS
55	Dataset of botulinum toxin A influence on interleukins under neuropathy. <i>Data in Brief</i> , 2016, 9, 1020-1023.	0.5	2
56	Microglial Inhibition Influences XCL1/XCR1 Expression and Causes Analgesic Effects in a Mouse Model of Diabetic Neuropathy. <i>Anesthesiology</i> , 2016, 125, 573-589.	1.3	37
57	Bifunctional Peptide-Based Opioid Agonist-Nociceptin Antagonist Ligands for Dual Treatment of Acute and Neuropathic Pain. <i>Journal of Medicinal Chemistry</i> , 2016, 59, 3777-3792.	2.9	36
58	Maraviroc reduces neuropathic pain through polarization of microglia and astroglia – Evidence from in vivo and in vitro studies. <i>Neuropharmacology</i> , 2016, 108, 207-219.	2.0	91
59	Intravenous lidocaine infusions in a multidirectional model of treatment of neuropathic pain patients. <i>Pharmacological Reports</i> , 2016, 68, 1069-1075.	1.5	17
60	Participation of pro- and anti-nociceptive interleukins in botulinum toxin A-induced analgesia in a rat model of neuropathic pain. <i>European Journal of Pharmacology</i> , 2016, 791, 377-388.	1.7	57
61	Direct and indirect pharmacological modulation of CCL2/CCR2 pathway results in attenuation of neuropathic pain – In vivo and in vitro evidence. <i>Journal of Neuroimmunology</i> , 2016, 297, 9-19.	1.1	54
62	Characteristics, diagnosis and therapeutic strategies for IgG4-related orbital disease. <i>Pharmacological Reports</i> , 2016, 68, 507-513.	1.5	3
63	Blockade of IL-18 signaling diminished neuropathic pain and enhanced the efficacy of morphine and buprenorphine. <i>Molecular and Cellular Neurosciences</i> , 2016, 71, 114-124.	1.0	65
64	Pharmacological kynurenine 3-monooxygenase enzyme inhibition significantly reduces neuropathic pain in a rat model. <i>Neuropharmacology</i> , 2016, 102, 80-91.	2.0	49
65	Treatment with a carbon monoxide-releasing molecule (CORM-2) inhibits neuropathic pain and enhances opioid effectiveness in rats. <i>Pharmacological Reports</i> , 2016, 68, 206-213.	1.5	25
66	Beneficial properties of maraviroc on neuropathic pain development and opioid effectiveness in rats. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2016, 64, 68-78.	2.5	60
67	Rola cytokin z rodziny interleukiny-1 w transmisji nocyceptywnej. <i>Bł</i> , 2016, 15, 39-47.	0.1	2
68	Zatokowy bł twarzoczaszki – trudnoŃci diagnostyczne w rŃnicowaniu. <i>Bł</i> , 2016, 15, 48-51.	0.1	0
69	Spontaneous cerebrospinal fluid leak at the clivus. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2015, 4, 593-599.	0.3	3
70	A new potential mechanism of action of tianeptine – the effect on microglial cell activation. <i>SpringerPlus</i> , 2015, 4, .	1.2	1
71	Anandamide, Acting via CB2 Receptors, Alleviates LPS-Induced Neuroinflammation in Rat Primary Microglial Cultures. <i>Neural Plasticity</i> , 2015, 2015, 1-10.	1.0	83
72	Parthenolide Relieves Pain and Promotes M2 Microglia/Macrophage Polarization in Rat Model of Neuropathy. <i>Neural Plasticity</i> , 2015, 2015, 1-15.	1.0	80

#	ARTICLE	IF	CITATIONS
73	The Role of Some Chemokines from the CXC Subfamily in a Mouse Model of Diabetic Neuropathy. <i>Journal of Diabetes Research</i> , 2015, 2015, 1-13.	1.0	32
74	Original article IgG4-related inflammatory orbital pseudotumors – a retrospective case series. <i>Folia Neuropathologica</i> , 2015, 2, 111-120.	0.5	7
75	IL-1 receptor antagonist improves morphine and buprenorphine efficacy in a rat neuropathic pain model. <i>European Journal of Pharmacology</i> , 2015, 764, 240-248.	1.7	47
76	Synthesis and biological evaluation of compact, conformationally constrained bifunctional opioid agonist – Neurokinin-1 antagonist peptidomimetics. <i>European Journal of Medicinal Chemistry</i> , 2015, 92, 64-77.	2.6	27
77	Prenatal stress is a vulnerability factor for altered morphology and biological activity of microglia cells. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 82.	1.8	108
78	Effects of chronic doxepin and amitriptyline administration in naïve mice and in neuropathic pain mice model. <i>Neuroscience</i> , 2015, 294, 38-50.	1.1	23
79	Dual Alleviation of Acute and Neuropathic Pain by Fused Opioid Agonist-Neurokinin 1 Antagonist Peptidomimetics. <i>ACS Medicinal Chemistry Letters</i> , 2015, 6, 1209-1214.	1.3	20
80	The influence of microglia activation on the efficacy of amitriptyline, doxepin, milnacipran, venlafaxine and fluoxetine in a rat model of neuropathic pain. <i>European Journal of Pharmacology</i> , 2015, 749, 115-123.	1.7	29
81	Immunoglobulin G4-Related Disease (IgG4-RD) in the Orbit: Mucosa-Associated Lymphoid Tissue (MALT)-Type Lymphomas. <i>Medical Science Monitor</i> , 2015, 21, 1043-1050.	0.5	11
82	PD98059 Influences Immune Factors and Enhances Opioid Analgesia in Model of Neuropathy. <i>PLoS ONE</i> , 2015, 10, e0138583.	1.1	44
83	Castleman's disease of the neck - case report. , 2015, 4, 40-43.		0
84	Marawirok jako potencjalny lek stosowany w terapii bólu neuropatycznego – dowody z badań podstawowych. <i>B&amp;A</i> , 2015, 16, 31-36.	0.1	0
85	Involvement of pro- and antinociceptive factors in minocycline analgesia in rat neuropathic pain model. <i>Journal of Neuroimmunology</i> , 2014, 277, 57-66.	1.1	81
86	Minocycline Enhances the Effectiveness of Nociceptin/Orphanin FQ during Neuropathic Pain. <i>BioMed Research International</i> , 2014, 2014, 1-12.	0.9	28
87	Minocycline prevents dynorphin-induced neurotoxicity during neuropathic pain in rats. <i>Neuropharmacology</i> , 2014, 86, 301-310.	2.0	46
88	Analgesic effects of antidepressants alone and after their local co-administration with morphine in a rat model of neuropathic pain. <i>Pharmacological Reports</i> , 2014, 66, 459-465.	1.5	6
89	Expression Profiling of Genes Modulated by Minocycline in a Rat Model of Neuropathic Pain. <i>Molecular Pain</i> , 2014, 10, 1744-8069-10-47.	1.0	40
90	Inhibition of intracellular signaling pathways NF- $\kappa$ B and MEK1/2 attenuates neuropathic pain development and enhances morphine analgesia. <i>Pharmacological Reports</i> , 2014, 66, 845-851.	1.5	56

#	ARTICLE	IF	CITATIONS
91	Delta-Opioid Receptor Analgesia Is Independent of Microglial Activation in a Rat Model of Neuropathic Pain. PLoS ONE, 2014, 9, e104420.	1.1	74
92	Rola chemokin w b $\alpha$ 3lu neuropatycznym. B $\alpha$ 3l, 2014, 15, 19-35.	0.1	4
93	Minocycline influences the anti-inflammatory interleukins and enhances the effectiveness of morphine under mice diabetic neuropathy. Journal of Neuroimmunology, 2013, 262, 35-45.	1.1	54
94	Effects of selective and non-selective inhibitors of nitric oxide synthase on morphine- and endomorphin-1-induced analgesia in acute and neuropathic pain in rats. Neuropharmacology, 2013, 75, 445-457.	2.0	82
95	Mechanisms and pharmacology of diabetic neuropathy – experimental and clinical studies. Pharmacological Reports, 2013, 65, 1601-1610.	1.5	79
96	Neuronal and immunological basis of action of antidepressants in chronic pain – clinical and experimental studies. Pharmacological Reports, 2013, 65, 1611-1621.	1.5	91
97	Importance of glial activation in neuropathic pain. European Journal of Pharmacology, 2013, 716, 106-119.	1.7	362
98	The glutamatergic system as a target for neuropathic pain relief. Experimental Physiology, 2013, 98, 372-384.	0.9	100
99	Preclinical Cancer Pain Models. , 2013, , 71-93.		0
100	The role of nociceptin and dynorphin in chronic pain: Implications of neuro-glial interaction. Neuropeptides, 2011, 45, 247-261.	0.9	73
101	Glial inhibitors influence the mRNA and protein levels of mGlu2/3, 5 and 7 receptors and potentiate the analgesic effects of their ligands in a mouse model of neuropathic pain. Pain, 2009, 147, 175-186.	2.0	35
102	Differential activation of spinal microglial and astroglial cells in a mouse model of peripheral neuropathic pain. European Journal of Pharmacology, 2009, 623, 65-72.	1.7	160
103	Attenuation of morphine tolerance by minocycline and pentoxifylline in naive and neuropathic mice. Brain, Behavior, and Immunity, 2009, 23, 75-84.	2.0	160
104	Interleukin-1alpha has antiallodynic and antihyperalgesic activities in a rat neuropathic pain model. Pain, 2008, 138, 587-597.	2.0	88
105	Glutamate receptor ligands attenuate allodynia and hyperalgesia and potentiate morphine effects in a mouse model of neuropathic pain. Pain, 2008, 139, 117-126.	2.0	110
106	Modulation of microglia can attenuate neuropathic pain symptoms and enhance morphine effectiveness. Pharmacological Reports, 2008, 60, 297-307.	1.5	147
107	Antinociceptive effect of antisense oligonucleotides against the vanilloid receptor VR1/TRPV1. Neurochemistry International, 2007, 50, 281-290.	1.9	81
108	Minocycline and pentoxifylline attenuate allodynia and hyperalgesia and potentiate the effects of morphine in rat and mouse models of neuropathic pain. European Journal of Pharmacology, 2007, 560, 142-149.	1.7	211



#	ARTICLE	IF	CITATIONS
109	Silencing of vanilloid receptor TRPV1 by RNAi reduces neuropathic and visceral pain in vivo. <i>Biochemical and Biophysical Research Communications</i> , 2006, 350, 238-243.	1.0	147
110	Effects of ORL1 Receptor Agonists and Antagonists in Nociception. <i>Journal of Neuropathic Pain &amp; Symptom Palliation</i> , 2006, 2, 29-44.	0.1	0
111	Analysis of the Dorsal Spinal Cord Synaptic Architecture by Combined Proteome Analysis and in Situ Hybridization. <i>Journal of Proteome Research</i> , 2005, 4, 238-249.	1.8	8
112	Morphine and endomorphin-1 differently influence pronociceptin/orphanin FQ system in neuropathic rats. <i>Pharmacology Biochemistry and Behavior</i> , 2004, 78, 171-178.	1.3	28
113	The Effects of Local Pentoxifylline and Propentofylline Treatment on Formalin-Induced Pain and Tumor Necrosis Factor- $\alpha$ Messenger RNA Levels in the Inflamed Tissue of the Rat Paw. <i>Anesthesia and Analgesia</i> , 2004, 98, 1566-1573.	1.1	52
114	Relationship of pronociceptin/orphanin FQ and the nociceptin receptor ORL1 with substance P and calcitonin gene-related peptide expression in dorsal root ganglion of the rat. <i>Neuroscience Letters</i> , 2003, 348, 190-194.	1.0	26
115	The role of $\delta$ -opioid receptor subtypes in neuropathic pain. <i>European Journal of Pharmacology</i> , 2001, 415, 31-37.	1.7	82
116	Spinal analgesic action of endomorphins in acute, inflammatory and neuropathic pain in rats. <i>European Journal of Pharmacology</i> , 1999, 367, 189-196.	1.7	123
117	Evidence for Fos involvement in the regulation of proenkephalin and prodynorphin gene expression in the rat hippocampus. <i>Molecular Brain Research</i> , 1998, 54, 243-251.	2.5	13
118	Chronic morphine increases biosynthesis of nitric oxide synthase in the rat spinal cord. <i>NeuroReport</i> , 1997, 8, 2743-2747.	0.6	56
119	Season-specific thymic architecture in the frog, <i>Rana temporaria</i> : Sem studies. <i>Developmental and Comparative Immunology</i> , 1996, 20, 129-137.	1.0	11
120	Plasticity of thymuses of ectothermic vertebrates. <i>Trends in Immunology</i> , 1996, 17, 442.	7.5	1
121	Age-dependent changes in thymuses in the European common frog, <i>Rana temporaria</i> . <i>The Journal of Experimental Zoology</i> , 1995, 273, 451-460.	1.4	16
122	Hybrid peptidomimetics for the use in neuropathic pain. , 0, , .		2