

# Clarice S Madruga

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8135271/publications.pdf>

Version: 2024-02-01

31  
papers

620  
citations

686830

13  
h-index

610482

24  
g-index

31  
all docs

31  
docs citations

31  
times ranked

991  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of Cocaine Use in Brazil: Data from the II Brazilian National Alcohol and Drugs Survey (BNADS). <i>Addictive Behaviors</i> , 2014, 39, 297-301.	1.7	118
2	Use of licit and illicit substances among adolescents in Brazil – A national survey. <i>Addictive Behaviors</i> , 2012, 37, 1171-1175.	1.7	74
3	Early handling, but not maternal separation, decreases emotional responses in two paradigms of fear without changes in mesolimbic dopamine. <i>Behavioural Brain Research</i> , 2006, 166, 241-246.	1.2	58
4	Contingency management is effective in promoting abstinence and retention in treatment among crack cocaine users in Brazil: A randomized controlled trial.. <i>Psychology of Addictive Behaviors</i> , 2016, 30, 536-543.	1.4	38
5	Early life exposure to violence and substance misuse in adulthood – The first Brazilian national survey. <i>Addictive Behaviors</i> , 2011, 36, 251-255.	1.7	32
6	Suicidal behavior among substance users: data from the Second Brazilian National Alcohol and Drug Survey (II BNADS). <i>Revista Brasileira De Psiquiatria</i> , 2019, 41, 437-440.	0.9	30
7	Intimate partner violence trends in Brazil: data from two waves of the Brazilian National Alcohol and Drugs Survey. <i>Revista Brasileira De Psiquiatria</i> , 2016, 38, 98-105.	0.9	26
8	Pathways from witnessing parental violence during childhood to involvement in intimate partner violence in adult life: The roles of depression and substance use. <i>Drug and Alcohol Review</i> , 2017, 36, 107-114.	1.1	26
9	Association between drug use and urban violence: Data from the II Brazilian National Alcohol and Drugs Survey (BNADS). <i>Addictive Behaviors Reports</i> , 2018, 7, 8-13.	1.0	25
10	Contingency management targeting abstinence is effective in reducing depressive and anxiety symptoms among crack cocaine-dependent individuals.. <i>Experimental and Clinical Psychopharmacology</i> , 2017, 25, 466-472.	1.3	22
11	Prevalence of and pathways to benzodiazepine use in Brazil: the role of depression, sleep, and sedentary lifestyle. <i>Revista Brasileira De Psiquiatria</i> , 2019, 41, 44-50.	0.9	18
12	How smokers may react to cigarette taxes and price increases in Brazil: data from a national survey. <i>BMC Public Health</i> , 2014, 14, 327.	1.2	17
13	Sociodemographic Characteristics, Patterns of Crack Use, Concomitant Substance Use Disorders, and Psychiatric Symptomatology in Treatment-Seeking Crack-Dependent Individuals in Brazil. <i>Journal of Psychoactive Drugs</i> , 2018, 50, 367-372.	1.0	15
14	Discrepant Trends in Income, Drinking, and Alcohol Problems in an Emergent Economy: Brazil 2006 to 2012. <i>Alcoholism: Clinical and Experimental Research</i> , 2015, 39, 863-871.	1.4	11
15	Crack cocaine users views regarding treatment with contingency management in Brazil. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2018, 13, 7.	1.0	11
16	Alcohol misuse among women in Brazil: recent trends and associations with unprotected sex, early pregnancy, and abortion. <i>Revista Brasileira De Psiquiatria</i> , 2019, 41, 131-137.	0.9	11
17	The Brazilian Drug Policy Situation: The Public Health Approach Based on Research Undertaken in a Developing Country. <i>Public Health Reviews</i> , 2013, 35, .	1.3	10
18	HIV and syphilis infections and associated factors among patients in treatment at a Specialist Alcohol, Tobacco, and Drugs Center in São Paulo – Trends in Psychiatry and Psychotherapy, 2020, 42, 1-6.	0.4	10

#	ARTICLE	IF	CITATIONS
19	Patrones de consumo de alcohol y problemas asociados en Brasil. Revista De Psicologia De La Salud, 2013, 25, 287.	0.2	9
20	Tobacco use among the elderly: The first Brazilian National Survey (BNAS). Aging and Mental Health, 2010, 14, 720-724.	1.5	8
21	Amphetamine-type stimulant use and conditional paths of consumption: data from the Second Brazilian National Alcohol and Drugs Survey. Revista Brasileira De Psiquiatria, 2017, 39, 201-207.	0.9	8
22	Effectiveness of incorporating contingency management into a public treatment program for people who use crack cocaine in Brazil. A single-blind randomized controlled trial. International Journal of Drug Policy, 2022, 99, 103464.	1.6	8
23	Correlated Factors and Prevalence of Alcohol Treatment in Brazil. Addictive Disorders and Their Treatment, 2015, 14, 40-46.	0.5	7
24	Estupros no Brasil e relações com o consumo de álcool: estimativas baseadas em autorrelato sigiloso. Cadernos De Saude Publica, 2019, 35, e00022118.	0.4	7
25	Violence and substance use in sexual minorities: Data from the Second Brazilian National Alcohol and Drugs Survey (II BNADS). Archives of Psychiatric Nursing, 2020, 34, 41-48.	0.7	6
26	Brazilian cross-cultural adaptation and validation of the List of Threatening Events Questionnaire (LTE-Q). Revista Brasileira De Psiquiatria, 2017, 39, 330-336.	0.9	5
27	The effect of religiosity on violence: Results from a Brazilian population-based representative survey of 4,607 individuals. PLoS ONE, 2020, 15, e0238020.	1.1	5
28	Acceptability and feasibility of incorporating contingency management into a public treatment program for homeless crack cocaine users in Brazil: A pilot study.. Experimental and Clinical Psychopharmacology, 2022, 30, 507-513.	1.3	3
29	Commentary on Andreuccetti <i>et al</i> . (2011): The gap between stricter blood alcohol concentration legislation and enforcement in Brazil. Addiction, 2011, 106, 2132-2133.	1.7	2
30	Estupros no Brasil e relações com o consumo de álcool: estimativas baseadas em autorrelato sigiloso. Cadernos De Saude Publica, 2019, 35, .	0.4	0
31	Sociodemographic and clinical profile of crack cocaine treatment-seeking individuals living in Crackland, Brazil. Jornal Brasileiro De Psiquiatria, 2022, 71, 50-55.	0.2	0