

# Jeroen A C M Goos

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8143773/publications.pdf>

Version: 2024-02-01

16  
papers

1,664  
citations

840776

11  
h-index

996975

15  
g-index

25  
all docs

25  
docs citations

25  
times ranked

3297  
citing authors

#	ARTICLE	IF	CITATIONS
1	Optimisation of the Synthesis and Cell Labelling Conditions for [89Zr]Zr-oxine and [89Zr]Zr-DFO-NCS: a Direct In Vitro Comparison in Cell Types with Distinct Therapeutic Applications. <i>Molecular Imaging and Biology</i> , 2021, 23, 952-962.	2.6	4
2	Delivery of polymeric nanostars for molecular imaging and endoradiotherapy through the enhanced permeability and retention (EPR) effect. <i>Theranostics</i> , 2020, 10, 567-584.	10.0	63
3	Expression of the immune modulator secretory leukocyte protease inhibitor (SLPI) in colorectal cancer liver metastases and matched primary tumors is associated with a poorer prognosis. <i>Oncolimmunology</i> , 2020, 9, 1832761.	4.6	10
4	pH-Responsive Polymers for Improving the Signal-to-Noise Ratio of Hypoxia PET Imaging with [18F]Fluoromisonidazole. <i>Macromolecular Rapid Communications</i> , 2020, 41, 2000061.	3.9	4
5	Design and preclinical evaluation of nanostars for the passive pretargeting of tumor tissue. <i>Nuclear Medicine and Biology</i> , 2020, 84-85, 63-72.	0.6	16
6	Temporal Modulation of HER2 Membrane Availability Increases Pertuzumab Uptake and Pretargeted Molecular Imaging of Gastric Tumors. <i>Journal of Nuclear Medicine</i> , 2019, 60, 1569-1578.	5.0	27
7	Biologically Targeted Magnetic Hyperthermia: Potential and Limitations. <i>Frontiers in Pharmacology</i> , 2018, 9, 831.	3.5	340
8	A prognostic classifier for patients with colorectal cancer liver metastasis, based on AURKA, PTGS2 and MMP9. <i>Oncotarget</i> , 2016, 7, 2123-2134.	1.8	17
9	Loss of KCNQ1 expression in stage II and stage III colon cancer is a strong prognostic factor for disease recurrence. <i>British Journal of Cancer</i> , 2016, 115, 1565-1574.	6.4	34
10	Glucose Transporter 1 (SLC2A1) and Vascular Endothelial Growth Factor A (VEGFA) Predict Survival After Resection of Colorectal Cancer Liver Metastasis. <i>Annals of Surgery</i> , 2016, 263, 138-145.	4.2	44
11	Molecular imaging of aurora kinase A (AURKA) expression: Synthesis and preclinical evaluation of radiolabeled alisertib (MLN8237). <i>Nuclear Medicine and Biology</i> , 2016, 43, 63-72.	0.6	9
12	MGL ligand expression is correlated to BRAF mutation and associated with poor survival of stage III colon cancer patients. <i>Oncotarget</i> , 2015, 6, 26278-26290.	1.8	39
13	Combination of Episulfide Ring-Opening Polymerization With ATRP for the Preparation of Amphiphilic Block Copolymers. <i>Macromolecular Rapid Communications</i> , 2013, 34, 156-162.	3.9	15
14	Biomimetic synthesis of calcium carbonate bilayers interfaced by a diblock copolymer template. <i>Zeitschrift Fur Kristallographie - Crystalline Materials</i> , 2012, 227, 739-743.	0.8	0
15	SATB2 in Combination With Cytokeratin 20 Identifies Over 95% of all Colorectal Carcinomas. <i>American Journal of Surgical Pathology</i> , 2011, 35, 937-948.	3.7	209
16	The Initial Stages of Template-Controlled CaCO <sub>3</sub> Formation Revealed by Cryo-TEM. <i>Science</i> , 2009, 323, 1455-1458.	12.6	831