

Rajarshi Banerjee

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8149461/rajarshi-banerjee-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75
papers

2,456
citations

26
h-index

49
g-index

104
ext. papers

3,374
ext. citations

4.9
avg, IF

4.9
L-index

#	Paper	IF	Citations
75	Precision medicine for liver tumours with quantitative MRI and whole genome sequencing (Precision1 trial): study protocol for observational cohort study.. <i>BMJ Open</i> , 2022 , 12, e057163	3	
74	Quantitative magnetic resonance imaging to aid clinical decision making in autoimmune hepatitis.. <i>EClinicalMedicine</i> , 2022 , 46, 101325	11.3	1
73	Quantitative MR in Paediatric Patients with Wilson Disease: A Case Series Review. <i>Children</i> , 2022 , 9, 613 2.8		
72	Multiorgan impairment in low-risk individuals with post-COVID-19 syndrome: a prospective, community-based study. <i>BMJ Open</i> , 2021 , 11, e048391	3	92
71	Repeatability and reproducibility of deep-learning-based liver volume and Couinaud segment volume measurement tool. <i>Abdominal Radiology</i> , 2021 , 1	3	4
70	Multiparametric MRI as a Noninvasive Monitoring Tool for Children With Autoimmune Hepatitis. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2021 , 72, 108-114	2.8	7
69	Comparison between magnetic resonance and ultrasound-derived indicators of hepatic steatosis in a pooled NAFLD cohort. <i>PLoS ONE</i> , 2021 , 16, e0249491	3.7	2
68	Effect of Mastiha supplementation on NAFLD: The MAST4HEALTH Randomised, Controlled Trial. <i>Molecular Nutrition and Food Research</i> , 2021 , 65, e2001178	5.9	6
67	Quantitative multiparametric MRI as a non-invasive stratification tool in children and adolescents with autoimmune liver disease. <i>Scientific Reports</i> , 2021 , 11, 15261	4.9	2
66	rs641738C>T near MBOAT7 is associated with liver fat, ALT and fibrosis in NAFLD: A meta-analysis. <i>Journal of Hepatology</i> , 2021 , 74, 20-30	13.4	24
65	Multiparametric Magnetic Resonance Imaging, Autoimmune Hepatitis, and Prediction of Disease Activity. <i>Hepatology Communications</i> , 2021 , 5, 1009-1020	6	6
64	Quantitative multiparametric magnetic resonance imaging can aid non-alcoholic steatohepatitis diagnosis in a Japanese cohort. <i>World Journal of Gastroenterology</i> , 2021 , 27, 609-623	5.6	8
63	Hepatic Steatosis, Rather Than Underlying Obesity, Increases the Risk of Infection and Hospitalization for COVID-19. <i>Frontiers in Medicine</i> , 2021 , 8, 636637	4.9	6
62	Mastiha has efficacy in immune-mediated inflammatory diseases through a microRNA-155 Th17 dependent action. <i>Pharmacological Research</i> , 2021 , 171, 105753	10.2	2
61	Patient understanding and experience of non-invasive imaging diagnostic techniques and the liver patient pathway. <i>Journal of Patient-Reported Outcomes</i> , 2021 , 5, 89	2.6	0
60	Diagnostic accuracy of elastography and magnetic resonance imaging in patients with NAFLD: A systematic review and meta-analysis. <i>Journal of Hepatology</i> , 2021 , 75, 770-785	13.4	19
59	New boundaries of liver imaging: from morphology to function. <i>European Journal of Internal Medicine</i> , 2020 , 79, 12-22	3.9	2

58	Quantitative MRCP Imaging: Accuracy, Repeatability, Reproducibility, and Cohort-Derived Normative Ranges. <i>Journal of Magnetic Resonance Imaging</i> , 2020 , 52, 807-820	5.6	11
57	Genome-wide and Mendelian randomisation studies of liver MRI yield insights into the pathogenesis of steatohepatitis. <i>Journal of Hepatology</i> , 2020 , 73, 241-251	13.4	28
56	S1178 Utility and Interpretation of the Quantitative MRI Metrics PDFF and cT1 as Biomarkers for Non-Alcoholic Steatohepatitis. <i>American Journal of Gastroenterology</i> , 2020 , 115, S589-S590	0.7	
55	Quantitative magnetic resonance imaging predicts individual future liver performance after liver resection for cancer. <i>PLoS ONE</i> , 2020 , 15, e0238568	3.7	6
54	The Effect of Multi-Parametric Magnetic Resonance Imaging in Standard of Care for Nonalcoholic Fatty Liver Disease: Protocol for a Randomized Control Trial. <i>JMIR Research Protocols</i> , 2020 , 9, e19189	2	2
53	Nutrition and Nonalcoholic Fatty Liver Disease: Current Perspectives. <i>Gastroenterology Clinics of North America</i> , 2020 , 49, 63-94	4.4	26
52	Ethnic Differences in Body Fat Deposition and Liver Fat Content in Two UK-Based Cohorts. <i>Obesity</i> , 2020 , 28, 2142-2152	8	1
51	Multiparametric MR mapping in clinical decision-making for diffuse liver disease. <i>Abdominal Radiology</i> , 2020 , 45, 3507-3522	3	13
50	Prognostic value of multiparametric magnetic resonance imaging, transient elastography and blood-based fibrosis markers in patients with chronic liver disease. <i>Liver International</i> , 2020 , 40, 3071-3082	7.9	17
49	A composite biomarker using multiparametric magnetic resonance imaging and blood analytes accurately identifies patients with non-alcoholic steatohepatitis and significant fibrosis. <i>Scientific Reports</i> , 2020 , 10, 15308	4.9	13
48	NGM282 Improves Liver Fibrosis and Histology in 12 Weeks in Patients With Nonalcoholic Steatohepatitis. <i>Hepatology</i> , 2020 , 71, 1198-1212	11.2	129
47	Correlations Between MRI Biomarkers PDFF and cT1 With Histopathological Features of Non-Alcoholic Steatohepatitis. <i>Frontiers in Endocrinology</i> , 2020 , 11, 575843	5.7	10
46	Genetic studies of abdominal MRI data identify genes regulating hepcidin as major determinants of liver iron concentration. <i>Journal of Hepatology</i> , 2019 , 71, 594-602	13.4	10
45	Repeatability and reproducibility of multiparametric magnetic resonance imaging of the liver. <i>PLoS ONE</i> , 2019 , 14, e0214921	3.7	41
44	Non-invasive assessment of portal hypertension by multi-parametric magnetic resonance imaging of the spleen: A proof of concept study. <i>PLoS ONE</i> , 2019 , 14, e0221066	3.7	19
43	AB064. P-35. Quantitative magnetic resonance cholangiopancreatography applications in primary sclerosing cholangitis and cholangiocarcinoma. <i>Hepatobiliary Surgery and Nutrition</i> , 2019 , 8, AB064-AB064 ^{2,1}		78
42	Hepatic iron is the major determinant of serum ferritin in NAFLD patients. <i>Liver International</i> , 2018 , 38, 164-173	7.9	38
41	Non-invasive assessment of liver disease in rats using multiparametric magnetic resonance imaging: a feasibility study. <i>Biology Open</i> , 2018 , 7,	2.2	11

40	Voxel-Wise Analysis of Paediatric Liver MRI. <i>Communications in Computer and Information Science</i> , 2018 , 57-62	0.3	1
39	Automated Detection of Cystic Lesions in Quantitative T1 Liver Images. <i>Communications in Computer and Information Science</i> , 2018 , 51-56	0.3	
38	Novel Quantitative Magnetic Resonance Imaging Features with Liver Function Tests to Distinguish Parenchymal and Biliary Disease. <i>Communications in Computer and Information Science</i> , 2018 , 37-43	0.3	1
37	Regional Assessment of Liver Disease Progression and Response to Therapy by Multi-time Point m-SLIC Correspondence. <i>Communications in Computer and Information Science</i> , 2018 , 44-50	0.3	
36	The interplay between metabolic alterations, diastolic strain rate and exercise capacity in mild heart failure with preserved ejection fraction: a cardiovascular magnetic resonance study. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2018 , 20, 88	6.9	33
35	Measurement of liver iron by magnetic resonance imaging in the UK Biobank population. <i>PLoS ONE</i> , 2018 , 13, e0209340	3.7	19
34	Study protocol: HepaT1ca - an observational clinical cohort study to quantify liver health in surgical candidates for liver malignancies. <i>BMC Cancer</i> , 2018 , 18, 890	4.8	6
33	Utility and variability of three non-invasive liver fibrosis imaging modalities to evaluate efficacy of GR-MD-02 in subjects with NASH and bridging fibrosis during a phase-2 randomized clinical trial. <i>PLoS ONE</i> , 2018 , 13, e0203054	3.7	36
32	Imaging endpoints for non-alcoholic steatohepatitis (NASH) therapeutic trials: A growing role for multiparametric MRI?. <i>Journal of Hepatology</i> , 2018 , 69, 755-756	13.4	1
31	Reply to: "Multiparametric magnetic resonance imaging to predict clinical outcomes in patients with chronic liver disease: A cautionary note on a promising technique". <i>Journal of Hepatology</i> , 2017 , 66, 457-458	13.4	1
30	A model for hepatic fibrosis: the competing effects of cell loss and iron on shortened modified Look-Locker inversion recovery T (shMOLLI-T) in the liver. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 45, 450-462	5.6	42
29	Interobserver Variability in Histologic Evaluation of Liver Fibrosis Using Categorical and Quantitative Scores. <i>American Journal of Clinical Pathology</i> , 2017 , 147, 364-369	1.9	31
28	Multiparametric magnetic resonance imaging for the assessment of non-alcoholic fatty liver disease severity. <i>Liver International</i> , 2017 , 37, 1065-1073	7.9	103
27	Characterisation of liver fat in the UK Biobank cohort. <i>PLoS ONE</i> , 2017 , 12, e0172921	3.7	51
26	Novel Insights into Complex Cardiovascular Pathologies using 4D Flow Analysis by Cardiovascular Magnetic Resonance Imaging. <i>Current Pharmaceutical Design</i> , 2017 , 23, 3262-3267	3.3	6
25	Ectopic and Visceral Fat Deposition in Lean and Obese Patients With Type 2 Diabetes. <i>Journal of the American College of Cardiology</i> , 2016 , 68, 53-63	15.1	105
24	Investigating a Liver Fat: Arterial Stiffening Pathway in Adult and Childhood Obesity. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016 , 36, 198-203	9.4	16
23	Multiparametric magnetic resonance imaging predicts clinical outcomes in patients with chronic liver disease. <i>Journal of Hepatology</i> , 2016 , 64, 308-315	13.4	127

22	Improvements in ECG accuracy for diagnosis of left ventricular hypertrophy in obesity. <i>Heart</i> , 2016 , 102, 1566-72	5.1	21
21	Fasting Plasma Insulin Concentrations Are Associated With Changes in Hepatic Fatty Acid Synthesis and Partitioning Prior to Changes in Liver Fat Content in Healthy Adults. <i>Diabetes</i> , 2016 , 65, 1858-67	0.9	29
20	Obese subjects show sex-specific differences in right ventricular hypertrophy. <i>Circulation: Cardiovascular Imaging</i> , 2015 , 8,	3.9	15
19	Evidence of a Direct Effect of Myocardial Steatosis on LV Hypertrophy and Diastolic Dysfunction in Adult and Adolescent Obesity. <i>JACC: Cardiovascular Imaging</i> , 2015 , 8, 1468-1470	8.4	14
18	Sex-Specific Differences in Hepatic Fat Oxidation and Synthesis May Explain the Higher Propensity for NAFLD in Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 4425-33	5.6	74
17	Menopausal Status and Abdominal Obesity Are Significant Determinants of Hepatic Lipid Metabolism in Women. <i>Journal of the American Heart Association</i> , 2015 , 4, e002258	6	34
16	Normalization of Visceral Fat and Complete Reversal of Cardiovascular Remodeling Accompany Gastric Bypass, not Banding. <i>Journal of the American College of Cardiology</i> , 2015 , 66, 2569-70	15.1	6
15	Observational study of regional aortic size referenced to body size: production of a cardiovascular magnetic resonance nomogram. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2014 , 16, 9	6.9	57
14	Multiparametric magnetic resonance for the non-invasive diagnosis of liver disease. <i>Journal of Hepatology</i> , 2014 , 60, 69-77	13.4	272
13	HIV is an independent predictor of aortic stiffness. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2014 , 16, 57	6.9	24
12	Structural and functional properties of deep abdominal subcutaneous adipose tissue explain its association with insulin resistance and cardiovascular risk in men. <i>Diabetes Care</i> , 2014 , 37, 821-9	14.6	111
11	Lower resting and total energy expenditure in postmenopausal compared with premenopausal women matched for abdominal obesity. <i>Journal of Nutritional Science</i> , 2014 , 3, e3	2.7	28
10	Normal variation of magnetic resonance T1 relaxation times in the human population at 1.5 T using ShMOLLI. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2013 , 15, 13	6.9	168
9	Gender-specific differences in left ventricular remodelling in obesity: insights from cardiovascular magnetic resonance imaging. <i>European Heart Journal</i> , 2013 , 34, 292-9	9.5	66
8	Response to letter regarding article, "myocardial tissue characterization using magnetic resonance noncontrast t1 mapping in hypertrophic and dilated cardiomyopathy". <i>Circulation: Cardiovascular Imaging</i> , 2013 , 6, e2	3.9	
7	Myocardial tissue characterization using magnetic resonance noncontrast t1 mapping in hypertrophic and dilated cardiomyopathy. <i>Circulation: Cardiovascular Imaging</i> , 2012 , 5, 726-33	3.9	230
6	Arterial stiffness: using simple surrogate measures to make sense of a biologically complex phenomenon. <i>Hypertension Research</i> , 2012 , 35, 155-6	4.7	6
5	Age and gender dependence of pre-contrast T1-relaxation times in normal human myocardium at 1.5T using ShMOLLI. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2012 , 14,	6.9	7

4	Contained left ventricular rupture after acute myocardial infarction revealed by cardiovascular magnetic resonance imaging. <i>Circulation</i> , 2012 , 125, 2278-80	16.7	9
3	rs641738C>T near MBOAT7 is positively associated with liver fat, ALT, and histological severity of NAFLD: a meta-analysis		3
2	High Liver Fat Associates with Higher Risk of Developing Symptomatic COVID-19 Infection - Initial UK Biobank Observations		6
1	Multi-organ impairment in low-risk individuals with long COVID		51