

# Onur Tasar

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8153173/publications.pdf>

Version: 2024-02-01

30  
papers

207  
citations

1040056

9  
h-index

1058476

14  
g-index

32  
all docs

32  
docs citations

32  
times ranked

470  
citing authors

#	ARTICLE	IF	CITATIONS
1	Early Detection of Bi-ventricular and Atrial Mechanical Dysfunction Using Two-Dimensional Speckle Tracking Echocardiography in Patients with Sarcoidosis. <i>Lung</i> , 2015, 193, 669-675.	3.3	30
2	Suboptimal use of non-vitamin K antagonist oral anticoagulants. <i>Medicine (United States)</i> , 2016, 95, e4672.	1.0	28
3	Predictors and outcomes of no-reflow phenomenon in patients with acute ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>Coronary Artery Disease</i> , 2019, 30, 270-276.	0.7	23
4	P wave peak time; a novel electrocardiographic parameter in the assessment of coronary no-reflow. <i>Journal of Electrocardiology</i> , 2017, 50, 584-590.	0.9	21
5	ReAl-life Multicenter Survey Evaluating Stroke prevention strategies in non-valvular atrial fibrillation (RAMSES study). <i>Anatolian Journal of Cardiology</i> , 2016, 16, 734-741.	0.9	17
6	Predictive role of left atrial and ventricular mechanical function in postoperative atrial fibrillation: a two-dimensional speckle-tracking echocardiography study. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2016, 44, 45-52.	0.5	13
7	Evaluation of Left Atrial Functions by 2-dimensional Speckle Tracking Echocardiography During Healthy Pregnancy. <i>Journal of Ultrasound in Medicine</i> , 2019, 38, 2981-2988.	1.7	11
8	Relationship between R-wave peak time and no-reflow in ST elevation myocardial infarction treated with a primary percutaneous coronary intervention. <i>Coronary Artery Disease</i> , 2017, 28, 326-331.	0.7	10
9	Association between BNP levels and new-onset atrial fibrillation. <i>Herz</i> , 2018, 43, 548-554.	1.1	10
10	Guideline-adherent therapy for stroke prevention in atrial fibrillation in different health care settings: Results from RAMSES study. <i>European Journal of Internal Medicine</i> , 2017, 40, 50-55.	2.2	9
11	A successful percutaneous closure of ventricular septal defect following septal myectomy in patients with hypertrophic obstructive cardiomyopathy. <i>Perfusion (United Kingdom)</i> , 2012, 27, 253-255.	1.0	6
12	Real-life use of digoxin in patients with non-valvular atrial fibrillation: data from the RAMSES study. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2016, 41, 711-717.	1.5	6
13	Real-world stroke prevention strategies in nonvalvular atrial fibrillation in patients with renal impairment. <i>European Journal of Clinical Investigation</i> , 2017, 47, 428-438.	3.4	5
14	Multimodality imaging of a left ventricular aneurysm in a patient with normal coronary arteries: Unusual localization. <i>Echocardiography</i> , 2017, 34, 1110-1111.	0.9	4
15	Assessment of the relationship between reperfusion success and T-peak to T-end interval in patients with ST elevation myocardial infarction treated with percutaneous coronary intervention. <i>Anatolian Journal of Cardiology</i> , 2018, 19, 50-57.	0.9	4
16	Impact of valvular heart disease on oral anticoagulant therapy in non-valvular atrial fibrillation: results from the RAMSES study. <i>Journal of Thrombosis and Thrombolysis</i> , 2017, 43, 157-165.	2.1	3
17	Predictors of coronary artery aneurysm after stent implantation in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. <i>International Journal of Cardiovascular Imaging</i> , 2014, 30, 1435-1444.	1.5	2
18	Coronary artery disease in outpatients with nonvalvular atrial fibrillation. <i>Coronary Artery Disease</i> , 2016, 27, 497-503.	0.7	2

#	ARTICLE	IF	CITATIONS
19	Evaluation of aneurysm after coronary stent implantation by optical coherence tomography. <i>Kardiologia Polska</i> , 2013, 71, 659-659.	0.6	1
20	Gender-Related Differences in Presentation and Treatment of Patients With Non-Valvular Atrial Fibrillation: Results from RAMSES study. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2017, 45, 16-25.	0.5	1
21	New and important guide in acute coronary syndrome: optical coherence tomography. <i>Anatolian Journal of Cardiology</i> , 2012, 13, 190.	0.4	0
22	Baseline SYNTAX score and long term outcomes in patients treated with primary percutaneous coronary intervention. <i>European Heart Journal</i> , 2013, 34, P2217-P2217.	2.2	0
23	An interesting diastolic jet in left ventricle. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 399, 413.	0.4	0
24	Baseline red cell distribution width and long term clinical outcomes in patients treated with primary percutaneous coronary intervention. <i>European Heart Journal</i> , 2013, 34, P427-P427.	2.2	0
25	Rescue administration of intracoronary thrombolytic therapy for drug-eluting stent thrombosis. <i>Herz</i> , 2014, 39, 647-650.	1.1	0
26	Optical coherence tomography guided successful fibrinolytic treatment without the need for percutaneous coronary intervention in a patient with acute ST-segment elevation myocardial infarction. <i>Postepy W Kardiologii Interwencyjnej</i> , 2018, 14, 199-201.	0.2	0
27	Optical coherence tomography-verified longer balloon inflation time may provide better stent apposition and optimal index parameters. <i>Herz</i> , 2020, 45, 369-374.	1.1	0
28	Confounding Factors About Microvolt T-wave Alternans Testing And Life Threatening Ventricular Arrhythmias. <i>Anatolian Journal of Cardiology</i> , 2018, 21, 51-52.	0.9	0
29	Rationale, design and methodology of the EPIC (Epidemiology of Polypharmacy and potential) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T 391-398.	0.5	0
30	One-Year Outcomes of Femoropopliteal Chronic Total Occlusions Treated With Percutaneous Provisional Approach: A Single Center Experience (Percutaneous Treatment of Femoropopliteal CTO). <i>European Journal of Therapeutics</i> , 2019, 25, 289-294.	0.1	0