

Swei Sunny Hann

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8153247/publications.pdf>

Version: 2024-02-01

38
papers

1,461
citations

279487

23
h-index

329751

37
g-index

40
all docs

40
docs citations

40
times ranked

1890
citing authors

#	ARTICLE	IF	CITATIONS
1	The regulation and interaction of colon cancer-associated transcriptâ€1 and miR7â€5p contribute to the inhibition of SP1 expression by solamargine in human nasopharyngeal carcinoma cells. <i>Phytotherapy Research</i> , 2020, 34, 201-213.	2.8	17
2	The regulation and interaction of PVT1 and miR181a-5p contributes to the repression of SP1 expression by the combination of XJD decoction and cisplatin in human lung cancer cells. <i>Biomedicine and Pharmacotherapy</i> , 2020, 121, 109632.	2.5	9
3	<p>Biological Roles and Mechanisms of Circular RNA in Human Cancers</p>. <i>OncoTargets and Therapy</i> , 2020, Volume 13, 2067-2092.	1.0	125
4	<p>Regulations of miR-183-5p and Snail-Mediated Shikonin-Reduced Epithelial-Mesenchymal Transition in Cervical Cancer Cells</p>. <i>Drug Design, Development and Therapy</i> , 2020, Volume 14, 577-589.	2.0	19
5	Novel regulation of miRâ€34â€5p and HOTAIR by the combination of berberine and gefitinib leading to inhibition of EMT in human lung cancer. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 5578-5592.	1.6	55
6	Novel reciprocal interaction of lncRNA HOTAIR and miRâ€214â€3p contribute to the solamargineâ€inhibited PDPK1 gene expression in human lung cancer. <i>Journal of Cellular and Molecular Medicine</i> , 2019, 23, 7749-7761.	1.6	23
7	The Reciprocal Interaction Between lncRNA CCAT1 and miR-375-3p Contribute to the Downregulation of IRF5 Gene Expression by Solasonine in HepG2 Human Hepatocellular Carcinoma Cells. <i>Frontiers in Oncology</i> , 2019, 9, 1081.	1.3	20
8	<p>Novel Tumor Suppressor lncRNA Growth Arrest-Specific 5 (GAS5) In Human Cancer</p>. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 8421-8436.	1.0	43
9	Interactions among lncRNAs, miRNAs and mRNA in colorectal cancer. <i>Biochimie</i> , 2019, 163, 58-72.	1.3	81
10	The functions and oncogenic roles of CCAT1 in human cancer. <i>Biomedicine and Pharmacotherapy</i> , 2019, 115, 108943.	2.5	46
11	The repression and reciprocal interaction of DNA methyltransferase 1 and specificity protein 1 contributes to the inhibition of MET expression by the combination of Chinese herbal medicine FZKA decoction and erlotinib. <i>Journal of Ethnopharmacology</i> , 2019, 239, 111928.	2.0	6
12	Reciprocal interaction of HOTAIR and SP1 together enhance the ability of Xiaoji decoction and gefitinib to inhibit EP4 expression. <i>Journal of Ethnopharmacology</i> , 2019, 237, 128-140.	2.0	15
13	<p>Interaction Of c-Jun And HOTAIR- Increased Expression Of p21 Converge In Polyphyllin I-Inhibited Growth Of Human Lung Cancer Cells</p>. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 10115-10127.	1.0	11
14	The enhancement of combination of berberine and metformin in inhibition of DNMT1 gene expression through interplay of SP1 and PDPK1. <i>Journal of Cellular and Molecular Medicine</i> , 2018, 22, 600-612.	1.6	21
15	HOTAIR-mediated reciprocal regulation of EZH2 and DNMT1 contribute to polyphyllin I-inhibited growth of castration-resistant prostate cancer cells in vitro and in vivo. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2018, 1862, 589-599.	1.1	48
16	Repression of PDK1- and lncRNA HOTAIR-Mediated EZH2 Gene Expression Contributes to the Enhancement of Atractylenolide 1 and Erlotinib in the Inhibition of Human Lung Cancer Cells. <i>Cellular Physiology and Biochemistry</i> , 2018, 49, 1615-1632.	1.1	25
17	Inactivation of Stat3 and crosstalk of miRNA155-5p and FOXO3a contribute to the induction of IGFBP1 expression by beta-elemene in human lung cancer. <i>Experimental and Molecular Medicine</i> , 2018, 50, 1-14.	3.2	25
18	Crosstalk of NF-ÎB/P65 and lncRNA HOTAIR-Mediated Repression of MUC1 Expression Contribute to Synergistic Inhibition of Castration-Resistant Prostate Cancer by Polyphyllin 1â€Enzalutamide Combination Treatment. <i>Cellular Physiology and Biochemistry</i> , 2018, 47, 759-773.	1.1	31

#	ARTICLE	IF	CITATIONS
19	HOTAIR: An Oncogenic Long Non-Coding RNA in Human Cancer. <i>Cellular Physiology and Biochemistry</i> , 2018, 47, 893-913.	1.1	212
20	Decoction of Chinese Herbal Medicine Fuzheng Kang-Ai Induces Lung Cancer Cell Apoptosis via STAT3/Bcl-2/Caspase-3 Pathway. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018, 2018, 1-14.	0.5	16
21	Interplay of DNA methyltransferase 1 and EZH2 through inactivation of Stat3 contributes to β -elemene-inhibited growth of nasopharyngeal carcinoma cells. <i>Scientific Reports</i> , 2017, 7, 509.	1.6	38
22	Emodin Increases Expression of Insulin-Like Growth Factor Binding Protein 1 through Activation of MEK/ERK/AMPK \pm and Interaction of PPAR δ and Sp1 in Lung Cancer. <i>Cellular Physiology and Biochemistry</i> , 2017, 41, 339-357.	1.1	46
23	Combination of Solamargine and Metformin Strengthens IGFBP1 Gene Expression Through Inactivation of Stat3 and Reciprocal Interaction Between FOXO3a and SP1. <i>Cellular Physiology and Biochemistry</i> , 2017, 43, 2310-2326.	1.1	16
24	Activation of ERK and Mutual Regulation of Stat3 and SP1 Contribute to Inhibition of PDK1 Expression by Atractylenolide-1 in Human Lung Cancer Cells. <i>Cellular Physiology and Biochemistry</i> , 2017, 43, 2353-2366.	1.1	20
25	Traditional Chinese medicine, Fuzheng Kang-Ai decoction, inhibits metastasis of lung cancer cells through the STAT3/MMP9 pathway. <i>Molecular Medicine Reports</i> , 2017, 16, 2461-2468.	1.1	26
26	Chinese Herbal Medicine Fuzheng Kang-Ai Decoction Inhibited Lung Cancer Cell Growth through AMPK \pm -Mediated Induction and Interplay of IGFBP1 and FOXO3a. <i>Evidence-based Complementary and Alternative Medicine</i> , 2016, 2016, 1-15.	0.5	10
27	Chinese herbal medicine Fuzheng Kang-Ai decoction sensitized the effect of gefitinib on inhibition of human lung cancer cells through inactivating PI3-K/Akt-mediated suppressing MUC1 expression. <i>Journal of Ethnopharmacology</i> , 2016, 194, 918-929.	2.0	20
28	Inter-regulation of IGFBP1 and FOXO3a unveils novel mechanism in ursolic acid-inhibited growth of hepatocellular carcinoma cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2016, 35, 59.	3.5	20
29	Inhibition of EZH2 via activation of SAPK/JNK and reduction of p65 and DNMT1 as a novel mechanism in inhibition of human lung cancer cells by polyphyllin I. <i>Journal of Experimental and Clinical Cancer Research</i> , 2016, 35, 112.	3.5	37
30	Activation of AMPK \pm mediates additive effects of solamargine and metformin on suppressing MUC1 expression in castration-resistant prostate cancer cells. <i>Scientific Reports</i> , 2016, 6, 36721.	1.6	31
31	Inactivation of PI3-K/Akt and reduction of SP1 and p65 expression increase the effect of solamargine on suppressing EP4 expression in human lung cancer cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015, 34, 154.	3.5	34
32	Activation of SAPK/JNK mediated the inhibition and reciprocal interaction of DNA methyltransferase 1 and EZH2 by ursolic acid in human lung cancer cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015, 34, 99.	3.5	32
33	β -elemene inhibited expression of DNA methyltransferase 1 through activation of ERK1/2 and AMPK \pm signalling pathways in human lung cancer cells: the role of Sp1. <i>Journal of Cellular and Molecular Medicine</i> , 2015, 19, 630-641.	1.6	94
34	Combination of curcumin and bicalutamide enhanced the growth inhibition of androgen-independent prostate cancer cells through SAPK/JNK and MEK/ERK1/2-mediated targeting NF- κ B/p65 and MUC1-C. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015, 34, 46.	3.5	48
35	Inhibition of integrin-linked kinase expression by emodin through crosstalk of AMPK \pm and ERK1/2 signaling and reciprocal interplay of Sp1 and c-Jun. <i>Cellular Signalling</i> , 2015, 27, 1469-1477.	1.7	40
36	Baicalein increases the expression and reciprocal interplay of RUNX3 and FOXO3a through crosstalk of AMPK \pm and MEK/ERK1/2 signaling pathways in human non-small cell lung cancer cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015, 34, 41.	3.5	43

#	ARTICLE	IF	CITATIONS
37	Repression of phosphoinositide-dependent protein kinase 1 expression by ciglitazone via Egr-1 represents a new approach for inhibition of lung cancer cell growth. <i>Molecular Cancer</i> , 2014, 13, 149.	7.9	24
38	Targeting EP4 by curcumin through cross talks of AMP-dependent kinase alpha and p38 mitogen-activated protein kinase signaling: The role of PGC-1 β and Sp1. <i>Cellular Signalling</i> , 2013, 25, 2566-2574.	1.7	32