

Maria WoÅnyñec

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8164521/publications.pdf>

Version: 2024-02-01

16
papers

100
citations

1477746

6
h-index

1473754

9
g-index

16
all docs

16
docs citations

16
times ranked

177
citing authors

#	ARTICLE	IF	CITATIONS
1	Cardiovascular Risk Factors Drive Impaired Fasting Glucose to Type 2 Diabetes: Findings After a 9-Year Follow-Up in the PURE Study in Poland. <i>Advances in Experimental Medicine and Biology</i> , 2022, , 1.	0.8	0
2	Obesity and Chosen Non-Communicable Diseases in PURE Poland Cohort Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 2701.	1.2	23
3	Magnetic resonance white matter changes in patients with obstructive sleep apnoea. The subset of the PURE-MIND (Prospective Urban and Rural Epidemiological) cohort study. <i>Archives of Medical Science</i> , 2021, , .	0.4	0
4	Patterns of Alcohol Consumption in the PURE Poland Cohort Study and Their Relationship with Health Problems. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4185.	1.2	6
5	Characteristic of FINDRISC Score and Association with Diabetes Development in 6-Year Follow-Up in PURE Poland Cohort Study. <i>Vascular Health and Risk Management</i> , 2021, Volume 17, 631-639.	1.0	2
6	Anthropometric Indices as Long-Term Predictors of Diabetes in Impaired Fasting Glucose Metabolism: Findings in the PURE Study in Poland. <i>Advances in Experimental Medicine and Biology</i> , 2021, , 79-88.	0.8	2
7	Changes in diabetes prevalence and corresponding risk factors - findings from 3- and 6-year follow-up of PURE Poland cohort study. <i>BMC Public Health</i> , 2020, 20, 843.	1.2	10
8	Prevalence and Risk of Obstructive Sleep Apnea and Arterial Hypertension in the Adult Population in Poland: An Observational Subset of the International Prospective Urban Rural Epidemiology (PURE) Study. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1222, 37-42.	0.8	9
9	Sociodemographic characteristic of changes in smoking patterns in rural and urban population of PURE Poland study: findings from 6-year follow up. <i>BMC Public Health</i> , 2019, 19, 6.	1.2	7
10	Comparison of intake of food groups between participants with normoglycemia, impaired fasting glucose, and type 2 diabetes in PURE Poland population. <i>International Journal of Diabetes in Developing Countries</i> , 2019, 39, 315-324.	0.3	6
11	Differences in prevalence of diabetes mellitus type 2 and impaired fasting glucose between urban and rural areas according to PURE Poland substudy. <i>International Journal of Diabetes in Developing Countries</i> , 2017, 37, 305-312.	0.3	3
12	Prevalence of hypertension in Polish population of PURE Poland study. <i>Journal of Health Inequalities</i> , 2016, 2, 157-162.	0.1	9
13	Serum CETP and PLTP activity in middle-aged men living in urban or rural area of the Lower Silesia region. PURE Poland sub-study. <i>Archives of Medical Science</i> , 2016, 4, 704-714.	0.4	2
14	Serum Lipid Transfer Proteins in Hypothyreotic Patients Are Inversely Correlated with Thyroid-Stimulating Hormone (TSH) Levels. <i>Medical Science Monitor</i> , 2016, 22, 4661-4669.	0.5	9
15	Lipid pattern in middle-aged inhabitants of the Lower Silesian region of Poland. The PURE Poland sub-study. <i>Annals of Agricultural and Environmental Medicine</i> , 2013, 20, 317-24.	0.5	4
16	Prevalence of diabetes mellitus and IFG in the prospective cohort 'PONS' study - baseline assessment. <i>Annals of Agricultural and Environmental Medicine</i> , 2011, 18, 265-9.	0.5	8