

Maren Falch Lindberg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8174051/publications.pdf>

Version: 2024-02-01

18
papers

270
citations

1039406

9
h-index

940134

16
g-index

18
all docs

18
docs citations

18
times ranked

295
citing authors

#	ARTICLE	IF	CITATIONS
1	Many Patients With Persistent Pain 1 Year After TKA Report Improvement by 5 to 7 Years: A Mixed-methods Study. <i>Clinical Orthopaedics and Related Research</i> , 2022, 480, 2075-2088.	0.7	4
2	Factors Correlated With Physical Function 1 Year After Total Knee Arthroplasty in Patients With Knee Osteoarthritis. <i>JAMA Network Open</i> , 2022, 5, e2219636.	2.8	12
3	Preoperative risk factors associated with chronic pain profiles following total knee arthroplasty. <i>European Journal of Pain</i> , 2021, 25, 680-692.	1.4	23
4	Authors'™ reply to the comment by Riddle et al. <i>European Journal of Pain</i> , 2021, 25, 1173-1174.	1.4	0
5	Development of an internet-delivered cognitive behavioral therapy program for use in combination with exercise therapy and education by patients at increased risk of chronic pain following total knee arthroplasty. <i>BMC Health Services Research</i> , 2021, 21, 1151.	0.9	9
6	The effectiveness of exercise therapy and education plus cognitive behavioral therapy, alone or in combination with total knee arthroplasty in patients with knee osteoarthritis – study protocol for the MultiKnee trial. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 1054.	0.8	7
7	High levels of preoperative pain and fatigue are red flags for moderate–severe pain 12 months after total knee arthroplasty—A longitudinal cohort study. <i>Musculoskeletal Care</i> , 2020, 19, 186-192.	0.6	8
8	More Severe Radiographic Osteoarthritis Is Associated With Increased Improvement in Patients'™ Health State Following a Total Knee Arthroplasty. <i>Journal of Arthroplasty</i> , 2020, 35, 3131-3137.	1.5	9
9	Predictors of chronic pain and level of physical function in total knee arthroplasty: a protocol for a systematic review and meta-analysis. <i>BMJ Open</i> , 2020, 10, e037674.	0.8	3
10	High pre- and postoperative symptom burden in non-responders to total knee arthroplasty. <i>PLoS ONE</i> , 2020, 15, e0233347.	1.1	4
11	Psychometric properties of a short version of Lee Fatigue Scale used as a generic PROM in persons with stroke or osteoarthritis: assessment using a Rasch analysis approach. <i>Health and Quality of Life Outcomes</i> , 2020, 18, 168.	1.0	5
12	Higher physical activity and lower pain levels before surgery predict non-improvement of knee pain 1 year after TKA. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 1698-1708.	2.3	21
13	The Impact of Demographic, Clinical, Symptom and Psychological Characteristics on the Trajectories of Acute Postoperative Pain After Total Knee Arthroplasty. <i>Pain Medicine</i> , 2017, 18, 124-139.	0.9	25
14	The relationship between pain with walking and self-rated health 12 months following total knee arthroplasty: a longitudinal study. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 75.	0.8	13
15	Factors Related to Postoperative Pain Trajectories following Total Knee Arthroplasty: A Longitudinal Study of Patients Admitted to a Russian Orthopaedic Clinic. <i>Pain Research and Treatment</i> , 2016, 2016, 1-12.	1.7	24
16	Preoperative Pain, Symptoms, and Psychological Factors related to Higher Acute Pain Trajectories during Hospitalization for Total Knee Arthroplasty. <i>PLoS ONE</i> , 2016, 11, e0161681.	1.1	38
17	Factors that can predict pain with walking, 12 months after total knee arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 600-606.	1.2	41
18	Pain characteristics and self-rated health after elective orthopaedic surgery – a cross-sectional survey. <i>Journal of Clinical Nursing</i> , 2013, 22, 1242-1253.	1.4	24