

# James Mugisha

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8179091/publications.pdf>

Version: 2024-02-01

100  
papers

3,567  
citations

172386

29  
h-index

155592

55  
g-index

100  
all docs

100  
docs citations

100  
times ranked

4845  
citing authors

#	ARTICLE	IF	CITATIONS
1	Physical and mental health impact of COVID-19 on children, adolescents, and their families: The Collaborative Outcomes study on Health and Functioning during Infection Times - Children and Adolescents (COH-FIT&A). <i>Journal of Affective Disorders</i> , 2022, 299, 367-376.	2.0	33
2	Towards Entrapment: An Escalating Reality for Children and Adolescents Living with HIV/AIDS in Uganda. , 2022, , 145-153.		0
3	Mental contrasting and implementation of physical activity intentions in Ugandan primary care patients with mental health problems: A real-world intervention involving support partners. <i>Psychiatry Research</i> , 2022, 307, 114335.	1.7	5
4	Less time spent walking and depressive symptoms are associated with more self-perceived disability in primary care patients: a cross-sectional study from Uganda. <i>Pan African Medical Journal</i> , 2022, 41, 173.	0.3	0
5	Suubi4Stigma study protocol: a pilot cluster randomized controlled trial to address HIV-associated stigma among adolescents living with HIV in Uganda. <i>Pilot and Feasibility Studies</i> , 2022, 8, 95.	0.5	8
6	The efficacy of meditation-based mind-body interventions for mental disorders: A meta-review of 17 meta-analyses of randomized controlled trials. <i>Journal of Psychiatric Research</i> , 2021, 134, 181-191.	1.5	35
7	The efficacy of a lay health workers ‘‘ led physical activity counselling program in patients with HIV and mental health problems: a real-world intervention from Uganda. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2021, 33, 1189-1195.	0.6	10
8	Sociocultural Aspects of Health Promotion in Palliative Care in Uganda. , 2021, , 303-312.		1
9	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with alcohol use disorder. <i>Alcohol</i> , 2021, 90, 74-79.	0.8	2
10	Physical activity, physical fitness and quality of life in outpatients with a psychotic disorder versus healthy matched controls in a low-income country. <i>Schizophrenia Research</i> , 2021, 229, 1-2.	1.1	3
11	Correlates of missed HIV appointments in low-resource settings: a study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 125-131.	0.3	0
12	Re-conceptualizing sustainable urban sanitation in Uganda: why the roots of ‘‘Slumification’’ must be dealt with. <i>BMC Public Health</i> , 2021, 21, 992.	1.2	6
13	Effectiveness and cost-effectiveness of integrating the management of depression into routine HIV Care in Uganda (the HIV+D trial): A protocol for a cluster-randomised trial. <i>International Journal of Mental Health Systems</i> , 2021, 15, 45.	1.1	4
14	Religion as Meaning-Making Resource in Understanding Suicidal Behavior in Ghana and Uganda. <i>Frontiers in Psychology</i> , 2021, 12, 549404.	1.1	4
15	Stakeholders’ perspectives on integrating the management of depression into routine HIV care in Uganda: qualitative findings from a feasibility study. <i>International Journal of Mental Health Systems</i> , 2021, 15, 63.	1.1	2
16	A Struggle for Survival: Meaning of Late Life in a Rural District in Uganda: A Qualitative Study. <i>Frontiers in Psychology</i> , 2021, 12, 699485.	1.1	0
17	Physical activity, physical fitness and quality of life in outpatients with major depressive disorder versus matched healthy controls: Data from a low-income country. <i>Journal of Affective Disorders</i> , 2021, 294, 802-804.	2.0	4
18	Internalised HIV-related stigma associated with physical inactivity in people with HIV and AIDS: A cross-sectional study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 238-243.	0.3	4

#	ARTICLE	IF	CITATIONS
19	“When will the doctor be around so that I come by?” Geo-socio effects on health care supply, access and utilisation: experiences from Kalangala Islands, Uganda. BMC Health Services Research, 2021, 21, 1163.	0.9	1
20	Moderate to severe generalized anxiety disorder symptoms are associated with physical inactivity in people with HIV/AIDS: a study from Uganda. International Journal of STD and AIDS, 2021, 32, 170-175.	0.5	6
21	A 10-hours workshop improves physical activity prescription for mental illness knowledge and confidence in health care professionals: a nation-wide multicentre study from Uganda. Disability and Rehabilitation, 2021, , 1-6.	0.9	5
22	The efficacy of physical activity counseling in Ugandan patients with HIV and a co-morbid mental disorder: a pilot study. AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV, 2020, 32, 758-761.	0.6	6
23	Assessing quality of life in older adults: psychometric properties of the OPQoL-brief questionnaire in a nursing home population. Health and Quality of Life Outcomes, 2020, 18, 1.	1.0	140
24	Test-retest reliability, validity, and correlates of the 2-min walk test in outpatients with depression. Physiotherapy Research International, 2020, 25, e1821.	0.7	8
25	Correlates of physical activity stages of change in people living with HIV in a Ugandan community. Disability and Rehabilitation, 2020, , 1-6.	0.9	4
26	Health care professionals’ perspectives on barriers to treatment seeking for formal health services among orphan children and adolescents with HIV/AIDS and mental distress in a rural district in central, Uganda. Child and Adolescent Psychiatry and Mental Health, 2020, 14, 26.	1.2	5
27	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. African Health Sciences, 2020, 20, 1407-1415.	0.3	0
28	Test-retest reliability and correlates of the Simple Physical Activity Questionnaire in Ugandan out-patients with psychosis. African Health Sciences, 2020, 20, 1438-1445.	0.3	1
29	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. African Health Sciences, 2020, 20, 1407-1415.	0.3	5
30	Test-retest reliability and correlates of the Simple Physical Activity Questionnaire in Ugandan out-patients with psychosis. African Health Sciences, 2020, 20, 1438-1445.	0.3	4
31	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with psychosis. Psychiatry Research, 2019, 282, 112619.	1.7	5
32	Motives for physical activity in the adoption and maintenance of physical activity in middle-aged and old age outpatients with a mental disorder: A cross-sectional study from a low-income country. Psychiatry Research, 2019, 282, 112620.	1.7	5
33	A quantitative assessment of the views of mental health professionals on exercise for people with mental illness: perspectives from a low-resource setting. African Health Sciences, 2019, 19, 2172.	0.3	9
34	Experience of implementing new mental health indicators within information systems in six low- and middle-income countries. BJPsych Open, 2019, 5, e71.	0.3	12
35	The experience of mental health service users in health system strengthening: lessons from Uganda. International Journal of Mental Health Systems, 2019, 13, 60.	1.1	11
36	Associations between physical inactivity, major depressive disorder, and alcohol use disorder in people living with HIV in a Ugandan fishing community. International Journal of STD and AIDS, 2019, 30, 1177-1184.	0.5	6

#	ARTICLE	IF	CITATIONS
37	Scaling up integrated primary mental health in six low- and middle-income countries: obstacles, synergies and implications for systems reform. <i>BJPsych Open</i> , 2019, 5, e69.	0.3	51
38	Livelihood Risk, Culture, and the HIV Interface: Evidence from Lakeshore Border Communities in Buliisa District, Uganda. <i>Journal of Tropical Medicine</i> , 2019, 2019, 1-10.	0.6	10
39	Associations of the Built Environment With Physical Activity and Sedentary Time in Ugandan Outpatients With Mental Health Problems. <i>Journal of Physical Activity and Health</i> , 2019, 16, 243-250.	1.0	11
40	Sedentary behavior and anxiety-induced sleep disturbance among 181,093 adolescents from 67 countries: a global perspective. <i>Sleep Medicine</i> , 2019, 58, 19-26.	0.8	33
41	Leisure-time sedentary behavior and suicide attempt among 126,392 adolescents in 43 countries. <i>Journal of Affective Disorders</i> , 2019, 250, 346-353.	2.0	32
42	Prevalence of behavioral disorders and attention deficit/hyperactive disorder among school going children in Southwestern Uganda. <i>BMC Psychiatry</i> , 2019, 19, 105.	1.1	12
43	Health care professionalsâ€™ perspectives on physical activity within the Ugandan mental health care system. <i>Mental Health and Physical Activity</i> , 2019, 16, 1-7.	0.9	12
44	Evaluation of performance and perceived utility of mental healthcare indicators in routine health information systems in five low- and middle-income countries. <i>BJPsych Open</i> , 2019, 5, e70.	0.3	13
45	Mental health financing challenges, opportunities and strategies in low- and middle-income countries: findings from the Emerald project. <i>BJPsych Open</i> , 2019, 5, e68.	0.3	41
46	Peer support for frequent users of inpatient mental health care in Uganda: protocol of a quasi-experimental study. <i>BMC Psychiatry</i> , 2019, 19, 374.	1.1	10
47	Sedentary Behavior and Quality of Life in People with Psychotic Disorders from a Low Income Country: A Study from Uganda. <i>Community Mental Health Journal</i> , 2019, 55, 714-720.	1.1	1
48	Barriers, attitudes, confidence, and knowledge of nurses regarding metabolic health screening and intervention in people with mental illness: a pilot study from Uganda. <i>African Health Sciences</i> , 2019, 19, 2546-2554.	0.3	4
49	Correlates of sedentary behavior in 2,375 people with depression from 6 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2018, 234, 97-104.	2.0	22
50	Motives for physical activity in the adoption and maintenance of physical activity in men with alcohol use disorders. <i>Psychiatry Research</i> , 2018, 261, 522-526.	1.7	9
51	Correlates of Physical Activity Among Middle-Aged and Older Adults With Hazardous Drinking Habits in Six Low- and Middle-Income Countries. <i>Journal of Aging and Physical Activity</i> , 2018, 26, 589-598.	0.5	5
52	Adherence to physical activity recommendations and physical and mental health risk in people with severe mental illness in Uganda. <i>Psychiatry Research</i> , 2018, 260, 236-240.	1.7	11
53	Global physical activity levels among people living with HIV: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2018, 40, 388-397.	0.9	100
54	Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. <i>Disability and Rehabilitation</i> , 2018, 40, 1618-1629.	0.9	65

#	ARTICLE	IF	CITATIONS
55	Correlates of physical activity among depressed older people in six low-income and middle-income countries: A community-based cross-sectional study. <i>International Journal of Geriatric Psychiatry</i> , 2018, 33, e314-e322.	1.3	15
56	Physical activity and suicidal ideation: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2018, 225, 438-448.	2.0	140
57	Exercise self-efficacy correlates in people with psychosis. <i>Psychiatry Research</i> , 2018, 262, 359-362.	1.7	5
58	Gender differences in motives for physical activity across the stages of change in Ugandan outpatients with psychosis. <i>Schizophrenia Research</i> , 2018, 197, 568-569.	1.1	4
59	Correlates of physical activity among community-dwelling individuals aged 65 years or older with anxiety in six low- and middle-income countries. <i>International Psychogeriatrics</i> , 2018, 30, 705-714.	0.6	10
60	Physical activity and HIV in sub-Saharan Africa: a systematic review of correlates and levels. <i>African Health Sciences</i> , 2018, 18, 394-406.	0.3	15
61	Pain and severe sleep disturbance in the general population: Primary data and meta-analysis from 240,820 people across 45 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2018, 53, 52-58.	1.2	14
62	Cultural responses towards the aftermath of suicide among the Acholi in Northern Uganda. <i>International Journal of Social Psychiatry</i> , 2018, 64, 545-553.	1.6	4
63	Physiotherapy for people with mental health problems in Sub-Saharan African countries: a systematic review. <i>Archives of Physiotherapy</i> , 2018, 8, 2.	0.7	8
64	Physical Activity Levels and Psychosis: A Mediation Analysis of Factors Influencing Physical Activity Target Achievement Among 204 186 People Across 46 Low- and Middle-Income Countries. <i>Schizophrenia Bulletin</i> , 2017, 43, sbw111.	2.3	49
65	Cardiorespiratory Fitness in Severe Mental Illness: A Systematic Review and Meta-analysis. <i>Sports Medicine</i> , 2017, 47, 343-352.	3.1	170
66	Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, 6.	2.0	115
67	Health systems context(s) for integrating mental health into primary health care in six Emerald countries: a situation analysis. <i>International Journal of Mental Health Systems</i> , 2017, 11, 7.	1.1	55
68	Perceived Stress and Its Relationship With Chronic Medical Conditions and Multimorbidity Among 229,293 Community-Dwelling Adults in 44 Low- and Middle-Income Countries. <i>American Journal of Epidemiology</i> , 2017, 186, 979-989.	1.6	62
69	Interest, competence, appearance, fitness and social relatedness as motives for physical activity in Ugandan outpatients with psychosis. <i>Mental Health and Physical Activity</i> , 2017, 13, 94-99.	0.9	6
70	Physical activity correlates among 24,230 people with depression across 46 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2017, 221, 81-88.	2.0	21
71	Strengthening mental health system governance in six low- and middle-income countries in Africa and South Asia: challenges, needs and potential strategies. <i>Health Policy and Planning</i> , 2017, 32, 699-709.	1.0	127
72	Sedentary Behavior in People Living With HIV: A Systematic Review and Meta-Analysis. <i>Journal of Physical Activity and Health</i> , 2017, 14, 571-577.	1.0	22

#	ARTICLE	IF	CITATIONS
73	Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. <i>World Psychiatry</i> , 2017, 16, 308-315.	4.8	600
74	Physical activity is associated with the physical, psychological, social and environmental quality of life in people with mental health problems in a low resource setting. <i>Psychiatry Research</i> , 2017, 258, 250-254.	1.7	25
75	Lifetime self-reported arthritis is associated with elevated levels of mental health burden: A multi-national cross sectional study across 46 low- and middle-income countries. <i>Scientific Reports</i> , 2017, 7, 7138.	1.6	11
76	Physical health policies and metabolic screening in mental health care systems of sub-Saharan African countries: a systematic review. <i>International Journal of Mental Health Systems</i> , 2017, 11, 31.	1.1	13
77	Physical activity and anxiety: A perspective from the World Health Survey. <i>Journal of Affective Disorders</i> , 2017, 208, 545-552.	2.0	211
78	A systematic review of physical activity policy recommendations and interventions for people with mental health problems in Sub-Saharan African countries. <i>Pan African Medical Journal</i> , 2017, 26, 104.	0.3	20
79	The experiences of caregivers of children living with HIV and AIDS in Uganda: a qualitative study. <i>Globalization and Health</i> , 2017, 13, 72.	2.4	35
80	Growing up HIV-positive in Uganda: "psychological immunodeficiency" A qualitative study. <i>BMC Psychology</i> , 2017, 5, 30.	0.9	15
81	Prevalence of suicidal ideation and attempt: associations with psychiatric disorders and HIV/AIDS in post-conflict Northern Uganda. <i>African Health Sciences</i> , 2017, 16, 1027.	0.3	18
82	Physical activity and sedentary behavior in people with bipolar disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 201, 145-152.	2.0	109
83	Indicators for routine monitoring of effective mental healthcare coverage in low- and middle-income settings: a Delphi study. <i>Health Policy and Planning</i> , 2016, 31, 1100-1106.	1.0	36
84	The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2016, 43, 63-70.	1.2	122
85	Cardiorespiratory fitness levels and moderators in people with HIV: A systematic review and meta-analysis. <i>Preventive Medicine</i> , 2016, 93, 106-114.	1.6	36
86	Physical multimorbidity and psychosis: comprehensive cross sectional analysis including 242,952 people across 48 low- and middle-income countries. <i>BMC Medicine</i> , 2016, 14, 189.	2.3	119
87	The prevalence of diabetes mellitus type 2 in people with alcohol use disorders: a systematic review and large scale meta-analysis. <i>Psychiatry Research</i> , 2016, 246, 394-400.	1.7	27
88	The Functional Exercise Capacity Is Associated With Global Functioning in People With Bipolar Disorder. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 673-677.	0.5	7
89	The Prevalence of Metabolic Syndrome in Alcohol Use Disorders: A Systematic Review and Meta-analysis. <i>Alcohol and Alcoholism</i> , 2016, 51, 515-521.	0.9	72
90	Towards understanding governance issues in integration of mental health into primary health care in Uganda. <i>International Journal of Mental Health Systems</i> , 2016, 10, 25.	1.1	58

#	ARTICLE	IF	CITATIONS
91	Service user and caregiver involvement in mental health system strengthening in low- and middle-income countries: systematic review. <i>BMC Health Services Research</i> , 2016, 16, 79.	0.9	175
92	Physical activity as a vital sign in patients with schizophrenia: Evidence and clinical recommendations. <i>Schizophrenia Research</i> , 2016, 170, 336-340.	1.1	32
93	Prevalence and factors associated with Posttraumatic Stress Disorder seven years after the conflict in three districts in northern Uganda (The Wayo-Nero Study). <i>BMC Psychiatry</i> , 2015, 15, 170.	1.1	47
94	Explaining the slow transition of child-appropriate dosage formulations from the global to national level in the context of Uganda: a qualitative study. <i>Journal of Pharmaceutical Policy and Practice</i> , 2015, 8, 19.	1.1	7
95	Major depressive disorder seven years after the conflict in northern Uganda: burden, risk factors and impact on outcomes (The Wayo-Nero Study). <i>BMC Psychiatry</i> , 2015, 15, 48.	1.1	39
96	Stock-outs, uncertainty and improvisation in access to healthcare in war-torn Northern Uganda. <i>Social Science and Medicine</i> , 2015, 146, 316-323.	1.8	20
97	The internal dialogue between the individual and the community: A discourse analysis of public views on suicide among the Baganda, Uganda. <i>International Journal of Culture and Mental Health</i> , 2014, 7, 122-136.	0.6	6
98	Religious Views on Suicide Among the Baganda, Uganda: A Qualitative Study. <i>Death Studies</i> , 2013, 37, 343-361.	1.8	17
99	Distancing: A traditional mechanism of dealing with suicide among the Baganda, Uganda. <i>Transcultural Psychiatry</i> , 2011, 48, 624-642.	0.9	34
100	Doing Qualitative Research on Suicide in a Developing Country. <i>Crisis</i> , 2011, 32, 15-23.	0.9	18