

# Per Holmich

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/8185404/per-holmich-publications-by-citations.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

221  
papers

6,955  
citations

43  
h-index

78  
g-index

246  
ext. papers

8,208  
ext. citations

5.7  
avg, IF

6.09  
L-index

| #   | Paper                                                                                                                                                                                                                               | IF   | Citations |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 221 | The Warwick Agreement on femoroacetabular impingement syndrome (FAI syndrome): an international consensus statement. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 1169-76                                          | 10.3 | 490       |
| 220 | Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. <i>American Journal of Sports Medicine</i> , <b>2011</b> , 39, 2296-303                                 | 6.8  | 366       |
| 219 | Effectiveness of active physical training as treatment for long-standing adductor-related groin pain in athletes: randomised trial. <i>Lancet, The</i> , <b>1999</b> , 353, 439-43                                                  | 40   | 357       |
| 218 | The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. <i>British Journal of Sports Medicine</i> , <b>2011</b> , 45, 478-91                                              | 10.3 | 328       |
| 217 | Doha agreement meeting on terminology and definitions in groin pain in athletes. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 768-74                                                                               | 10.3 | 277       |
| 216 | Clinical assessment of hip strength using a hand-held dynamometer is reliable. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2010</b> , 20, 493-501                                                            | 4.6  | 267       |
| 215 | Long-standing groin pain in sportspeople falls into three primary patterns, a "clinical entity" approach: a prospective study of 207 patients. <i>British Journal of Sports Medicine</i> , <b>2007</b> , 41, 247-52; discussion 252 | 10.3 | 171       |
| 214 | Evidence based prevention of hamstring injuries in sport. <i>British Journal of Sports Medicine</i> , <b>2005</b> , 39, 319-23                                                                                                      | 10.3 | 152       |
| 213 | Hip- and knee-strength assessments using a hand-held dynamometer with external belt-fixation are inter-tester reliable. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2013</b> , 21, 550-5                             | 5.5  | 135       |
| 212 | Validity, reliability and responsiveness of patient-reported outcome questionnaires when assessing hip and groin disability: a systematic review. <i>British Journal of Sports Medicine</i> , <b>2010</b> , 44, 1186-96             | 10.3 | 128       |
| 211 | Diagnostic accuracy of clinical tests for the diagnosis of hip femoroacetabular impingement/labral tear: a systematic review with meta-analysis. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 811                  | 10.3 | 120       |
| 210 | Patient-Reported Outcome (PRO) questionnaires for young to middle-aged adults with hip and groin disability: a systematic review of the clinimetric evidence. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 812     | 10.3 | 113       |
| 209 | Clinical examination of athletes with groin pain: an intraobserver and interobserver reliability study. <i>British Journal of Sports Medicine</i> , <b>2004</b> , 38, 446-51                                                        | 10.3 | 109       |
| 208 | Hip adduction and abduction strength profiles in elite soccer players: implications for clinical evaluation of hip adductor muscle recovery after injury. <i>American Journal of Sports Medicine</i> , <b>2011</b> , 39, 121-6      | 6.8  | 103       |
| 207 | Incidence and clinical presentation of groin injuries in sub-elite male soccer. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 1245-50                                                                               | 10.3 | 102       |
| 206 | Diagnosis of Acute Groin Injuries: A Prospective Study of 110 Athletes. <i>American Journal of Sports Medicine</i> , <b>2015</b> , 43, 1857-64                                                                                      | 6.8  | 95        |
| 205 | Which factors differentiate athletes with hip/groin pain from those without? A systematic review with meta-analysis. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 810                                              | 10.3 | 91        |

|     |                                                                                                                                                                                                                                                 |      |    |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 204 | Exercise program for prevention of groin pain in football players: a cluster-randomized trial. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2010</b> , 20, 814-21                                                         | 4.6  | 91 |
| 203 | Prevention of overuse injuries by a concurrent exercise program in subjects exposed to an increase in training load: a randomized controlled trial of 1020 army recruits. <i>American Journal of Sports Medicine</i> , <b>2008</b> , 36, 663-70 | 6.8  | 81 |
| 202 | Groin Problems in Male Soccer Players Are More Common Than Previously Reported. <i>American Journal of Sports Medicine</i> , <b>2017</b> , 45, 1304-1308                                                                                        | 6.8  | 78 |
| 201 | Prevalence of radiological signs of femoroacetabular impingement in patients presenting with long-standing adductor-related groin pain. <i>British Journal of Sports Medicine</i> , <b>2011</b> , 45, 6-9                                       | 10.3 | 77 |
| 200 | Eccentric hip adduction and abduction strength in elite soccer players and matched controls: a cross-sectional study. <i>British Journal of Sports Medicine</i> , <b>2011</b> , 45, 10-3                                                        | 10.3 | 76 |
| 199 | Study quality on groin injury management remains low: a systematic review on treatment of groin pain in athletes. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 813                                                             | 10.3 | 75 |
| 198 | The Adductor Strengthening Programme prevents groin problems among male football players: a cluster-randomised controlled trial. <i>British Journal of Sports Medicine</i> , <b>2019</b> , 53, 150-157                                          | 10.3 | 66 |
| 197 | Prevalence and severity of hip and groin pain in sub-elite male football: a cross-sectional cohort study of 695 players. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2017</b> , 27, 107-114                              | 4.6  | 65 |
| 196 | EMG evaluation of hip adduction exercises for soccer players: implications for exercise selection in prevention and treatment of groin injuries. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 1108-14                          | 10.3 | 65 |
| 195 | Copenhagen hip and groin outcome score (HAGOS) in male soccer: reference values for hip and groin injury-free players. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 557-9                                                      | 10.3 | 60 |
| 194 | Radiological findings in symphyseal and adductor-related groin pain in athletes: a critical review of the literature. <i>British Journal of Sports Medicine</i> , <b>2013</b> , 47, 611-9                                                       | 10.3 | 59 |
| 193 | Effects of evidence-based prevention training on neuromuscular and biomechanical risk factors for ACL injury in adolescent female athletes: a randomised controlled trial. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 552-7  | 10.3 | 58 |
| 192 | Eccentric and Isometric Hip Adduction Strength in Male Soccer Players With and Without Adductor-Related Groin Pain: An Assessor-Blinded Comparison. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2014</b> , 2, 2325967114521778           | 3.5  | 57 |
| 191 | Return to Sport and Performance After Hip Arthroscopy for Femoroacetabular Impingement in 18- to 30-Year-Old Athletes: A Cross-sectional Cohort Study of 189 Athletes. <i>American Journal of Sports Medicine</i> , <b>2018</b> , 46, 2578-2587 | 6.8  | 56 |
| 190 | Hip strength assessment using handheld dynamometry is subject to intertester bias when testers are of different sex and strength. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2013</b> , 23, 487-93                      | 4.6  | 56 |
| 189 | High injury incidence in adolescent female soccer. <i>American Journal of Sports Medicine</i> , <b>2014</b> , 42, 2487-94                                                                                                                       | 6.8  | 53 |
| 188 | Athletic groin pain: a systematic review of surgical diagnoses, investigations and treatment. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 1181-6                                                                              | 10.3 | 52 |
| 187 | Large eccentric strength increase using the Copenhagen Adduction exercise in football: A randomized controlled trial. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2016</b> , 26, 1334-1342                               | 4.6  | 52 |

|     |                                                                                                                                                                                                                                                                                             |      |    |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 186 | The diagnostic and prognostic value of ultrasonography in soccer players with acute hamstring injuries. <i>American Journal of Sports Medicine</i> , <b>2014</b> , 42, 399-404                                                                                                              | 6.8  | 51 |
| 185 | Associations between abnormal ultrasound color Doppler measures and tendon pain symptoms in badminton players during a season: a prospective cohort study. <i>American Journal of Sports Medicine</i> , <b>2012</b> , 40, 548-55                                                            | 6.8  | 51 |
| 184 | Danish Hip Arthroscopy Registry (DHAR): the outcome of patients with femoroacetabular impingement (FAI). <i>Journal of Hip Preservation Surgery</i> , <b>2017</b> , 4, 170-177                                                                                                              | 2    | 50 |
| 183 | Acute hamstring injuries in Danish elite football: a 12-month prospective registration study among 374 players. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2010</b> , 20, 588-92                                                                                    | 4.6  | 50 |
| 182 | Dislocation following total hip arthroplasty. <i>Archives of Orthopaedic and Traumatic Surgery Archiv für Orthopädische Und Unfall-Chirurgie</i> , <b>1985</b> , 103, 375-7                                                                                                                 |      | 50 |
| 181 | Effects of the Nordic Hamstring exercise on sprint capacity in male football players: a randomized controlled trial. <i>Journal of Sports Sciences</i> , <b>2018</b> , 36, 1663-1672                                                                                                        | 3.6  | 47 |
| 180 | Eccentric strengthening effect of hip-adductor training with elastic bands in soccer players: a randomised controlled trial. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 332-8                                                                                            | 10.3 | 46 |
| 179 | MRI findings in soccer players with long-standing adductor-related groin pain and asymptomatic controls. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 681-91                                                                                                               | 10.3 | 45 |
| 178 | Patellofemoral Pain in Adolescence and Adulthood: Same Same, but Different?. <i>Sports Medicine</i> , <b>2015</b> , 45, 1489-95                                                                                                                                                             | 10.6 | 40 |
| 177 | Consensus recommendations on the classification, definition and diagnostic criteria of hip-related pain in young and middle-aged active adults from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 631-641 | 10.3 | 39 |
| 176 | Hip strength and range of motion: Normal values from a professional football league. <i>Journal of Science and Medicine in Sport</i> , <b>2017</b> , 20, 339-343                                                                                                                            | 4.4  | 39 |
| 175 | Symptoms of nerve dysfunction after hip arthroscopy: an under-reported complication?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , <b>2014</b> , 30, 202-7                                                                                                            | 5.4  | 39 |
| 174 | Reproducibility of ultrasound and magnetic resonance imaging measurements of tendon size. <i>Acta Radiologica</i> , <b>2006</b> , 47, 954-9                                                                                                                                                 | 2    | 38 |
| 173 | Copenhagen five-second squeeze: a valid indicator of sports-related hip and groin function. <i>British Journal of Sports Medicine</i> , <b>2017</b> , 51, 594-599                                                                                                                           | 10.3 | 36 |
| 172 | Patient-reported outcome measures for hip-related pain: a review of the available evidence and a consensus statement from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 848-857                           | 10.3 | 36 |
| 171 | Musculoskeletal Screening Tests and Bony Hip Morphology Cannot Identify Male Professional Soccer Players at Risk of Groin Injuries: A 2-Year Prospective Cohort Study. <i>American Journal of Sports Medicine</i> , <b>2018</b> , 46, 1294-1305                                             | 6.8  | 36 |
| 170 | Characteristics of acute groin injuries in the adductor muscles: A detailed MRI study in athletes. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2018</b> , 28, 667-676                                                                                                | 4.6  | 36 |
| 169 | Danish Hip Arthroscopy Registry: an epidemiologic and perioperative description of the first 2000 procedures. <i>Journal of Hip Preservation Surgery</i> , <b>2016</b> , 3, 138-45                                                                                                          | 2    | 35 |

|     |                                                                                                                                                                                                                                                                                            |      |    |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 168 | Ultrasound guided electrocoagulation in patients with chronic non-insertional Achilles tendinopathy: a pilot study. <i>British Journal of Sports Medicine</i> , <b>2006</b> , 40, 761-6                                                                                                    | 10.3 | 35 |
| 167 | Quality assessment of radiological measurements of trochlear dysplasia; a literature review. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2018</b> , 26, 746-755                                                                                                             | 5.5  | 34 |
| 166 | Including the Copenhagen Adduction Exercise in the FIFA 11+ Provides Missing Eccentric Hip Adduction Strength Effect in Male Soccer Players: A Randomized Controlled Trial. <i>American Journal of Sports Medicine</i> , <b>2017</b> , 45, 3052-3059                                       | 6.8  | 34 |
| 165 | Ethnic Differences in Bony Hip Morphology in a Cohort of 445 Professional Male Soccer Players. <i>American Journal of Sports Medicine</i> , <b>2016</b> , 44, 2967-2974                                                                                                                    | 6.8  | 34 |
| 164 | Surgical criteria for femoroacetabular impingement syndrome: a scoping review. <i>British Journal of Sports Medicine</i> , <b>2017</b> , 51, 1605-1610                                                                                                                                     | 10.3 | 33 |
| 163 | The effect of protein and carbohydrate supplementation on strength training outcome of rehabilitation in ACL patients. <i>Journal of Orthopaedic Research</i> , <b>2006</b> , 24, 2114-23                                                                                                  | 3.8  | 33 |
| 162 | Minimum reporting standards for clinical research on groin pain in athletes. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 775-81                                                                                                                                          | 10.3 | 32 |
| 161 | Clinical Examination, Diagnostic Imaging, and Testing of Athletes With Groin Pain: An Evidence-Based Approach to Effective Management. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2018</b> , 48, 239-249                                                               | 4.2  | 32 |
| 160 | Patient-Reported Outcomes Within the First Year After Hip Arthroscopy and Rehabilitation for Femoroacetabular Impingement and/or Labral Injury: The Difference Between Getting Better and Getting Back to Normal. <i>American Journal of Sports Medicine</i> , <b>2018</b> , 46, 2607-2614 | 6.8  | 32 |
| 159 | Continued significant effect of physical training as treatment for overuse injury: 8- to 12-year outcome of a randomized clinical trial. <i>American Journal of Sports Medicine</i> , <b>2011</b> , 39, 2447-51                                                                            | 6.8  | 32 |
| 158 | Diagnostic Accuracy of Imaging Modalities and Injection Techniques for the Diagnosis of Femoroacetabular Impingement/Labral Tear: A Systematic Review With Meta-analysis. <i>American Journal of Sports Medicine</i> , <b>2017</b> , 45, 2665-2677                                         | 6.8  | 31 |
| 157 | The pyramidalis-anterior pubic ligament-adductor longus complex (PLAC) and its role with adductor injuries: a new anatomical concept. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2017</b> , 25, 3969-3977                                                                  | 5.5  | 31 |
| 156 | Micromechanical properties and collagen composition of ruptured human achilles tendon. <i>American Journal of Sports Medicine</i> , <b>2013</b> , 41, 437-43                                                                                                                               | 6.8  | 30 |
| 155 | Can standardised clinical examination of athletes with acute groin injuries predict the presence and location of MRI findings?. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 1541-1547                                                                                    | 10.3 | 29 |
| 154 | Sprint and jump performance in elite male soccer players following a 10-week Nordic Hamstring exercise Protocol: a randomised pilot study. <i>BMC Research Notes</i> , <b>2017</b> , 10, 669                                                                                               | 2.3  | 28 |
| 153 | Pubic apophysitis: a previously undescribed clinical entity of groin pain in athletes. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 828-34                                                                                                                                | 10.3 | 27 |
| 152 | Glenohumeral and scapulothoracic strength impairments exists in patients with subacromial impingement, but these are not reflected in the shoulder pain and disability index. <i>BMC Musculoskeletal Disorders</i> , <b>2017</b> , 18, 302                                                 | 2.8  | 27 |
| 151 | Isolated ulnar shaft fractures. Comparison of treatment by a functional brace and long-arm cast. <i>Journal of Bone and Joint Surgery: British Volume</i> , <b>1992</b> , 74, 757-9                                                                                                        |      | 27 |

|     |                                                                                                                                                                                                                                                                                |      |    |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 150 | Danish Hip Arthroscopy Registry: predictors of outcome in patients with femoroacetabular impingement (FAI). <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2019</b> , 27, 3110-3120                                                                                | 5.5  | 27 |
| 149 | Reliability of MRI assessment of acute musculotendinous groin injuries in athletes. <i>European Radiology</i> , <b>2017</b> , 27, 1486-1495                                                                                                                                    | 8    | 26 |
| 148 | Hip arthroscopy with labral repair for femoroacetabular impingement: short-term outcomes. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2014</b> , 22, 744-9                                                                                                      | 5.5  | 26 |
| 147 | Return to Sport After Criteria-Based Rehabilitation of Acute Adductor Injuries in Male Athletes: A Prospective Cohort Study. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2020</b> , 8, 2325967119897247                                                                 | 3.5  | 23 |
| 146 | The Influence of Hamstring Muscle Peak Torque and Rate of Torque Development for Sprinting Performance in Football Players: A Cross-Sectional Study. <i>International Journal of Sports Physiology and Performance</i> , <b>2019</b> , 14, 665-673                             | 3.5  | 23 |
| 145 | Hip joint pathology: relationship between patient history, physical tests, and arthroscopy findings in clinical practice. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2017</b> , 27, 342-350                                                            | 4.6  | 22 |
| 144 | Adductor canal blockade for moderate to severe pain after arthroscopic knee surgery: a randomized controlled trial. <i>Acta Anaesthesiologica Scandinavica</i> , <b>2014</b> , 58, 1220-7                                                                                      | 1.9  | 22 |
| 143 | Hip abduction strength training in the clinical setting: with or without external loading?. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2010</b> , 20 Suppl 2, 70-7                                                                                     | 4.6  | 22 |
| 142 | Acute patellofemoral pain: aggravating activities, clinical examination, MRI and ultrasound findings. <i>British Journal of Sports Medicine</i> , <b>2008</b> , 42, 64-7; discussion 67                                                                                        | 10.3 | 22 |
| 141 | The elite marathon runner: problems during and after competition. <i>British Journal of Sports Medicine</i> , <b>1988</b> , 22, 19-21                                                                                                                                          | 10.3 | 21 |
| 140 | Important clinical descriptors to include in the examination and assessment of patients with femoroacetabular impingement syndrome: an international and multi-disciplinary Delphi survey. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2017</b> , 25, 1975-1986 | 5.5  | 19 |
| 139 | Multidisciplinary Assessment of 100 Athletes With Groin Pain Using the Doha Agreement: High Prevalence of Adductor-Related Groin Pain in Conjunction With Multiple Causes. <i>Clinical Journal of Sport Medicine</i> , <b>2018</b> , 28, 364-369                               | 3.2  | 19 |
| 138 | Does bony hip morphology affect the outcome of treatment for patients with adductor-related groin pain? Outcome 10 years after baseline assessment. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 1240-4                                                       | 10.3 | 19 |
| 137 | Terminology and definitions on groin pain in athletes: building agreement using a short Delphi method. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 825-7                                                                                                     | 10.3 | 18 |
| 136 | Muscle-tendon-related pain in 100 patients with hip dysplasia: prevalence and associations with self-reported hip disability and muscle strength. <i>Journal of Hip Preservation Surgery</i> , <b>2018</b> , 5, 39-46                                                          | 2    | 18 |
| 135 | Preseason Adductor Squeeze Strength in 303 Spanish Male Soccer Athletes: A Cross-sectional Study. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2018</b> , 6, 2325967117747275                                                                                            | 3.5  | 18 |
| 134 | Femoroacetabular Impingement Surgery Is on the Rise-But What Is the Next Step?. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2016</b> , 46, 406-8                                                                                                            | 4.2  | 18 |
| 133 | The Copenhagen Standardised MRI protocol to assess the pubic symphysis and adductor regions of athletes: outline and intratester and intertester reliability. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 692-9                                              | 10.3 | 18 |

|     |                                                                                                                                                                                                                                                                                                  |      |    |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 132 | Physiotherapist-led treatment for young to middle-aged active adults with hip-related pain: consensus recommendations from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 504-511                               | 10.3 | 18 |
| 131 | Activity Modification and Load Management of Adolescents With Patellofemoral Pain: A Prospective Intervention Study Including 151 Adolescents. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, 1629-1637                                                                          | 6.8  | 17 |
| 130 | Demographic and Radiographic Factors Associated With Intra-articular Hip Cartilage Injury: A Cross-sectional Study of 1511 Hip Arthroscopy Procedures. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, 2617-2625                                                                  | 6.8  | 17 |
| 129 | Prevalence and severity of groin problems in Spanish football: A prospective study beyond the time-loss approach. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2020</b> , 30, 914-921                                                                                      | 4.6  | 17 |
| 128 | Characteristics of acute groin injuries in the hip flexor muscles - a detailed MRI study in athletes. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2018</b> , 28, 677-685                                                                                                  | 4.6  | 16 |
| 127 | Efficacy of early controlled motion of the ankle compared with immobilisation in non-operative treatment of patients with an acute Achilles tendon rupture: an assessor-blinded, randomised controlled trial. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 719-724              | 10.3 | 16 |
| 126 | Large strengthening effect of a hip-flexor training programme: a randomized controlled trial. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2016</b> , 24, 2346-52                                                                                                                  | 5.5  | 15 |
| 125 | Iliotibial band autograft versus bone-patella-tendon-bone autograft, a possible alternative for ACL reconstruction: a 15-year prospective randomized controlled trial. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2014</b> , 22, 2094-101                                        | 5.5  | 15 |
| 124 | Non-elite marathon runners: health, training and injuries. <i>British Journal of Sports Medicine</i> , <b>1989</b> , 23, 177-8                                                                                                                                                                   | 10.3 | 15 |
| 123 | Ultrasound Doppler of the Achilles tendon before and after injection of an ultrasound contrast agent--findings in asymptomatic subjects. <i>Ultraschall in Der Medizin</i> , <b>2007</b> , 28, 52-6                                                                                              | 3.8  | 13 |
| 122 | Is Bony Hip Morphology Associated With Range of Motion and Strength in Asymptomatic Male Soccer Players?. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2018</b> , 48, 250-259                                                                                                  | 4.2  | 12 |
| 121 | Different injury pattern in goalkeepers compared to field players: a three-year epidemiological study of professional football. <i>Journal of Science and Medicine in Sport</i> , <b>2014</b> , 17, 34-8                                                                                         | 4.4  | 12 |
| 120 | Clinical recovery of two hip adductor longus ruptures: a case-report of a soccer player. <i>BMC Research Notes</i> , <b>2013</b> , 6, 205                                                                                                                                                        | 2.3  | 12 |
| 119 | Plaster cast compared with bridging external fixation for distal radius fractures of the Colles type. <i>International Orthopaedics</i> , <b>2001</b> , 24, 358-60                                                                                                                               | 3.8  | 12 |
| 118 | DYNAMIC HIP ADDUCTION, ABDUCTION AND ABDOMINAL EXERCISES FROM THE HOLMICH GROIN-INJURY PREVENTION PROGRAM ARE INTENSE ENOUGH TO BE CONSIDERED STRENGTHENING EXERCISES - A CROSS-SECTIONAL STUDY. <i>International Journal of Sports Physical Therapy</i> , <b>2017</b> , 12, 371-380             | 1.4  | 12 |
| 117 | Pain, Sports Participation, and Physical Function in Adolescents With Patellofemoral Pain and Osgood-Schlatter Disease: A Matched Cross-sectional Study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2020</b> , 50, 149-157                                                   | 4.2  | 12 |
| 116 | Standardised measurement of physical capacity in young and middle-aged active adults with hip-related pain: recommendations from the first International Hip-related Pain Research Network (IHiPRN) meeting, Zurich, 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 702-710 | 10.3 | 12 |
| 115 | Multicentre study on capsular closure versus non-capsular closure during hip arthroscopy in Danish patients with femoroacetabular impingement (FAI): protocol for a randomised controlled trial. <i>BMJ Open</i> , <b>2018</b> , 8, e019176                                                      | 3    | 12 |

|     |                                                                                                                                                                                                                                                                                                                                                                                |      |    |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 114 | The association between specific sports activities and sport performance following hip arthroscopy for femoroacetabular impingement syndrome: A secondary analysis of a cross-sectional cohort study including 184 athletes. <i>Journal of Hip Preservation Surgery</i> , <b>2019</b> , 6, 124-133                                                                             | 2    | 11 |
| 113 | The iHOT-33: how valid is it?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , <b>2012</b> , 28, 1194-5; author reply 1195-6                                                                                                                                                                                                                                | 5.4  | 11 |
| 112 | Concussion surveillance: do low concussion rates in the Qatar Professional Football League reflect a true difference or emphasize challenges in knowledge translation?. <i>Clinical Journal of Sport Medicine</i> , <b>2015</b> , 25, 73-4                                                                                                                                     | 3.2  | 10 |
| 111 | Associations Between Initial Clinical Examination and Imaging Findings and Return-to-Sport in Male Athletes With Acute Adductor Injuries: A Prospective Cohort Study. <i>American Journal of Sports Medicine</i> , <b>2020</b> , 48, 1151-1159                                                                                                                                 | 6.8  | 10 |
| 110 | Muscle-tendon-related abnormalities detected by ultrasonography are common in symptomatic hip dysplasia. <i>Archives of Orthopaedic and Trauma Surgery</i> , <b>2018</b> , 138, 1059-1067                                                                                                                                                                                      | 3.6  | 10 |
| 109 | Patient-reported outcome and muscle-tendon pain after periacetabular osteotomy are related: 1-year follow-up in 82 patients with hip dysplasia. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2019</b> , 90, 40-45                                                                                                                                    | 4.3  | 10 |
| 108 | Study protocol for a randomised controlled trial of meniscal surgery compared with exercise and patient education for treatment of meniscal tears in young adults. <i>BMJ Open</i> , <b>2017</b> , 7, e017436                                                                                                                                                                  | 3    | 9  |
| 107 | Measurement qualities of hip and groin outcome scores: letter to the editor. <i>American Journal of Sports Medicine</i> , <b>2014</b> , 42, NP7-10                                                                                                                                                                                                                             | 6.8  | 9  |
| 106 | Effect of Medialization of the Trochlear Groove and Lateralization of the Tibial Tubercle on TT-TG Distance: A Cross-sectional Study of Dysplastic and Nondysplastic Knees. <i>American Journal of Sports Medicine</i> , <b>2021</b> , 49, 970-974                                                                                                                             | 6.8  | 9  |
| 105 | Does the physical activity profile change in patients with hip dysplasia from before to 1 year after periacetabular osteotomy?. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2018</b> , 89, 622-627                                                                                                                                                  | 4.3  | 9  |
| 104 | Increased external hip-rotation strength relates to reduced dynamic knee control in females: paradox or adaptation?. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2011</b> , 21, e215-21                                                                                                                                                                 | 4.6  | 8  |
| 103 | Insulin-dependent diabetes mellitus and marathon running. <i>British Journal of Sports Medicine</i> , <b>1987</b> , 21, 51-2                                                                                                                                                                                                                                                   | 10.3 | 8  |
| 102 | Efficacy of early controlled motion of the ankle compared with no motion after non-operative treatment of an acute Achilles tendon rupture: study protocol for a randomized controlled trial. <i>Trials</i> , <b>2016</b> , 17, 564                                                                                                                                            | 2.8  | 8  |
| 101 | Rasch validation of the Danish version of the shoulder pain and disability index (SPADI) in patients with rotator cuff-related disorders. <i>Quality of Life Research</i> , <b>2019</b> , 28, 795-800                                                                                                                                                                          | 3.7  | 8  |
| 100 | Conservative treatment for patients with subacromial impingement: Changes in clinical core outcomes and their relation to specific rehabilitation parameters. <i>PeerJ</i> , <b>2018</b> , 6, e4400                                                                                                                                                                            | 3.1  | 8  |
| 99  | Groin injuries in athletes--development of clinical entities, treatment, and prevention. <i>Danish Medical Journal</i> , <b>2015</b> , 62, B5184                                                                                                                                                                                                                               | 3.8  | 8  |
| 98  | Mechanistic pain profiling in young adolescents with patellofemoral pain before and after treatment: a prospective cohort study. <i>Pain</i> , <b>2020</b> , 161, 1065-1071                                                                                                                                                                                                    | 8    | 7  |
| 97  | The Strengthening Exercises in Shoulder Impingement trial (The SExSI-trial) investigating the effectiveness of a simple add-on shoulder strengthening exercise programme in patients with long-lasting subacromial impingement syndrome: Study protocol for a pragmatic, assessor blinded, parallel-group, randomised, controlled trial. <i>Trials</i> , <b>2019</b> , 19, 154 | 2.8  | 7  |



|    |                                                                                                                                                                                                                                                                                          |      |   |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 96 | Increased medial foot loading during drop jump in subjects with patellofemoral pain. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2014</b> , 22, 2301-7                                                                                                                    | 5.5  | 7 |
| 95 | Injury incidence and burden in a youth elite football academy: a four-season prospective study of 551 players aged from under 9 to under 19 years. <i>British Journal of Sports Medicine</i> , <b>2021</b> , 55, 493-500                                                                 | 10.3 | 7 |
| 94 | First-time anterior cruciate ligament injury in adolescent female elite athletes: a prospective cohort study to identify modifiable risk factors. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2021</b> , 1                                                                | 5.5  | 7 |
| 93 | Hip Strength Testing of Soccer Players With Long-Standing Hip and Groin Pain: What are the Clinical Implications of Pain During Testing?. <i>Clinical Journal of Sport Medicine</i> , <b>2016</b> , 26, 210-5                                                                            | 3.2  | 7 |
| 92 | Reliability of the Copenhagen Achilles length measure (CALM) on patients with an Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2020</b> , 28, 281-290                                                                                              | 5.5  | 7 |
| 91 | Activity Modification and Knee Strengthening for Osgood-Schlatter Disease: A Prospective Cohort Study. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2020</b> , 8, 2325967120911106                                                                                                 | 3.5  | 7 |
| 90 | Self-reported previous knee injury and low knee function increase knee injury risk in adolescent female football. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2016</b> , 26, 919-26                                                                               | 4.6  | 6 |
| 89 | Epidemiology of Groin Injuries in Athletes <b>2014</b> , 13-21                                                                                                                                                                                                                           |      | 6 |
| 88 | MEASURES OF HIP MUSCLE STRENGTH AND RATE OF FORCE DEVELOPMENT USING A FIXATED HANDHELD DYNAMOMETER: INTRA-TESTER INTRA-DAY RELIABILITY OF A CLINICAL SET-UP. <i>International Journal of Sports Physical Therapy</i> , <b>2019</b> , 14, 715-723                                         | 1.4  | 6 |
| 87 | How Many Patients Achieve an Acceptable Symptom State After Hip Arthroscopy for Femoroacetabular Impingement Syndrome? A Cross-sectional Study Including PASS Cutoff Values for the HAGOS and iHOT-33. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2021</b> , 9, 2325967121995267 | 3.5  | 6 |
| 86 | Custom-Made Foot Orthoses Decrease Medial Foot Loading During Drop Jump in Individuals With Patellofemoral Pain. <i>Clinical Journal of Sport Medicine</i> , <b>2016</b> , 26, 335-7                                                                                                     | 3.2  | 6 |
| 85 | Progression of Strength, Flexibility, and Palpation Pain During Rehabilitation of Athletes With Acute Adductor Injuries: A Prospective Cohort Study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2021</b> , 51, 126-134                                               | 4.2  | 6 |
| 84 | Groin injuries in athletes [New stepping stones. <i>Sports Orthopaedics and Traumatology</i> , <b>2017</b> , 33, 106-112].                                                                                                                                                               | 2.4  | 5 |
| 83 | Proximal hamstring tendon avulsion treatment choice depends on a combination of clinical and imaging-related factors: a worldwide survey on current clinical practice and decision-making. <i>Journal of ISAKOS</i> , <b>2019</b> , 4, 175-180                                           | 1.1  | 5 |
| 82 | Acute and sub-acute effects of repetitive kicking on hip adduction torque in injury-free elite youth soccer players. <i>Journal of Sports Sciences</i> , <b>2014</b> , 32, 1357-64                                                                                                       | 3.6  | 5 |
| 81 | Neuromuscular Coordination Deficit Persists 12 Months after ACL Reconstruction But Can Be Modulated by 6 Weeks of Kettlebell Training: A Case Study in Women's Elite Soccer. <i>Case Reports in Orthopedics</i> , <b>2017</b> , 2017, 4269575                                            | 0.4  | 5 |
| 80 | The use and handling of acrylic bone cement in Danish orthopaedic departments. <i>Basic and Clinical Pharmacology and Toxicology</i> , <b>1993</b> , 72, 332-5                                                                                                                           |      | 5 |
| 79 | A comparison of hamstring muscle activity during different screening tests for non-contact ACL injury. <i>Knee</i> , <b>2016</b> , 23, 362-6                                                                                                                                             | 2.6  | 5 |

|    |                                                                                                                                                                                                                                                                                                                            |     |   |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 78 | Men are stronger than women-also in the hip. <i>Journal of Science and Medicine in Sport</i> , <b>2013</b> , 16, E1-3                                                                                                                                                                                                      | 4.4 | 4 |
| 77 | Reliability and normative values of the foot line test: a technique to assess foot posture. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2007</b> , 37, 703-7                                                                                                                                            | 4.2 | 4 |
| 76 | Prevalence and Impact of Diabetes Mellitus on the Frozen Shoulder. <i>Southern Medical Journal</i> , <b>2018</b> , 111, 654-659                                                                                                                                                                                            | 0.6 | 4 |
| 75 | Ultrasound findings in adductor related groin pain. <i>Ultraschall in Der Medizin</i> , <b>2006</b> , 27, 509-11                                                                                                                                                                                                           | 3.8 | 4 |
| 74 | Completeness and data validity in the Danish Achilles tendon Database. <i>Danish Medical Journal</i> , <b>2019</b> , 66,                                                                                                                                                                                                   | 3.8 | 4 |
| 73 | Risk of Deep Vein Thrombosis After Acute Achilles Tendon Rupture: A Secondary Analysis of a Randomized Controlled Trial Comparing Early Controlled Motion of the Ankle Versus Immobilization. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2020</b> , 8, 2325967120915909                                            | 3.5 | 3 |
| 72 | Individualized treatment for acute Achilles tendon rupture based on the Copenhagen Achilles Rupture Treatment Algorithm (CARTA): a study protocol for a multicenter randomized controlled trial. <i>Trials</i> , <b>2020</b> , 21, 399                                                                                     | 2.8 | 3 |
| 71 | Patients who are candidates for subacromial decompression have more pronounced range of motion deficits, but do not differ in self-reported shoulder function, strength or pain compared to non-candidates. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2018</b> , 26, 2505-2511                            | 5.5 | 3 |
| 70 | Normative profiles for hip strength and flexibility in elite footballers. <i>Journal of Science and Medicine in Sport</i> , <b>2014</b> , 18, e32-e33                                                                                                                                                                      | 4.4 | 3 |
| 69 | MAXIMAL HIP AND KNEE MUSCLE STRENGTH ARE NOT RELATED TO NEUROMUSCULAR PRE-ACTIVITY DURING SIDECUTTING MANEUVER: A CROSS-SECTIONAL STUDY. <i>International Journal of Sports Physical Therapy</i> , <b>2018</b> , 13, 66-76                                                                                                 | 1.4 | 3 |
| 68 | Maximal hip muscle strength and rate of torque development 6-30 months after hip arthroscopy for femoroacetabular impingement syndrome: A cross-sectional study. <i>Journal of Science and Medicine in Sport</i> , <b>2021</b> , 24, 1110-1115                                                                             | 4.4 | 3 |
| 67 | Effectiveness of Adding a Large Dose of Shoulder Strengthening to Current Nonoperative Care for Subacromial Impingement: A Pragmatic, Double-Blind Randomized Controlled Trial (SESI Trial). <i>American Journal of Sports Medicine</i> , <b>2021</b> , 49, 3040-3049                                                      | 6.8 | 3 |
| 66 | Graft Failure, Revision ACLR, and Reoperation Rates After ACLR With Quadriceps Tendon Versus Hamstring Tendon Autografts: A Registry Study With Review of 475 Patients. <i>American Journal of Sports Medicine</i> , <b>2021</b> , 49, 2136-2143                                                                           | 6.8 | 3 |
| 65 | Isometric hip strength impairments in patients with hip dysplasia are improved but not normalized 1 year after periacetabular osteotomy: a cohort study of 82 patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2021</b> , 92, 285-291                                                      | 4.3 | 3 |
| 64 | Is the Prognosis of Osgood-Schlatter Poorer Than Anticipated? A Prospective Cohort Study With 24-Month Follow-up. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2021</b> , 9, 23259671211022239                                                                                                                       | 3.5 | 3 |
| 63 | Outcome after acute Achilles tendon rupture is not negatively affected by female sex and age over 65 years. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2020</b> , 28, 3994-4002                                                                                                                            | 5.5 | 2 |
| 62 | HAGOS Could Be Important in the Evaluation of Patients Undergoing Hip Arthroscopy-Why Ignore It in a Sports Medicine Update When the Scientific Data Suggests Otherwise? Letter to the Editor. <i>American Journal of Sports Medicine</i> , <b>2018</b> , 46, NP6-NP7                                                      | 6.8 | 2 |
| 61 | Passive Knee Stability After Anterior Cruciate Ligament Reconstruction Using the Endobutton or ToggleLoc With ZipLoop as a Femoral Fixation Device: A Comparison of 1654 Patients From the Danish Knee Ligament Reconstruction Registry. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2018</b> , 6, 2325967118778507 | 3.5 | 2 |

|    |                                                                                                                                                                                                                                                                                             |      |   |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 60 | Electromyography Activation Levels of the 3 Gluteus Medius Subdivisions During Manual Strength Testing. <i>Journal of Sport Rehabilitation</i> , <b>2015</b> , 24, 244-51                                                                                                                   | 1.7  | 2 |
| 59 | Preventing Groin Injuries 91-113                                                                                                                                                                                                                                                            |      | 2 |
| 58 | MEASURES OF HIP MUSCLE STRENGTH AND RATE OF FORCE DEVELOPMENT USING A FIXATED HANDHELD DYNAMOMETER: INTRA-TESTER INTRA-DAY RELIABILITY OF A CLINICAL SET-UP. <i>International Journal of Sports Physical Therapy</i> , <b>2019</b> , 14, 715-723                                            | 1.4  | 2 |
| 57 | Analgesic use in adolescents with patellofemoral pain or Osgood-Schlatter Disease: a secondary cross-sectional analysis of 323 subjects. <i>Scandinavian Journal of Pain</i> , <b>2021</b> ,                                                                                                | 1.9  | 2 |
| 56 | An Updated Description of More Than 5,000 Procedures from the Danish Hip Arthroscopy Registry. <i>Journal of Bone and Joint Surgery - Series A</i> , <b>2020</b> , 102, 43-50                                                                                                               | 5.6  | 2 |
| 55 | SPRINT PERFORMANCE IN FOOTBALL (SOCCER) PLAYERS WITH AND WITHOUT A PREVIOUS HAMSTRING STRAIN INJURY: AN EXPLORATIVE CROSS-SECTIONAL STUDY. <i>International Journal of Sports Physical Therapy</i> , <b>2020</b> , 15, 947-957                                                              | 1.4  | 2 |
| 54 | The Achilles Tendon Length Measure and the Achilles Tendon Resting Angle show acceptable construct validity using the Copenhagen Achilles Length Measure as gold standard. <i>Foot and Ankle Surgery</i> , <b>2021</b> , 27, 655-659                                                        | 3.1  | 2 |
| 53 | Association of Skeletal Maturity and Injury Risk in Elite Youth Soccer Players: A 4-Season Prospective Study With Survival Analysis. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2021</b> , 9, 2325967121999113                                                                      | 3.5  | 2 |
| 52 | 12 Efficacy of early controlled motion of the ankle in non-operative treatment of acute achilles tendon rupture. An assessor-blinded RCT <b>2019</b> ,                                                                                                                                      |      | 2 |
| 51 | Infographic. The Adductor Strengthening Programme prevents groin problems among male football players. <i>British Journal of Sports Medicine</i> , <b>2019</b> , 53, 45-46                                                                                                                  | 10.3 | 2 |
| 50 | Diabetes and treatment with orally administrated corticosteroids negatively affect treatment outcome at follow-up after acute Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2021</b> , 29, 1584-1592                                                  | 5.5  | 2 |
| 49 | Associations between clinical findings and MRI injury extent in male athletes with acute adductor injuries - A cross-sectional study. <i>Journal of Science and Medicine in Sport</i> , <b>2021</b> , 24, 454-462                                                                           | 4.4  | 2 |
| 48 | The heel-rise work test overestimates the performed work with 21-25% after an Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2021</b> , 29, 1604-1611                                                                                                  | 5.5  | 2 |
| 47 | No clinically relevant difference between operative and non-operative treatment in tendon elongation measured with the Achilles tendon resting angle (ATRA) 1 year after acute Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2021</b> , 29, 1617-1626 | 5.5  | 2 |
| 46 | Hamstring and Quadriceps Muscle Strength in Youth to Senior Elite Soccer: A Cross-Sectional Study Including 125 Players. <i>International Journal of Sports Physiology and Performance</i> , <b>2021</b> , 1-7                                                                              | 3.5  | 2 |
| 45 | Groin problems from pre- to in-season: a prospective study on 386 male Spanish footballers. <i>Research in Sports Medicine</i> , <b>2021</b> , 29, 498-504                                                                                                                                  | 3.8  | 1 |
| 44 | Effects of a lighter, smaller football on acute match injuries in adolescent female football: a pilot cluster-randomized controlled trial. <i>Journal of Sports Medicine and Physical Fitness</i> , <b>2018</b> , 58, 644-650                                                               | 1.4  | 1 |
| 43 | SPECIFIC EXERCISE TARGETING THE SEMITENDINOSUS IN FEMALE ACL-RECONSTRUCTED ATHLETES. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 623.1-623                                                                                                                                | 10.3 | 1 |

|    |                                                                                                                                                                                                                                                                                                          |      |   |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 42 | Renal laceration secondary to blunt trauma in soccer. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2007</b> , 3, 292-293                                                                                                                                                           | 4.6  | 1 |
| 41 | Hip adduction strength and provoked groin pain: A comparison of long-lever squeeze testing using the ForceFrame and the Copenhagen 5-Second-Squeeze test.. <i>Physical Therapy in Sport</i> , <b>2022</b> , 55, 28-36                                                                                    | 3    | 1 |
| 40 | The 45-second anterior knee pain provocation test: A quick test of knee pain and sporting function in 10-14-year-old adolescents with patellofemoral pain. <i>Physical Therapy in Sport</i> , <b>2021</b> , 53, 28-33                                                                                    | 3    | 1 |
| 39 | MAXIMAL HIP AND KNEE MUSCLE STRENGTH ARE NOT RELATED TO NEUROMUSCULAR PRE-ACTIVITY DURING SIDECUTTING MANEUVER: A CROSS-SECTIONAL STUDY. <i>International Journal of Sports Physical Therapy</i> , <b>2018</b> , 13, 66-76                                                                               | 1.4  | 1 |
| 38 | Neither heel-rise Height (HRH) nor Achilles tendon resting angle (ATRA) show strong correlations to patient limitations and return to previous activities one year after acute Achilles tendon rupture. <i>Foot and Ankle Surgery</i> , <b>2021</b> ,                                                    | 3.1  | 1 |
| 37 | Validation of the Copenhagen Hip and Groin Outcome Score (HAGOS) using modern test theory across different cultures and languages: a cross-sectional study of 452 male athletes with groin pain. <i>British Journal of Sports Medicine</i> , <b>2021</b> ,                                               | 10.3 | 1 |
| 36 | Capsular closure in patients with femoroacetabular impingement syndrome (FAIS): results of a matched-cohort study from the Danish hip arthroscopy registry. <i>Journal of Hip Preservation Surgery</i> , <b>2021</b> , 7, 474-482                                                                        | 2    | 1 |
| 35 | Can't See the Right Forest Plot for the Wrong Trees!. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , <b>2020</b> , 36, 2787-2789                                                                                                                                                     | 5.4  | 1 |
| 34 | Past-season, pre-season and in-season risk assessment of groin problems in male football players: a prospective full-season study. <i>British Journal of Sports Medicine</i> , <b>2021</b> ,                                                                                                             | 10.3 | 1 |
| 33 | One-Year Clinical and Imaging Follow-up After Exercise-Based Treatment for Acute Complete Adductor Longus Tendon Avulsions in Athletes: A Prospective Case Series. <i>American Journal of Sports Medicine</i> , <b>2021</b> , 49, 3004-3013                                                              | 6.8  | 1 |
| 32 | Infographic. Consensus recommendations on the classification, definition and diagnostic criteria of hip-related pain in young and middle-aged active adults from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2021</b> , 55, 115-117 | 10.3 | 1 |
| 31 | A Delphi survey and international e-survey evaluating the Doha agreement meeting classification system in groin pain: Where are we 5 years later?. <i>Journal of Science and Medicine in Sport</i> , <b>2022</b> , 25, 3-8                                                                               | 4.4  | 1 |
| 30 | Femoroacetabular impingement syndrome and labral injuries: grading the evidence on diagnosis and non-operative treatment-a statement paper commissioned by the Danish Society of Sports Physical Therapy (DSSF). <i>British Journal of Sports Medicine</i> , <b>2021</b> , 55, 1301-1310                 | 10.3 | 1 |
| 29 | Current Status of Trans-Arterial Embolization in Pain Management of Musculoskeletal Inflammatory Conditions - An Evidence-Based Review. <i>CardioVascular and Interventional Radiology</i> , <b>2021</b> , 44, 1699-1708                                                                                 | 2.7  | 1 |
| 28 | Anatomical retraction of the semitendinosus muscle following harvest of the distal semitendinosus tendon for ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2021</b> , 1                                                                                                 | 5.5  | 1 |
| 27 | Rapid Spike in Hip Adduction Strength in Early Adolescent Footballers: A Study of 125 Elite Male Players From Youth to Senior.. <i>International Journal of Sports Physiology and Performance</i> , <b>2022</b> , 1-8                                                                                    | 3.5  | 1 |
| 26 | Return to sports after femoroacetabular Impingement-Operation. <i>Arthroscopie</i> , <b>2016</b> , 29, 22-31                                                                                                                                                                                             | 0.1  | 0 |
| 25 | THE INFLUENCE OF HIP JOINT CONTROL ON KNEE JOINT VALGUS MOMENT IN YOUNG FEMALE ELITE ATHLETES. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 566.1-566                                                                                                                                   | 10.3 | 0 |

|    |                                                                                                                                                                                                                                                                                       |      |   |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 24 | Proximal hamstring tendon avulsion: state of the art. <i>Journal of ISAKOS</i> , <b>2021</b> , 6, 237-246                                                                                                                                                                             | 1.1  | 0 |
| 23 | Treatment of osteoarthritis with autologous, micro-fragmented adipose tissue: a study protocol for a randomized controlled trial. <i>Trials</i> , <b>2021</b> , 22, 748                                                                                                               | 2.8  | 0 |
| 22 | Arthroscopic treatment for femoroacetabular impingement syndrome (FAIS) in adolescents-5-year follow-up.. <i>Journal of Hip Preservation Surgery</i> , <b>2021</b> , 8, 249-254                                                                                                       | 2    | 0 |
| 21 | Classifying radiographic changes of the pubic symphysis in male athletes: Development and reproducibility of a new scoring protocol. <i>European Journal of Radiology</i> , <b>2021</b> , 134, 109452                                                                                 | 4.7  | 0 |
| 20 | Effectiveness of Adding a Large Dose of Shoulder Strengthening to Current Nonoperative Care for Subacromial Impingement: A Pragmatic, Double-Blind Randomized Controlled Trial (SExSI Trial): Response.. <i>American Journal of Sports Medicine</i> , <b>2022</b> , 50, NP20-NP23     | 6.8  | 0 |
| 19 | INCLUDING THE COPENHAGEN ADDUCTION EXERCISE IN THE FIFA 11+ PROVIDES MISSING ECCENTRIC HIP ADDUCTION STRENGTH EFFECT: A RANDOMISED CONTROLLED TRIAL. <i>British Journal of Sports Medicine</i> , <b>2017</b> , 51, 327.1-327                                                          | 10.3 |   |
| 18 | Clinical Examination and Diagnosis of Extra-Articular Hip and Groin Pain <b>2019</b> , 79-94                                                                                                                                                                                          |      |   |
| 17 | Muscular Function and Treatment of Musculotendinous Groin Pain <b>2019</b> , 95-105                                                                                                                                                                                                   |      |   |
| 16 | Groin Injuries <b>2019</b> , 223-231                                                                                                                                                                                                                                                  |      |   |
| 15 | Use of Sports Hernia to Describe Groin Pain in Athletes. <i>JAMA Surgery</i> , <b>2020</b> , 155, 895                                                                                                                                                                                 | 5.4  |   |
| 14 | Infographic. Can standardised clinical examination of athletes with acute groin injuries predict the presence and location of MRI findings?. <i>British Journal of Sports Medicine</i> , <b>2018</b> , 52, 892-893                                                                    | 10.3 |   |
| 13 | Hip, Groin, and Abdominal Injuries in Handball <b>2018</b> , 243-260                                                                                                                                                                                                                  |      |   |
| 12 | Fractures in sports. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2007</b> , 2, 44-47                                                                                                                                                                           | 4.6  |   |
| 11 | Ischial apophysiolysis: a case report and a short review. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2007</b> , 3, 131-133                                                                                                                                    | 4.6  |   |
| 10 | Silent ischaemia and vigorous exercise. <i>British Journal of Sports Medicine</i> , <b>1989</b> , 23, 133-4                                                                                                                                                                           | 10.3 |   |
| 9  | Acute and Long-Standing Groin Injuries <b>2022</b> , 113-122                                                                                                                                                                                                                          |      |   |
| 8  | Correspondence: Isometric hip strength impairments in patients with hip dysplasia are improved but not normalized 1 year after periacetabular osteotomy: a cohort study of 82 patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2021</b> , 92, 760-762 | 4.3  |   |
| 7  | Injuries in the Pelvis, Groin, Hip and Thigh <b>2015</b> , 551-562                                                                                                                                                                                                                    |      |   |

6 Groin Pain in Athletes: Assessment and Nonsurgical Treatment **2017**, 315-322

|   |                                                                                                                                                                                                                                                                                                         |      |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 5 | Pubic apophysitis: 6 questions that need answers before I'm convinced it's a new clinical condition.<br><i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 1421-1422                                                                                                                         | 10.3 |
| 4 | Now you see it - Now you don't: A letter to the editor concerning "Surgery is no more effective than conservative treatment for femoroacetabular impingement syndrome: Systematic review and meta-analysis of randomized controlled trials". <i>Clinical Rehabilitation</i> , <b>2021</b> , 35, 464-466 | 3.3  |
| 3 | Six weeks of intensive rehearsals for the Swan Lake ballet shows ultrasound tissue characterization changes of the Achilles tendons in dancers. <i>Scandinavian Journal of Medicine and Science in Sports</i> , <b>2021</b> , 31, 2133-2143                                                             | 4.6  |
| 2 | Cross-sectional Study of EMG and EMG Rise During Fast and Slow Hamstring Exercises.<br><i>International Journal of Sports Physical Therapy</i> , <b>2021</b> , 16, 1033-1042                                                                                                                            | 1.4  |
| 1 | Training and injuries among world elite junior badminton players - Identifying the problems.<br><i>Asia-Pacific Journal of Sports Medicine, Arthroscopy, Rehabilitation and Technology</i> , <b>2021</b> , 26, 21-26                                                                                    | 1.2  |