Antonio RodrÃ-guez-Poncelas

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8186831/publications.pdf

Version: 2024-02-01

20 papers

403 citations

933447 10 h-index 752698 20 g-index

25 all docs

25 docs citations

25 times ranked

676 citing authors

| # | Article | IF | CITATIONS |
|----|--|-----------|---------------|
| 1 | Analysis of the location of retinal lesions in central retinographies of patients with Type 2 diabetes. Acta Ophthalmologica, 2020, 98, e13-e21. | 1.1 | 7 |
| 2 | What is the optimal cut-off threshold in self-home blood pressure measurement?. Medicine (United) Tj ETQq0 C | 0 rgBT /0 | verlock 10 Tf |
| 3 | The dynamic behaviour of metabolic syndrome and its components in an eight-year population-based cohort from the Mediterranean. PLoS ONE, 2017, 12, e0176665. | 2.5 | 15 |
| 4 | Noninsulin Antidiabetic Drugs for Patients with Type 2 Diabetes Mellitus: Are We Respecting Their Contraindications?. Journal of Diabetes Research, 2016, 2016, 1-9. | 2.3 | 14 |
| 5 | Chronic Kidney Disease and Diabetic Retinopathy in Patients with Type 2 Diabetes. PLoS ONE, 2016, 11, e0149448. | 2.5 | 63 |
| 6 | Comparison of different vascular risk engines in the identification of type 2 diabetes patients with high cardiovascular risk. BMC Cardiovascular Disorders, 2015, 15, 121. | 1.7 | 7 |
| 7 | Prevalence of diabetic retinopathy in individuals with type 2 diabetes who had recorded diabetic retinopathy from retinal photographs in Catalonia (Spain). British Journal of Ophthalmology, 2015, 99, 1628-1633. | 3.9 | 40 |
| 8 | Metabolic control and cardiovascular risk factors in type 2 diabetes mellitus patients according to diabetes duration. Family Practice, 2015, 32, 27-34. | 1.9 | 21 |
| 9 | Are There Heightâ€Dependent Differences in Subclinical Vascular Disease in Hypertensive Patients?. Journal of Clinical Hypertension, 2014, 16, 70-76. | 2.0 | 0 |
| 10 | Impact of chronic kidney disease on the prevalence of cardiovascular disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. BMC Nephrology, 2014, 15, 150. | 1.8 | 16 |
| 11 | Retinal arteriole-to-venule ratio changes and target organ disease evolution in newly diagnosed hypertensive patients at 1-year follow-up. Journal of the American Society of Hypertension, 2014, 8, 83-93. | 2.3 | 9 |
| 12 | Executive function impairment in diabetics. Alzheimer Realidades E Investigaci \tilde{A}^3 n En Demencia, 2014, , 37-42. | 0.1 | 0 |
| 13 | Prevalence of chronic kidney disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. BMC Nephrology, 2013, 14, 46. | 1.8 | 100 |
| 14 | Prevalence of chronic kidney disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. Primary Care Diabetes, 2013, 7, 74. | 1.8 | 0 |
| 15 | Why is cardiovascular risk stratification important in hypertensive patients?. Blood Pressure, 2012, 21, 182-190. | 1.5 | 4 |
| 16 | Chronic kidney disease in the type 2 diabetic patients: prevalence and associated variables in a random sample of 2642 patients of a Mediterranean area. BMC Nephrology, 2012, 13, 87. | 1.8 | 48 |
| 17 | Evolution of Target Organ Damage by Different Values of Self-Blood Pressure Measurement in Untreated Hypertensive Patients. American Journal of Hypertension, 2012, 25, 1256-63. | 2.0 | 9 |
| 18 | Isolated clinical hypertension diagnosis. Blood Pressure Monitoring, 2011, 16, 11-15. | 0.8 | 8 |

ANTONIO PODRÃCHEZ PONCEIAS

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Proportion of isolated clinical hypertension in primary care settings. Comparison of target organ damage in patients with isolated clinical hypertension and patients with sustained arterial hypertension. Blood Pressure, 2007, 16, 354-361. | 1.5 | O |
| 20 | Assessment of selfâ€monitoring of blood pressure in the diagnosis of isolated clinic hypertension. Blood Pressure, 2006, 15, 227-236. | 1.5 | 15 |