

Ho-Shik Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8196674/publications.pdf>

Version: 2024-02-01

24
papers

359
citations

933447

10
h-index

839539

18
g-index

25
all docs

25
docs citations

25
times ranked

621
citing authors

#	ARTICLE	IF	CITATIONS
1	Fabry disease exacerbates renal interstitial fibrosis after unilateral ureteral obstruction via impaired autophagy and enhanced apoptosis. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 208-219.	2.2	14
2	The Levels of Circulating MicroRNAs at 6-Hour Cardiac Arrest Can Predict 6-Month Poor Neurological Outcome. <i>Diagnostics</i> , 2021, 11, 1905.	2.6	3
3	Induction of p53-Dependent Apoptosis by Prostaglandin A2. <i>Biomolecules</i> , 2020, 10, 492.	4.0	19
4	Gene editing particle system as a therapeutic approach for drug-resistant colorectal cancer. <i>Nano Research</i> , 2020, 13, 1576-1585.	10.4	9
5	Inhibition of p300/CBP-Associated Factor Attenuates Renal Tubulointerstitial Fibrosis through Modulation of NF- κ B and Nrf2. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1554.	4.1	42
6	Empagliflozin Contributes to Polyuria via Regulation of Sodium Transporters and Water Channels in Diabetic Rat Kidneys. <i>Frontiers in Physiology</i> , 2019, 10, 271.	2.8	27
7	Dâ€™Pinitol alleviates cyclosporine Aâ€™induced renal tubulointerstitial fibrosis via activating Sirt1 and Nrf2 antioxidant pathways. <i>International Journal of Molecular Medicine</i> , 2018, 41, 1826-1834.	4.0	19
8	PGA2-induced expression of HO-1 is mediated by transcriptional upregulation of Nrf2. <i>Molecular and Cellular Toxicology</i> , 2018, 14, 391-398.	1.7	0
9	Thyocyteâ€™specific deletion of insulin and IGFâ€™1 receptors induces papillary thyroid carcinomaâ€™like lesions through EGFR pathway activation. <i>International Journal of Cancer</i> , 2018, 143, 2458-2469.	5.1	10
10	PGA2 induces the expression of HO-1 by activating p53 in HCT116 cells. <i>Molecular and Cellular Toxicology</i> , 2017, 13, 189-196.	1.7	2
11	Treatment combining aliskiren with paricalcitol is effective against progressive renal tubulointerstitial fibrosis via dual blockade of intrarenal renin. <i>PLoS ONE</i> , 2017, 12, e0181757.	2.5	17
12	Identification of a novel GLA mutation (Y88C) in a Korean family with Fabry nephropathy: a case report. <i>BMC Medical Genetics</i> , 2016, 17, 76.	2.1	4
13	Deletion of IGF-1 Receptors in Cardiomyocytes Attenuates Cardiac Aging in Male Mice. <i>Endocrinology</i> , 2016, 157, 336-345.	2.8	75
14	T-type calcium channel blocker attenuates unilateral ureteral obstruction-induced renal interstitial fibrosis by activating the Nrf2 antioxidant pathway. <i>American Journal of Translational Research (discontinued)</i> , 2016, 8, 4574-4585.	0.0	7
15	PGA2-induced HO-1 attenuates G2M arrest by modulating GADD45â€™ expression. <i>Molecular and Cellular Toxicology</i> , 2015, 11, 465-474.	1.7	8
16	Silencing of KIF14 interferes with cell cycle progression and cytokinesis by blocking the p27Kip1 ubiquitination pathway in hepatocellular carcinoma. <i>Experimental and Molecular Medicine</i> , 2014, 46, e97-e97.	7.7	45
17	Nutlin-3 induces BCL2A1 expression by activating ELK1 through the mitochondrial p53-ROS-ERK1/2 pathway. <i>International Journal of Oncology</i> , 2014, 45, 675-682.	3.3	16
18	Nutlin-3 induces HO-1 expression by activating JNK in a transcription-independent manner of p53. <i>International Journal of Oncology</i> , 2014, 44, 761-768.	3.3	11

#	ARTICLE	IF	CITATIONS
19	Anthocyanins of Black Soybean (cv. Cheongja 3) Induce Autophagy via AMPK Activation in U2OS cells. FASEB Journal, 2012, 26, 543.2.	0.5	0
20	Nutlin-3 activates MEK1/2-ERK1/2 pathway via p53-induced ROS accumulation. FASEB Journal, 2012, 26, 761.11.	0.5	0
21	Sodium nitroprusside induces autophagic cell death in glutathione-depleted osteoblasts. Molecular and Cellular Toxicology, 2010, 6, 41-49.	1.7	1
22	Prostaglandin A2 induces caspase-independent apoptosis in hepatocellular carcinoma cells. The Korean Journal of Hepatology, 2005, 11, 72-9.	1.5	1
23	Induction of apoptosis dependent on caspase activities and growth arrest in HL-60 cells by PGA2. Prostaglandins and Other Lipid Mediators, 2002, 70, 169-183.	1.9	14
24	Induction of apoptosis in human leukemia cells by 3-deazaadenosine is mediated by caspase-3-like activity. Experimental and Molecular Medicine, 2000, 32, 197-203.	7.7	15