

# Masayuki Iki

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8199962/masayuki-iki-publications-by-year.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

82  
papers

2,229  
citations

24  
h-index

45  
g-index

86  
ext. papers

2,571  
ext. citations

4.1  
avg, IF

4.2  
L-index

#	Paper	IF	Citations
82	Guideline adherence by physicians for management of glucocorticoid-induced osteoporosis in Japan: a nationwide health insurance claims database study.. <i>Osteoporosis International</i> , <b>2022</b> , 33, 1097	5.3	1
81	Combined results of three physical performance tests predict incident fracture independently of aBMD in community-dwelling elderly Japanese men: Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Bone</i> , <b>2022</b> , 154, 116240	4.7	
80	Delayed initiation of anti-osteoporosis medications increases subsequent hip and vertebral fractures in patients on long-term glucocorticoid therapy: A nationwide health insurance claims database study in Japan.. <i>Bone</i> , <b>2022</b> , 160, 116396	4.7	0
79	Associations between trunk-to-peripheral fat ratio and cardiometabolic risk factors in elderly Japanese men: baseline data from the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) study. <i>Environmental Health and Preventive Medicine</i> , <b>2021</b> , 26, 35	4.2	0
78	Determinants of bone health in elderly Japanese men: study design and key findings of the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) cohort study. <i>Environmental Health and Preventive Medicine</i> , <b>2021</b> , 26, 51	4.2	3
77	Predictive ability of novel volumetric and geometric indices derived from dual-energy X-ray absorptiometric images of the proximal femur for hip fracture compared with conventional areal bone mineral density: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2021</b> , 32, 2289-2299	5.3	2
76	Circulating osteocalcin levels were not significantly associated with the risk of incident type 2 diabetes mellitus in elderly Japanese men: The Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Bone</i> , <b>2021</b> , 147, 115912	4.7	0
75	Dietary calcium and vitamin K are associated with osteoporotic fracture risk in middle-aged and elderly Japanese women, but not men: the Murakami Cohort Study. <i>British Journal of Nutrition</i> , <b>2021</b> , 125, 319-328	3.6	1
74	Insufficient persistence of and adherence to osteoporosis pharmacotherapy in Japan. <i>Journal of Bone and Mineral Metabolism</i> , <b>2021</b> , 39, 501-509	2.9	3
73	Calcium Intake and Bone Mineral Acquisition during the Pubertal Growth Spurt: Three-Year Follow-Up of the Kitakata Kids Health Study in Japan. <i>Journal of Nutritional Science and Vitaminology</i> , <b>2020</b> , 66, 158-167	1.1	4
72	Insufficient increase in bone mineral density testing rates and pharmacotherapy after hip fracture in Japan. <i>Journal of Bone and Mineral Metabolism</i> , <b>2020</b> , 38, 589-596	2.9	6
71	Relationships between serum uric acid concentrations, uric acid lowering medications, and vertebral fracture in community-dwelling elderly Japanese men: Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Bone</i> , <b>2020</b> , 139, 115519	4.7	4
70	Total physical activity and risk of chronic low back and knee pain in middle-aged and elderly Japanese people: The Murakami cohort study. <i>European Journal of Pain</i> , <b>2020</b> , 24, 863-872	3.7	4
69	Decreased bone mineral density and osteoporotic fractures are associated with the development of echogenic plaques in the carotid arteries over a 10-year follow-up period: The Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Maturitas</i> , <b>2020</b> , 131, 40-47	5	2
68	Natto Intake is Inversely Associated with Osteoporotic Fracture Risk in Postmenopausal Japanese Women. <i>Journal of Nutrition</i> , <b>2020</b> , 150, 599-605	4.1	15
67	Response to "Increased risk of osteoporotic fractures in patients with gastric cancer and post-gastrectomy" by Lai SW, et al. <i>Bone</i> , <b>2020</b> , 132, 115213	4.7	
66	Trunk-to-peripheral fat ratio predicts a subsequent blood pressure in normal-weight pubertal boys: a 3-year follow-up of the Kitakata Kids Health Study. <i>Environmental Health and Preventive Medicine</i> , <b>2020</b> , 25, 41	4.2	2

65	Response to "Is grip strength an actual/significant predictor of areal bone mineral density by itself? Potential confounders and/or effect modifiers". <i>Journal of Bone and Mineral Metabolism</i> , <b>2020</b> , 38, 137-138		2.9	
64	Hyperglycemic status is associated with an elevated risk of osteoporotic fracture in community-dwelling elderly Japanese men: The Fujiwara-kyo osteoporosis risk in men (FORMEN) cohort study. <i>Bone</i> , <b>2019</b> , 121, 100-106	4.7		11
63	For making a declaration of countermeasures against the falling birth rate from the Japanese Society for Hygiene: summary of discussion in the working group on academic research strategy against an aging society with low birth rate. <i>Environmental Health and Preventive Medicine</i> , <b>2019</b> , 24, 14	4.2		14
62	Increased risk of osteoporotic fracture in community-dwelling elderly men 20 or more years after gastrectomy: The Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Bone</i> , <b>2019</b> , 127, 250-259	4.7		9
61	Association between hand-grip strength and site-specific risks of major osteoporotic fracture: Results from the Japanese Population-based Osteoporosis Cohort Study. <i>Maturitas</i> , <b>2019</b> , 130, 13-20	5		11
60	Low bone mineral density is associated with an elevated risk of developing increased arterial stiffness: A 10-year follow-up of Japanese women from the Japanese Population-based Osteoporosis (JPOS) cohort study. <i>Maturitas</i> , <b>2019</b> , 119, 39-45	5		5
59	Estimates of hip fracture incidence in Japan using the National Health Insurance Claim Database in 2012-2015. <i>Osteoporosis International</i> , <b>2019</b> , 30, 975-983	5.3		40
58	Does Trabecular Bone Score (TBS) improve the predictive ability of FRAX for major osteoporotic fractures according to the Japanese Population-Based Osteoporosis (JPOS) cohort study?. <i>Journal of Bone and Mineral Metabolism</i> , <b>2019</b> , 37, 161-170	2.9		8
57	Muscle strength is associated with bone health independently of muscle mass in postmenopausal women: the Japanese population-based osteoporosis study. <i>Journal of Bone and Mineral Metabolism</i> , <b>2019</b> , 37, 53-59	2.9		16
56	Ratio of Endogenous Secretory Receptor for Advanced Glycation End Products to Pentosidine Predicts Fractures in Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2018</b> , 103, 85-94	5.6		13
55	The Murakami Cohort Study of vitamin D for the prevention of musculoskeletal and other age-related diseases: a study protocol. <i>Environmental Health and Preventive Medicine</i> , <b>2018</b> , 23, 28	4.2		15
54	Epidemiological profiles of chronic low back and knee pain in middle-aged and elderly Japanese from the Murakami cohort. <i>Journal of Pain Research</i> , <b>2018</b> , 11, 3161-3169	2.9		11
53	Predicting bone mineral acquisition during puberty: data from a 3-year follow-up study in Hamamatsu, Japan. <i>Journal of Bone and Mineral Metabolism</i> , <b>2017</b> , 35, 185-191	2.9		4
52	Relative Importance of Central and Peripheral Adiposities on Cardiometabolic Variables in Females: A Japanese Population-Based Study. <i>Journal of Clinical Densitometry</i> , <b>2017</b> , 20, 58-65	3.5		3
51	Total 25-hydroxyvitamin D levels predict fracture risk: results from the 15-year follow-up of the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2017</b> , 28, 1903-1913	5.3		35
50	Hyperglycemia is associated with increased bone mineral density and decreased trabecular bone score in elderly Japanese men: The Fujiwara-kyo osteoporosis risk in men (FORMEN) study. <i>Bone</i> , <b>2017</b> , 105, 18-25	4.7		28
49	Incident fracture associated with increased risk of mortality even after adjusting for frailty status in elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Osteoporosis International</i> , <b>2017</b> , 28, 871-880	5.3		5
48	A Meta-Analysis of Trabecular Bone Score in Fracture Risk Prediction and Its Relationship to FRAX. <i>Journal of Bone and Mineral Research</i> , <b>2016</b> , 31, 940-8	6.3		346

47	Greater milk intake is associated with lower bone turnover, higher bone density, and higher bone microarchitecture index in a population of elderly Japanese men with relatively low dietary calcium intake: Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Study. <i>Osteoporosis International</i> , <b>2015</b> , 26, 1585-94	5.3	17
46	Trabecular bone score may improve FRAX <sup>®</sup> prediction accuracy for major osteoporotic fractures in elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Cohort Study. <i>Osteoporosis International</i> , <b>2015</b> , 26, 1841-8	5.3	39
45	Age-related normative values of trabecular bone score (TBS) for Japanese women: the Japanese Population-based Osteoporosis (JPOS) study. <i>Osteoporosis International</i> , <b>2015</b> , 26, 245-52	5.3	28
44	Cohort Profile: The Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>International Journal of Epidemiology</i> , <b>2015</b> , 44, 405-14	7.8	14
43	Fat mass is positively associated with bone mass in relatively thin adolescents: data from the Kitakata Kids Health Study. <i>Bone</i> , <b>2014</b> , 64, 298-302	4.7	10
42	Trabecular bone score (TBS) predicts vertebral fractures in Japanese women over 10 years independently of bone density and prevalent vertebral deformity: the Japanese Population-Based Osteoporosis (JPOS) cohort study. <i>Journal of Bone and Mineral Research</i> , <b>2014</b> , 29, 399-407	6.3	114
41	Applicability of homeostasis model assessment of insulin resistance to patients with hyperglycemia: reply to Kawada. <i>Osteoporosis International</i> , <b>2013</b> , 24, 2735	5.3	
40	Renal function and bone mineral density in community-dwelling elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Study. <i>Bone</i> , <b>2013</b> , 56, 61-6	4.7	14
39	Biochemical markers for bone turnover predict risk of vertebral fractures in postmenopausal women over 10 years: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2013</b> , 24, 887-97	5.3	23
38	Association between vitamin K intake from fermented soybeans, natto, and bone mineral density in elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) study. <i>Osteoporosis International</i> , <b>2012</b> , 23, 705-14	5.3	55
37	Serum undercarboxylated osteocalcin levels are inversely associated with glycemic status and insulin resistance in an elderly Japanese male population: Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Study. <i>Osteoporosis International</i> , <b>2012</b> , 23, 761-70	5.3	98
36	Increased ratio of trunk to appendicular fat and increased blood pressure: study of a general population of Hamamatsu children. <i>Circulation Journal</i> , <b>2012</b> , 76, 2848-54	2.9	17
35	A screening model for low bone mass in elderly Japanese men using quantitative ultrasound measurements: Fujiwara-Kyo Study. <i>Journal of Clinical Densitometry</i> , <b>2012</b> , 15, 343-50	3.5	
34	Japanese 2011 guidelines for prevention and treatment of osteoporosis--executive summary. <i>Archives of Osteoporosis</i> , <b>2012</b> , 7, 3-20	2.9	228
33	Response to Effects of Vitamin K intake on gamma-carboxylated proteins, bone fractures, and vascular calcifications. <i>Osteoporosis International</i> , <b>2012</b> , 23, 1639-1640	5.3	0
32	Association between weight changes and changes in hip geometric indices in the Japanese female population during 10-year follow-up: Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2012</b> , 23, 1581-91	5.3	7
31	Alcohol intake and bone status in elderly Japanese men: baseline data from the Fujiwara-kyo osteoporosis risk in men (FORMEN) study. <i>Bone</i> , <b>2011</b> , 49, 275-80	4.7	31
30	Tracking of appendicular bone mineral density for 6 years including the pubertal growth spurt: Japanese Population-based Osteoporosis kids cohort study. <i>Journal of Bone and Mineral Metabolism</i> , <b>2011</b> , 29, 208-16	2.9	13

29	Impact of smoking on bone mineral density and bone metabolism in elderly men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) study. <i>Osteoporosis International</i> , <b>2011</b> , 22, 133-41	5.3	48
28	Age-specific reference values of hip geometric indices from a representative sample of the Japanese female population: Japanese Population-based Osteoporosis (JPOS) Study. <i>Osteoporosis International</i> , <b>2011</b> , 22, 1987-96	5.3	13
27	Fracture risk prediction using FRAX <sup>®</sup> : a 10-year follow-up survey of the Japanese Population-Based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2011</b> , 22, 3037-45	5.3	38
26	Peroxisome proliferator-activated receptor gamma polymorphism is related to peak bone mass: the JPOS study. <i>Osteoporosis International</i> , <b>2010</b> , 21, 321-9	5.3	12
25	Prevalent vertebral deformity independently increases incident vertebral fracture risk in middle-aged and elderly Japanese women: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2010</b> , 21, 1513-22	5.3	28
24	Smoking among premenopausal women is associated with increased risk of low bone status: the JPOS Study. <i>Journal of Bone and Mineral Metabolism</i> , <b>2010</b> , 28, 320-7	2.9	15
23	Design and baseline characteristics of a prospective cohort study for determinants of osteoporotic fracture in community-dwelling elderly Japanese men: the Fujiwara-kyo osteoporosis risk in men (FORMEN) study. <i>BMC Musculoskeletal Disorders</i> , <b>2009</b> , 10, 165	2.8	29
22	Low bone mass is associated with carotid atherosclerosis in postmenopausal women: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2009</b> , 20, 53-60	5.3	51
21	Arm span increases predictive value of models for prevalent vertebral deformities: the Japanese Population-based Osteoporosis (JPOS) Study. <i>Maturitas</i> , <b>2009</b> , 64, 241-5	5	4
20	Use of anthropometric indicators in screening for undiagnosed vertebral fractures: a cross-sectional analysis of the Fukui Osteoporosis Cohort (FOC) study. <i>BMC Musculoskeletal Disorders</i> , <b>2008</b> , 9, 157	2.8	15
19	Which element of physical activity is more important for determining bone growth in Japanese children and adolescents: the degree of impact, the period, the frequency, or the daily duration of physical activity?. <i>Journal of Bone and Mineral Metabolism</i> , <b>2008</b> , 26, 366-72	2.9	17
18	Biochemical markers of bone turnover may predict progression to osteoporosis in osteopenic women: the JPOS Cohort Study. <i>Journal of Bone and Mineral Metabolism</i> , <b>2007</b> , 25, 122-9	2.9	15
17	Efficacy of optimization of vitamin D in preventing osteoporosis and osteoporotic fractures: A systematic review. <i>Environmental Health and Preventive Medicine</i> , <b>2006</b> , 11, 155-70	4.2	5
16	Intake of fermented soybeans, natto, is associated with reduced bone loss in postmenopausal women: Japanese Population-Based Osteoporosis (JPOS) Study. <i>Journal of Nutrition</i> , <b>2006</b> , 136, 1323-8	4.1	115
15	Trunk muscle strength is a strong predictor of bone loss in postmenopausal women. <i>Clinical Orthopaedics and Related Research</i> , <b>2006</b> , 443, 66-72	2.2	13
14	Biochemical markers of bone turnover predict bone loss in perimenopausal women but not in postmenopausal women-the Japanese Population-based Osteoporosis (JPOS) Cohort Study. <i>Osteoporosis International</i> , <b>2006</b> , 17, 1086-95	5.3	43
13	Efficacy of Optimization of Vitamin D in Preventing Osteoporosis and Osteoporotic Fractures: A Systematic Review. <i>Environmental Health and Preventive Medicine</i> , <b>2006</b> , 11, 155-170	4.2	
12	Development of a food-frequency questionnaire to measure the dietary calcium intake of adult Japanese women. <i>Tohoku Journal of Experimental Medicine</i> , <b>2005</b> , 207, 217-22	2.4	26

11	Effects of the Cdx-2 polymorphism of the vitamin D receptor gene and lifestyle factors on bone mineral density in a representative sample of Japanese women: the Japanese Population-based Osteoporosis (JPOS) Study. <i>Calcified Tissue International</i> , <b>2005</b> , 77, 339-47	3.9	18
10	Effects of pubertal development, height, weight, and grip strength on the bone mineral density of the lumbar spine and hip in peripubertal Japanese children: Kyoto kids increase density in the skeleton study (Kyoto KIDS study). <i>Journal of Bone and Mineral Metabolism</i> , <b>2005</b> , 23, 463-9	2.9	22
9	Prediction of bone mineral density from vitamin D receptor polymorphisms is uncertain in representative samples of Japanese Women. The Japanese Population-based Osteoporosis (JPOS) Study. <i>International Journal of Epidemiology</i> , <b>2004</b> , 33, 979-88	7.8	29
8	Ultrasound bone densitometry of the calcaneus, determined with Sahara, in healthy Japanese adolescents: Japanese Population-based Osteoporosis (JPOS) Study. <i>Journal of Bone and Mineral Metabolism</i> , <b>2004</b> , 22, 248-53	2.9	11
7	Reference database of biochemical markers of bone turnover for the Japanese female population. Japanese Population-based Osteoporosis (JPOS) Study. <i>Osteoporosis International</i> , <b>2004</b> , 15, 981-91	5.3	66
6	Age-specific values and cutoff levels for the diagnosis of osteoporosis in quantitative ultrasound measurements at the calcaneus with SAHARA in healthy Japanese women: Japanese population-based osteoporosis (JPOS) study. <i>Calcified Tissue International</i> , <b>2002</b> , 71, 1-9	3.9	19
5	Greater trunk muscle torque reduces postmenopausal bone loss at the spine independently of age, body size, and vitamin D receptor genotype in Japanese women. <i>Calcified Tissue International</i> , <b>2002</b> , 71, 300-7	3.9	34
4	Bone mineral density of the spine, hip and distal forearm in representative samples of the Japanese female population: Japanese Population-Based Osteoporosis (JPOS) Study. <i>Osteoporosis International</i> , <b>2001</b> , 12, 529-37	5.3	137
3	Reference data of forearm bone mineral density in healthy Japanese male and female subjects in the second decade based on calendar age and puberty onset: Japanese Population Based Osteoporosis (JPOS) study. <i>Osteoporosis International</i> , <b>2000</b> , 11, 858-65	5.3	10
2	Age, menopause, bone turnover markers and lumbar bone loss in healthy Japanese women. <i>Maturitas</i> , <b>1996</b> , 25, 59-67	5	42
1	Update of the fracture risk prediction tool FRAX: a systematic review of potential cohorts and analysis plan. <i>Osteoporosis International</i> ,	5.3	0