

# Farzad Bagherzadeh Kasmani

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8205904/publications.pdf>

Version: 2024-02-01

14  
papers

247  
citations

1170033

9  
h-index

1181555

14  
g-index

14  
all docs

14  
docs citations

14  
times ranked

309  
citing authors

#	ARTICLE	IF	CITATIONS
1	Excess dietary tryptophan mitigates aflatoxicosis in growing quails. <i>Journal of Animal Physiology and Animal Nutrition</i> , 2019, 103, 1462-1473.	1.0	7
2	<i>Plantago ovata</i> in broiler chicken nutrition: Performance, carcass criteria, intestinal morphology, immunity, and intestinal bacterial population. <i>Journal of Animal Physiology and Animal Nutrition</i> , 2018, 102, e353-e363.	1.0	10
3	Estimation of lysine requirements for growing Japanese quails. <i>Journal of Animal Physiology and Animal Nutrition</i> , 2018, 102, 557-563.	1.0	7
4	<i>Nigella sativa</i> (black cumin seed) as a biological detoxifier in diet contaminated with aflatoxin B <sub>1</sub> . <i>Journal of Animal Physiology and Animal Nutrition</i> , 2017, 101, e77-e86.	1.0	12
5	Growth responses of breast and leg muscles to essential amino acids in broiler chicks. <i>Animal</i> , 2016, 10, 390-395.	1.3	7
6	A simple estimation of ideal profile of essential amino acids and metabolizable energy for growing Japanese quail. <i>Journal of Animal Physiology and Animal Nutrition</i> , 2016, 100, 680-685.	1.0	1
7	Methionine requirement of growing Japanese quails. <i>Animal Feed Science and Technology</i> , 2016, 212, 122-128.	1.1	22
8	Effects of a multi-strain probiotics against aflatoxicosis in growing Japanese quails. <i>Livestock Science</i> , 2015, 177, 110-116.	0.6	14
9	<i>Mentha piperita</i> (peppermint) in growing Japanese quails diet: Performance, carcass attributes, morphology and microbial populations of intestine. <i>Animal Feed Science and Technology</i> , 2015, 207, 104-111.	1.1	19
10	<i>Mentha piperita</i> (peppermint) in growing Japanese quails diet: Serum biochemistry, meat quality, humoral immunity. <i>Animal Feed Science and Technology</i> , 2015, 206, 57-66.	1.1	32
11	Estimation of lysine requirements of growing Japanese quail during the fourth and fifth weeks of age. <i>Poultry Science</i> , 2015, 94, 1923-1927.	1.5	9
12	Effects of dietary <i>Mentha spicata</i> on performance, blood metabolites, meat quality and microbial ecosystem of small intestine in growing Japanese quail. <i>Animal Feed Science and Technology</i> , 2014, 194, 89-98.	1.1	30
13	Estimation of Optimal Lysine in Quail Chicks During the Second and Third Weeks of Age. <i>Italian Journal of Animal Science</i> , 2013, 12, e84.	0.8	6
14	A novel aflatoxin-binding <i>Bacillus</i> probiotic: Performance, serum biochemistry, and immunological parameters in Japanese quail. <i>Poultry Science</i> , 2012, 91, 1846-1853.	1.5	71