Irwin Nazareth

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8208782/publications.pdf

Version: 2024-02-01

224 papers

22,738 citations

18482 62 h-index 9103 144 g-index

230 all docs

230 docs citations

times ranked

230

33097 citing authors

#	Article	IF	CITATIONS
1	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. Applied Health Economics and Health Policy, 2022, 20, 269-282.	2.1	6
2	Diagnostic accuracy and usability of the EMBalance decision support system for vestibular disorders in primary care: proof of concept randomised controlled study results. Journal of Neurology, 2022, 269, 2584-2598.	3.6	5
3	Exploratory randomised trial of face-to-face and mobile phone counselling against usual care for tobacco cessation in Indian primary care: a randomised controlled trial protocol for project CERTAIN. BMJ Open, 2022, 12, e048628.	1.9	1
4	Developing and internally validating a prognostic model (P Risk) to improve the prediction of psychosis in a primary care population using electronic health records: The MAPPED study. Schizophrenia Research, 2022, 246, 241-249.	2.0	1
5	Cigarette smoking as a risk factor for schizophrenia or all non-affective psychoses. Psychological Medicine, 2021, 51, 1373-1381.	4.5	10
6	Long-term systemic glucocorticoid therapy and weight gain: a population-based cohort study. Rheumatology, 2021, 60, 1502-1511.	1.9	14
7	Trends in generalised anxiety disorders and symptoms in primary care: UK population-based cohort study. British Journal of Psychiatry, 2021, 218, 158-164.	2.8	32
8	Time trends in the incidence of clinically diagnosed type 2 diabetes and pre-diabetes in the UK 2009–2018: a retrospective cohort study. BMJ Open Diabetes Research and Care, 2021, 9, e001989.	2.8	14
9	Serum urate and lung cancer: a cohort study and Mendelian randomization using UK Biobank. Respiratory Research, 2021, 22, 179.	3.6	4
10	Maintenance or Discontinuation of Antidepressants in Primary Care. New England Journal of Medicine, 2021, 385, 1257-1267.	27.0	75
11	Use of a personalised depression intervention in primary care to prevent anxiety: a secondary study of a cluster randomised trial. British Journal of General Practice, 2021, 71, e95-e104.	1.4	8
12	Diagnostic accuracy and usability of the EMBalance decision support system for vestibular disorders in primary care: proof of concept randomised controlled study results. Journal of Neurology, 2021, , .	3.6	0
13	Antidepressant medication to prevent depression relapse in primary care: the ANTLER RCT. Health Technology Assessment, 2021, 25, 1-62.	2.8	9
14	Cardiovascular outcomes of type 2 diabetic patients treated with DPPâ€⁴4 inhibitors versus sulphonylureas as add-on to metformin in clinical practice. Scientific Reports, 2021, 11, 23826.	3.3	1
15	Effectiveness of cognitive–behavioural therapy for depression in advanced cancer: CanTalk randomised controlled trial. British Journal of Psychiatry, 2020, 216, 213-221.	2.8	24
16	Genetically raised serum bilirubin levels and lung cancer: a cohort study and Mendelian randomisation using UK Biobank. Thorax, 2020, 75, 955-964.	5.6	32
17	Improving the assessment and management of obesity in UK children and adolescents: the PROMISE research programme including a RCT. Programme Grants for Applied Research, 2020, 8, 1-264.	1.0	4
18	Observational studies of treatment effectiveness: worthwhile or worthless?. Clinical Epidemiology, 2019, Volume 11, 35-42.	3.0	31

#	Article	IF	CITATIONS
19	Comparative effectiveness of sitagliptin vs sulphonylureas in older people. Age and Ageing, 2019, 48, 725-732.	1.6	2
20	Barriers to recruitment when conducting a commissioned randomised controlled trial of medication versus psychological therapy for generalised anxiety disorder: some lessons learned. Trials, 2019, 20, 284.	1.6	5
21	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpRession): study protocol for a randomised controlled trial. Trials. 2019. 20. 319.	1.6	12
22	Family physicians' views on participating in prevention of major depression. The predictD-EVAL qualitative study. PLoS ONE, 2019, 14, e0217621.	2.5	7
23	<p>Health indicator recording in UK primary care electronic health records: key implications for handling missing data</p> . Clinical Epidemiology, 2019, Volume 11, 157-167.	3.0	38
24	Manualised cognitive–behavioural therapy in treating depression in advanced cancer: the CanTalk RCT. Health Technology Assessment, 2019, 23, 1-106.	2.8	18
25	Primary care management of cardiovascular risk for people with severe mental illnesses: the Primrose research programme including cluster RCT. Programme Grants for Applied Research, 2019, 7, 1-98.	1.0	6
26	Clinical and cost-effectiveness of an intervention for reducing cholesterol and cardiovascular risk for people with severe mental illness in English primary care: a cluster randomised controlled trial. Lancet Psychiatry,the, 2018, 5, 145-154.	7.4	66
27	A personalized intervention to prevent depression in primary care: cost-effectiveness study nested into a clustered randomized trial. BMC Medicine, 2018, 16, 28.	5.5	21
28	Costâ€effectiveness of personal tailored risk information and taster sessions to increase the uptake of the NHS stop smoking services: the Start2quit randomized controlled trial. Addiction, 2018, 113, 708-718.	3.3	8
29	Landmark Models for Optimizing the Use of Repeated Measurements of Risk Factors in Electronic Health Records to Predict Future Disease Risk. American Journal of Epidemiology, 2018, 187, 1530-1538.	3.4	35
30	Factors influencing recording of drug misuse in primary care: a qualitative study of GPs in England. British Journal of General Practice, 2018, 68, e234-e244.	1.4	8
31	Cost-effectiveness of a community-delivered multicomponent intervention compared with enhanced standard care of obese adolescents: cost-utility analysis alongside a randomised controlled trial (the) Tj $ETQq1\ 1$	0. 7 84314	l rgBT /Overl
32	Weight change over two years in people prescribed olanzapine, quetiapine and risperidone in UK primary care: Cohort study in THIN, a UK primary care database. Journal of Psychopharmacology, 2018, 32, 1098-1103.	4.0	9
33	Impact of Financial Incentives on Alcohol Consumption Recording in Primary Health Care Among Adults with Schizophrenia and Other Psychoses: A Cross-Sectional and Retrospective Cohort Study. Alcohol and Alcoholism, 2017, 52, 197-205.	1.6	3
34	Predicting the onset of hazardous alcohol drinking in primary care: development and validation of a simple risk algorithm. British Journal of General Practice, 2017, 67, e280-e292.	1.4	2
35	Implementing the compassion intervention, a model for integrated care for people with advanced dementia towards the end of life in nursing homes: a naturalistic feasibility study. BMJ Open, 2017, 7, e015515.	1.9	58
36	Effectiveness and cost-effectiveness of a cardiovascular risk prediction algorithm for people with severe mental illness (PRIMROSE). BMJ Open, 2017, 7, e018181.	1.9	13

#	Article	IF	Citations
37	A community-based motivational personalised lifestyle intervention to reduce BMI in obese adolescents: results from the Healthy Eating and Lifestyle Programme (HELP) randomised controlled trial. Archives of Disease in Childhood, 2017, 102, 695-701.	1.9	28
38	The role of self-regulatory skills and automaticity on the effectiveness of a brief weight loss habit-based intervention: secondary analysis of the 10 top tips randomised trial. International Journal of Behavioral Nutrition and Physical Activity, 2017, 14, 119.	4.6	70
39	Effects of chronic exposure of hydroxychloroquine/ chloroquine on the risk of cancer, metastasis, and death: a population-based cohort study on patients with connective tissue diseases. Clinical Epidemiology, 2017, Volume 9, 545-554.	3.0	10
40	Antiepileptic drugs prescribed in pregnancy and prevalence of major congenital malformations: comparative prevalence studies. Clinical Epidemiology, 2017, Volume 9, 95-103.	3.0	30
41	Effectiveness of sitagliptin compared to sulfonylureas for type 2 diabetes mellitus inadequately controlled on metformin: a systematic review and meta-analysis. BMJ Open, 2017, 7, e017260.	1.9	17
42	Start2quit: a randomised clinical controlled trial to evaluate the effectiveness and cost-effectiveness of using personal tailored risk information and taster sessions to increase the uptake of the NHS Stop Smoking Services. Health Technology Assessment, 2017, 21, 1-206.	2.8	11
43	Pilot of a randomised controlled trial of the selective serotonin reuptake inhibitor sertraline versus cognitive behavioural therapy for anxiety symptoms in people with generalised anxiety disorder who have failed to respond to low-intensity psychological treatments as defined by the National Institute for Health and Care Excellence guidelines. Health Technology Assessment, 2017, 21, 1-138.	2.8	6
44	Time trends in the prescription of statins for the primary prevention of cardiovascular disease in the United Kingdom: a cohort study using The Health Improvement Network primary care data. Clinical Epidemiology, 2016, 8, 123.	3.0	70
45	An algorithm for identification and classification of individuals with type 1 and type 2 diabetes mellitus in a large primary care database. Clinical Epidemiology, 2016, Volume 8, 373-380.	3.0	36
46	Common Infections in Patients Prescribed Systemic Glucocorticoids in Primary Care: A Population-Based Cohort Study. PLoS Medicine, 2016, 13, e1002024.	8.4	114
47	Intervention to Prevent Major Depression in Primary Care. Annals of Internal Medicine, 2016, 164, 656.	3.9	23
48	Trends in incidence, prevalence and prescribing in type 2 diabetes mellitus between 2000 and 2013 in primary care: a retrospective cohort study. BMJ Open, 2016, 6, e010210.	1.9	224
49	Monitoring of alcohol consumption in primary care among adults with bipolar disorder: A cross-sectional and retrospective cohort study. Journal of Affective Disorders, 2016, 198, 83-87.	4.1	2
50	Risks associated with antipsychotic treatment in pregnancy: Comparative cohort studies based on electronic health records. Schizophrenia Research, 2016, 176, 349-356.	2.0	36
51	Initiation and duration of selective serotonin reuptake inhibitor prescribing over time: UK cohort study. British Journal of Psychiatry, 2016, 209, 421-426.	2.8	29
52	Development of a model for integrated care at the end of life in advanced dementia: A whole systems UK-wide approach. Palliative Medicine, 2016, 30, 279-295.	3.1	34
53	Evaluating the clinical and cost effectiveness of a behaviour change intervention for lowering cardiovascular disease risk for people with severe mental illnesses in primary care (PRIMROSE study): study protocol for a cluster randomised controlled trial. Trials, 2016, 17, 80.	1.6	17
54	The clinical and cost effectiveness of cognitive behavioural therapy plus treatment as usual for the treatment of depression in advanced cancer (CanTalk): study protocol for a randomised controlled trial. Trials, 2016, 17, 113.	1.6	8

#	Article	IF	Citations
55	Suicidality in primary care patients who present with sadness and anhedonia: a prospective European study. BMC Psychiatry, 2016, 16, 94.	2.6	9
56	Cohort Profile: The Flu Watch Study. International Journal of Epidemiology, 2016, 46, dyv370.	1.9	22
57	Risks and benefits of psychotropic medication in pregnancy: cohort studies based on UK electronic primary care health records. Health Technology Assessment, 2016, 20, 1-176.	2.8	82
58	Monitoring of Patients on Long-Term Glucocorticoid Therapy. Medicine (United States), 2015, 94, e647.	1.0	48
59	Lithium Prescribing during Pregnancy: A UK Primary Care Database Study. PLoS ONE, 2015, 10, e0121024.	2.5	19
60	Cardiovascular Risk Prediction Models for People With Severe Mental Illness. JAMA Psychiatry, 2015, 72, 143.	11.0	127
61	Initiation rates of statin therapy for the primary prevention of cardiovascular disease: an assessment of differences between countries of the UK and between regions within England. BMJ Open, 2015, 5, e007207-e007207.	1.9	12
62	The risk for major depression conferred by childhood maltreatment is multiplied by <i>BDNF</i> and <i>SERT</i> genetic vulnerability: a replication study. Journal of Psychiatry and Neuroscience, 2015, 40, 187-196.	2.4	41
63	Natural T Cell–mediated Protection against Seasonal and Pandemic Influenza. Results of the Flu Watch Cohort Study. American Journal of Respiratory and Critical Care Medicine, 2015, 191, 1422-1431.	5 . 6	229
64	Synthetic Glucocorticoids and Early Variations of Blood Pressure: A Population-Based Cohort Study. Journal of Clinical Endocrinology and Metabolism, 2015, 100, 2777-2783.	3.6	10
65	Patients' Opinions about Knowing Their Risk for Depression and What to Do about It. The PredictD-Qualitative Study. PLoS ONE, 2014, 9, e92008.	2.5	15
66	Community-based recruitment of patients with COPD into clinical research. Thorax, 2014, 69, 951-952.	5 . 6	5
67	Cost-Effectiveness of Computer-Tailored Smoking Cessation Advice in Primary Care: A Randomized Trial (ESCAPE). Nicotine and Tobacco Research, 2014, 16, 270-278.	2.6	16
68	Prescribing of antipsychotics in UK primary care: a cohort study. BMJ Open, 2014, 4, e006135.	1.9	144
69	Trap of trends to statistical significance: likelihood of near significant P value becoming more significant with extra data. BMJ, The, 2014, 348, g2215-g2215.	6.0	90
70	Smoker, ex-smoker or non-smoker? The validity of routinely recorded smoking status in UK primary care: a cross-sectional study. BMJ Open, 2014, 4, e004958.	1.9	59
71	Discontinuation of antipsychotic medication in pregnancy: A cohort study. Schizophrenia Research, 2014, 159, 218-225.	2.0	51
72	Serum uric acid and the risk of respiratory disease: a population-based cohort study. Thorax, 2014, 69, 1021-1026.	5 . 6	60

#	Article	IF	CITATIONS
73	Evaluation of twoâ€fold fully conditional specification multiple imputation for longitudinal electronic health record data. Statistics in Medicine, 2014, 33, 3725-3737.	1.6	46
74	Socio-demographic variation in chest pain incidence and subsequent coronary heart disease in primary care in the United Kingdom. European Journal of Preventive Cardiology, 2014, 21, 566-575.	1.8	11
75	Psychological distress, gender and dietary factors in South Asians: a cross-sectional survey. Public Health Nutrition, 2014, 17, 1538-1546.	2.2	19
76	Comparative community burden and severity of seasonal and pandemic influenza: results of the Flu Watch cohort study. Lancet Respiratory Medicine, the, 2014, 2, 445-454.	10.7	341
77	Erectile dysfunction. BMJ, The, 2014, 348, g129-g129.	6.0	69
78	Prenatal exposure to selective serotonin reuptake inhibitors and autistic symptoms in young children: another red herring?. British Journal of Psychiatry, 2014, 205, 105-106.	2.8	10
79	Trends in management and outcomes of COPD patients in primary care, 2000–2009: a retrospective cohort study. Npj Primary Care Respiratory Medicine, 2014, 24, 14015.	2.6	37
80	Assessment of an incentivised scheme to provide annual health checks in primary care for adults with intellectual disability: a longitudinal cohort study. Lancet Psychiatry,the, 2014, 1, 522-530.	7.4	58
81	Gender Differences in Survival and the Use of Primary Care Prior to Diagnosis of Three Cancers: An Analysis of Routinely Collected UK General Practice Data. PLoS ONE, 2014, 9, e101562.	2.5	31
82	Predicting the Onset of Anxiety Syndromes at 12 Months in Primary Care Attendees. The PredictA-Spain Study. PLoS ONE, 2014, 9, e106370.	2.5	12
83	Risk factors for onset of multiple or long major depressive episodes versus single and short episodes. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 1067-1075.	3.1	13
84	Preventing the onset of major depression based on the level and profile of risk of primary care attendees: protocol of a cluster randomised trial (the predictD-CCRT study). BMC Psychiatry, 2013, 13, 171.	2.6	20
85	Developing and evaluating complex interventions: The new Medical Research Council guidance. International Journal of Nursing Studies, 2013, 50, 587-592.	5.6	1,138
86	Relative incidence of common cancers in people with severe mental illness. Cohort study in the United Kingdom THIN primary care database. Schizophrenia Research, 2013, 143, 44-49.	2.0	36
87	Psychometric properties of the List of Threatening Experiencesâ€"LTE and its association with psychosocial factors and mental disorders according to different scoring methods. Journal of Affective Disorders, 2013, 150, 931-940.	4.1	90
88	<scp>G</scp> ilbert's syndrome and the risk of death: A populationâ€based cohort study. Journal of Gastroenterology and Hepatology (Australia), 2013, 28, 1643-1647.	2.8	61
89	Safety of AS03-adjuvanted split-virion H1N1 (2009) pandemic influenza vaccine: a prospective cohort study. BMJ Open, 2013, 3, e001912.	1.9	11
90	Suicide-related events in young people following prescription of SSRIs and other antidepressants: a self-controlled case series analysis. BMJ Open, 2013, 3, e003247.	1.9	16

#	Article	IF	Citations
91	Do men consult less than women? An analysis of routinely collected UK general practice data. BMJ Open, 2013, 3, e003320.	1.9	377
92	ESCAPE: a randomised controlled trial of computerâ€ŧailored smoking cessation advice in primary care. Addiction, 2013, 108, 811-819.	3.3	18
93	Alcohol consumption screening of newly-registered patients in primary care: a cross-sectional analysis. British Journal of General Practice, 2013, 63, e706-e712.	1.4	26
94	Use of long-term antibiotic treatment in COPD patients in the UK: a retrospective cohort study. Primary Care Respiratory Journal: Journal of the General Practice Airways Group, 2013, 22, 271-277.	2.3	20
95	Strategies to improve retention in randomised trials. The Cochrane Library, 2013, , MR000032.	2.8	166
96	Response to Letter Regarding Article, "Cardiovascular Events as a Function of Serum Bilirubin Levels in a Large, Statin-Treated Cohort― Circulation, 2013, 127, e861.	1.6	0
97	Mortality and Medical Care after Bereavement: A General Practice Cohort Study. PLoS ONE, 2013, 8, e52561.	2.5	32
98	Association of Common Mental Disorders and Quality of Life with the Frequency of Attendance in Slovenian Family Medicine Practices: Longitudinal Study. PLoS ONE, 2013, 8, e54241.	2.5	17
99	Recording of Severe Mental Illness in United Kingdom Primary Care, 2000–2010. PLoS ONE, 2013, 8, e82365.	2.5	69
100	The Sexunzipped Trial: Optimizing the Design of Online Randomized Controlled Trials. Journal of Medical Internet Research, 2013, 15, e278.	4.3	47
101	Severe Neuropsychiatric Outcomes Following Discontinuation of Long-Term Glucocorticoid Therapy. Journal of Clinical Psychiatry, 2013, 74, e281-e286.	2.2	24
102	Depression, anxiety and physical function: exploring the strength of causality. Journal of Epidemiology and Community Health, 2012, 66, e25-e25.	3.7	47
103	Suicidal Behavior and Severe Neuropsychiatric Disorders Following Glucocorticoid Therapy in Primary Care. American Journal of Psychiatry, 2012, 169, 491-497.	7.2	228
104	Factors predicting recruitment to a UK wide primary care smoking cessation study (the ESCAPE trial). Family Practice, 2012, 29, 110-117.	1.9	28
105	Cardiovascular Events as a Function of Serum Bilirubin Levels in a Large, Statin-Treated Cohort. Circulation, 2012, 126, 2556-2564.	1.6	122
106	Does recognition of depression in primary care affect outcome? The PREDICT-NL study. Family Practice, 2012, 29, 16-23.	1.9	33
107	Selective Serotonin Reuptake Inhibitors and Risk for Major Congenital Anomalies. Obstetrics and Gynecology, 2012, 119, 182-183.	2.4	2
108	Response to Meduri and Shih Letter. American Journal of Psychiatry, 2012, 169, 867-868.	7.2	0

#	Article	IF	CITATIONS
109	Differential Impact of Risk Factors for Women and Men on the Risk of Major Depressive Disorder. Annals of Epidemiology, 2012, 22, 388-396.	1.9	26
110	Study protocol for the 10 Top Tips (10TT) Trial: Randomised controlled trial of habit-based advice for weight control in general practice. BMC Public Health, 2012, 12, 667.	2.9	82
111	Evaluating the effectiveness of using personal tailored risk information and taster sessions to increase the uptake of smoking cessation services: study protocol for a randomised controlled trial. Trials, 2012, 13, 195.	1.6	8
112	Trends in Depression and Antidepressant Prescribing in Children and Adolescents: A Cohort Study in The Health Improvement Network (THIN). PLoS ONE, 2012, 7, e33181.	2.5	78
113	The Role of Primary Care in Service Provision for People with Severe Mental Illness in the United Kingdom. PLoS ONE, 2012, 7, e36468.	2.5	118
114	Recent Trends in the Incidence of Anxiety Diagnoses and Symptoms in Primary Care. PLoS ONE, 2012, 7, e41670.	2.5	55
115	Antiepileptic Drugs during Pregnancy in Primary Care: A UK Population Based Study. PLoS ONE, 2012, 7, e52339.	2.5	54
116	National changes in oral antipsychotic treatment for people with schizophrenia in primary care between 1998 and 2007 in the United Kingdom. Pharmacoepidemiology and Drug Safety, 2012, 21, 161-169.	1.9	27
117	Risk of cardiovascular events in people prescribed glucocorticoids with iatrogenic Cushing's syndrome: cohort study. BMJ, The, 2012, 345, e4928-e4928.	6.0	88
118	Recent life events pose greatest risk for onset of major depressive disorder during mid-life. Journal of Affective Disorders, 2012, 136, 505-513.	4.1	19
119	The natural course and outcome of major depressive disorder in primary care: the PREDICT-NL study. Social Psychiatry and Psychiatric Epidemiology, 2012, 47, 87-95.	3.1	92
120	Access to Cancer Screening in People with Learning Disabilities in the UK: Cohort Study in the Health Improvement Network, a Primary Care Research Database. PLoS ONE, 2012, 7, e43841.	2.5	50
121	Should men be screened and treated for postnatal depression?. Expert Review of Neurotherapeutics, 2011, 11, 1-3.	2.8	19
122	Inequalities in the provision of cardiovascular screening to people with severe mental illnesses in primary care. Schizophrenia Research, 2011, 129, 104-110.	2.0	55
123	Managing Sexual Difficulties: A Qualitative Investigation of Coping Strategies. Journal of Sex Research, 2011, 48, 325-333.	2.5	27
124	Pregnancy and safety outcomes in women vaccinated with an ASO3-adjuvanted split virion H1N1 (2009) pandemic influenza vaccine during pregnancy: A prospective cohort study. Vaccine, 2011, 29, 6358-6365.	3.8	69
125	Pregnancy as a Major Determinant for Discontinuation of Antidepressants. Journal of Clinical Psychiatry, 2011, 72, 979-985.	2.2	158
126	Prescribing Trends in Bipolar Disorder: Cohort Study in the United Kingdom THIN Primary Care Database 1995–2009. PLoS ONE, 2011, 6, e28725.	2.5	100

#	Article	IF	Citations
127	Polymorphic variation at the serotonin 1-A receptor gene is associated with comorbid depression and generalized anxiety. Psychiatric Genetics, 2011, 21, 195-201.	1.1	48
128	Cardiovascular Screening of People With Severe Mental Illness in England: Views of Service Users and Providers. Psychiatric Services, 2011, 62, 1338-1345.	2.0	14
129	Do women consult more than men? A review of gender and consultation for back pain and headache. Journal of Health Services Research and Policy, 2011, 16, 108-117.	1.7	130
130	Scripting sexual function: a qualitative investigation. Sociology of Health and Illness, 2011, 33, 540-553.	2.1	28
131	Associations between unemployment and major depressive disorder: Evidence from an international, prospective study (the predict cohort). Social Science and Medicine, 2011, 73, 1627-1634.	3.8	105
132	Assessing the efficacy of the healthy eating and lifestyle programme (HELP) compared with enhanced standard care of the obese adolescent in the community: study protocol for a randomized controlled trial. Trials, 2011, 12, 242.	1.6	22
133	Cochrane Review: Interactive computerâ€based interventions for sexual health promotion. Evidence-Based Child Health: A Cochrane Review Journal, 2011, 6, 2192-2268.	2.0	3
134	Commentary on †Interactive computer†based interventions for sexual health promotion†with a response from the review authors. Evidence-Based Child Health: A Cochrane Review Journal, 2011, 6, 2271-2274.	2.0	1
135	Prevalence of long-term oral glucocorticoid prescriptions in the UK over the past 20 years. Rheumatology, 2011, 50, 1982-1990.	1.9	227
136	Serum Bilirubin and Risk of Respiratory Disease and Death. JAMA - Journal of the American Medical Association, 2011, 305, 691.	7.4	174
137	Self-Reported Long-Term Needs After Stroke. Stroke, 2011, 42, 1398-1403.	2.0	362
138	The influence of socio-demographic characteristics on consultation for back pain-a review of the literature. Family Practice, 2011, 28, 163-171.	1.9	14
139	Heavy Episodic Drinking in Europe: A Cross Section Study in Primary Care in Six European Countries. Alcohol and Alcoholism, 2011, 46, 600-606.	1.6	22
140	Development and Validation of a Risk Model for Prediction of Hazardous Alcohol Consumption in General Practice Attendees: The PredictAL Study. PLoS ONE, 2011, 6, e22175.	2.5	6
141	The pattern of physical comorbidity and the psychosocial determinants of depression: a prospective cohort study on a representative sample of family practice attendees in Slovenia. Mental Health in Family Medicine, 2011, 8, 147-55.	0.2	6
142	Interactive computer-based interventions for sexual health promotion. The Cochrane Library, 2010, , CD006483.	2.8	130
143	Polyclinics and polysystems: evolving ambiguity, evidence required. British Journal of General Practice, 2010, 60, 400-401.	1.4	6
144	â€~At the moment it is just a couple of eccentrics doing it': concordance in day-to-day practice. International Journal of Pharmacy Practice, 2010, 13, 273-280.	0.6	4

#	Article	IF	Citations
145	Longstanding disease, disability or infirmity and depression in primary care. Wiener Klinische Wochenschrift, 2010, 122, 567-571.	1.9	3
146	Impact of a nurse-led intervention to improve screening for cardiovascular risk factors in people with severe mental illnesses. Phase-two cluster randomised feasibility trial of community mental health teams. BMC Health Services Research, 2010, 10, 61.	2.2	63
147	The Patient Health Questionnaire-9 for detection of major depressive disorder in primary care: consequences of current thresholds in a crosssectional study. BMC Family Practice, 2010, 11, 98.	2.9	147
148	Does dissemination extend beyond publication: a survey of a cross section of public funded research in the UK. Implementation Science, 2010, 5, 61.	6.9	52
149	Disseminating research findings: what should researchers do? A systematic scoping review of conceptual frameworks. Implementation Science, 2010, 5, 91.	6.9	171
150	Issues in multiple imputation of missing data for large general practice clinical databases. Pharmacoepidemiology and Drug Safety, 2010, 19, 618-626.	1.9	99
151	Angina in primary care in Goa, India: sex differences and associated risk factors. Heart Asia, 2010, 2, 28-35.	1.1	2
152	Oral antibiotic prescribing during pregnancy in primary care: UK population-based study. Journal of Antimicrobial Chemotherapy, 2010, 65, 2238-2246.	3.0	68
153	Psychosocial and sociodemographic predictors of attrition in a longitudinal study of major depression in primary care: the predictD-Spain study. Journal of Epidemiology and Community Health, 2010, 64, 874-884.	3.7	27
154	Incidence of Maternal and Paternal Depression in Primary Care. JAMA Pediatrics, 2010, 164, 1038-44.	3.0	156
155	Comparison of risk factors for the onset and maintenance of depression. British Journal of Psychiatry, 2010, 196, 13-17.	2.8	31
156	Promoting self care for minor illness. BMJ: British Medical Journal, 2010, 340, c2913-c2913.	2.3	11
157	Survival of people with clinical diagnosis of dementia in primary care: cohort study. BMJ: British Medical Journal, 2010, 341, c3584-c3584.	2.3	171
158	Effects of a Financial Incentive on Health Researchers' Response to an Online Survey: a Randomized Controlled Trial. Journal of Medical Internet Research, 2010, 12, e13.	4.3	28
159	The Potential of Web-based Interventions for Heart Disease Self-Management: A Mixed Methods Investigation. Journal of Medical Internet Research, 2010, 12, e56.	4.3	47
160	Is there a place for mental health research in general practice?. London Journal of Primary Care, 2009, 2, 5-7.	0.9	0
161	A clinical prediction rule for detecting major depressive disorder in primary care: the PREDICT-NL study. Family Practice, 2009, 26, 241-250.	1.9	17
162	Highâ€activity variants of the uMAOA polymorphism increase the risk for depression in a large primary care sample. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2009, 150B, 395-402.	1.7	44

#	Article	IF	CITATIONS
163	"Some of These Words I Can't Pronounceâ€. A Qualitative Exploration of the Readability of Generic and Tailored Selfâ€help Material for Quitting Smoking. Journal of Applied Biobehavioral Research, 2009, 14, 1-14.	2.0	3
164	Perception of Computerâ€tailored Feedback for Smoking Cessation: Qualitative Findings from Focus Groups. Journal of Applied Biobehavioral Research, 2009, 14, 15-29.	2.0	1
165	Recent trends in the incidence of recorded depression in primary care. British Journal of Psychiatry, 2009, 195, 520-524.	2.8	135
166	Major paternal depression and child consultation for developmental and behavioural problems. British Journal of General Practice, 2009, 59, 180-185.	1.4	16
167	Economic evaluation of arthritis self management in primary care. BMJ, The, 2009, 339, b3532-b3532.	6.0	39
168	Bespoke cohort studies needed. BMJ: British Medical Journal, 2009, 339, b3512-b3512.	2.3	3
169	A Comparison of Father and Mother Report of Child Behaviour on the Strengths and Difficulties Questionnaire. Child Psychiatry and Human Development, 2008, 39, 399-413.	1.9	66
170	Associations between paternal depression and behaviour problems in children of $4\hat{a}\in 6$ years. European Child and Adolescent Psychiatry, 2008, 17, 306-315.	4.7	48
171	A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. BMC Psychiatry, 2008, 8, 70.	2.6	1,554
172	Relative risk of diabetes, dyslipidaemia, hypertension and the metabolic syndrome in people with severe mental illnesses: Systematic review and metaanalysis. BMC Psychiatry, 2008, 8, 84.	2.6	161
173	Impact of demographic factors on recognition of persons with depression and anxiety in primary care in Slovenia. BMC Psychiatry, 2008, 8, 96.	2.6	15
174	Effectiveness of computer-tailored Smoking Cessation Advice in Primary Care (ESCAPE): a Randomised Trial. Trials, 2008, 9, 23.	1.6	19
175	Predicting the onset and persistence of episodes of depression in primary health care. The predictD-Spain study: Methodology. BMC Public Health, 2008, 8, 256.	2.9	23
176	Suicide and severe mental illnesses. Cohort study within the UK general practice research database. Schizophrenia Research, 2008, 99, 134-138.	2.0	69
177	Applying user-generated quality criteria to develop an Internet intervention for patients with heart disease. Journal of Telemedicine and Telecare, 2008, 14, 124-127.	2.7	6
178	Panic disorder and risk of new onset coronary heart disease, acute myocardial infarction, and cardiac mortality: cohort study using the general practice research database. European Heart Journal, 2008, 29, 2981-2988.	2.2	78
179	Developing and evaluating complex interventions: the new Medical Research Council guidance. BMJ, The, 2008, 337, a1655.	6.0	7,454
180	Why promote the findings of single research studies?. BMJ: British Medical Journal, 2008, 336, 722-722.	2.3	14

#	Article	IF	Citations
181	Development and Validation of an International Risk Prediction Algorithm for Episodes of Major Depression in General Practice Attendees. Archives of General Psychiatry, 2008, 65, 1368.	12.3	134
182	Prevalence of common mental disorders in general practice attendees across Europe. British Journal of Psychiatry, 2008, 192, 362-367.	2.8	227
183	Assessing the feasibility of proactive recruitment of smokers to an intervention in general practice for smoking cessation using computer-tailored feedback reports. Family Practice, 2007, 24, 395-400.	1.9	20
184	Attitudes to depression and its treatment in primary care. Psychological Medicine, 2007, 37, 1239-1248.	4.5	29
185	Relative Risk of Cardiovascular and Cancer Mortality in People With Severe Mental Illness From the United Kingdom's General Practice Research Database. Archives of General Psychiatry, 2007, 64, 242.	12.3	527
186	Treatment of depression in primary care. British Journal of Psychiatry, 2007, 191, 164-169.	2.8	28
187	Information from the Internet and the doctor-patient relationship: the patient perspective $\hat{a}\in$ a qualitative study. BMC Family Practice, 2007, 8, 47.	2.9	144
188	Physical activity, dietary habits and Coronary Heart Disease risk factor knowledge amongst people with severe mental illness. Social Psychiatry and Psychiatric Epidemiology, 2007, 42, 787-793.	3.1	163
189	Women's Views of their Sexual Difficulties: Agreement and Disagreement with Clinical Diagnoses. Archives of Sexual Behavior, 2007, 36, 281-288.	1.9	93
190	Patterns of presentations of dizziness in primary care—a cross-sectional cluster analysis study. Journal of Psychosomatic Research, 2006, 60, 395-401.	2.6	29
191	Risk for coronary heart disease in people with severe mental illness. British Journal of Psychiatry, 2006, 188, 271-277.	2.8	99
192	A Feasibility Study on Recruiting Fathers of Young Children to Examine the Impact of Paternal Depression on Child Development. Child Psychiatry and Human Development, 2006, 36, 295-309.	1.9	26
193	Prevention of coronary heart disease in people with severe mental illnesses: a qualitative study of patient and professionals' preferences for care. BMC Psychiatry, 2006, 6, 16.	2.6	53
194	The health of people classified as lesbian, gay and bisexual attending family practitioners in London: a controlled study. BMC Public Health, 2006, 6, 127.	2.9	46
195	Prediction of depression in European general practice attendees: the PREDICT study. BMC Public Health, 2006, 6, 6.	2.9	71
196	The 5â€HTTLPR s/s genotype at the serotonin transporter gene (SLC6A4) increases the risk for depression in a large cohort of primary care attendees: The PREDICTâ€gene study. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2006, 141B, 912-917.	1.7	83
197	Self management of arthritis in primary care: randomised controlled trial. BMJ: British Medical Journal, 2006, 333, 879.	2.3	168
198	HARMFUL ALCOHOL USE IN GOA, INDIA, AND ITS ASSOCIATIONS WITH VIOLENCE: A STUDY IN PRIMARY CARE. Alcohol and Alcoholism, 2006, 42, 131-137.	1.6	54

#	Article	IF	Citations
199	Internet Interventions for Long-Term Conditions: Patient and Caregiver Quality Criteria. Journal of Medical Internet Research, 2006, 8, e13.	4.3	101
200	Internet Interventions or Patient Education Web Sites? – Author's Reply. Journal of Medical Internet Research, 2006, 8, e19.	4.3	2
201	The association of paternal mood and infant temperament: A pilot study. British Journal of Developmental Psychology, 2005, 23, 609-621.	1.7	35
202	Patient preferences in randomised controlled trials: Conceptual framework and implications for research. Social Science and Medicine, 2005, 61, 685-695.	3.8	70
203	Impact of Participant and Physician Intervention Preferences on Randomized Trials. JAMA - Journal of the American Medical Association, 2005, 293, 1089.	7.4	234
204	Interactive Health Communication Applications for people with chronic disease. The Cochrane Library, 2005, , CD004274.	2.8	456
205	"Is Cybermedicine Killing You?" - A Response From the Authors of the Cochrane Review. Journal of Medical Internet Research, 2005, 7, e40.	4.3	1
206	Sexuality of health practitioners is complex. BMJ: British Medical Journal, 2004, 328, 1567.4-1568.	2.3	0
207	Problems with sexual function in people attending London general practitioners: cross sectional study. BMJ: British Medical Journal, 2003, 327, 423-0.	2.3	187
208	Evaluation of a complex intervention for changing professional behaviour: the Evidence Based Out Reach (EBOR) Trial. Journal of Health Services Research and Policy, 2002, 7, 230-238.	1.7	30
209	Pharmacist follow-up telephone call reduces return visits to hospital. Evidence-Based Healthcare and Public Health, 2002, 6, 125-126.	0.0	0
210	A randomised controlled trial of the effect of educational outreach by community pharmacists on prescribing in UK general practice. British Journal of General Practice, 2002, 52, 290-5.	1.4	51
211	Sexual dysfunction after treatment for testicular cancer. Journal of Psychosomatic Research, 2001, 51, 735-743.	2.6	78
212	A pharmacy discharge plan for hospitalized elderly patients—a randomized controlled trial. Age and Ageing, 2001, 30, 33-40.	1.6	249
213	GPs' views on their management of sexual dysfunction. Family Practice, 2001, 18, 516-518.	1.9	122
214	Panic Disorder with Agoraphobia Associated with Dizziness: Characteristic Symptoms and Psychosocial Sequelae. Journal of Nervous and Mental Disease, 2001, 189, 321-327.	1.0	38
215	Designing trials of interventions to change professional practice in primary care: lessons from an exploratory study of two change strategies. BMJ: British Medical Journal, 2000, 320, 1580-1583.	2.3	26
216	Medical education in the community - the UNITRA experience. Medical Education, 1999, 33, 722-724.	2.1	14

#	Article	IF	CITATIONS
217	A Randomized Trial of Evidence-Based Outreach (EBOR). Contemporary Clinical Trials, 1999, 20, 479-492.	1.9	23
218	Psychiatric dysfunction and dizziness. Lancet, The, 1999, 353, 2069.	13.7	4
219	\hat{l}^2 -blockers in heart failure. Lancet, The, 1999, 353, 1011.	13.7	6
220	Monitoring Psychosis in General Practice: A Controlled Trial. British Journal of Psychiatry, 1996, 169, 475-482.	2.8	23
221	Withdrawal of triazolam's product licence: effect on patients 18 months later. Addiction, 1995, 90, 927-934.	3.3	3
222	Drug users' views on general practitioners. BMJ: British Medical Journal, 1995, 310, 302-302.	2.3	26
223	The urethral syndrome: A controlled evaluation. Journal of Psychosomatic Research, 1993, 37, 737-743.	2.6	10
224	A Guide to Psychiatry in Primary Care. By Patricia R. Casey Hampshire: Wrightson Biomedical Publishing Ltd. 1990. 269 pp. £19.50 British Journal of Psychiatry, 1991, 159, 753-754.	2.8	O