Siu Tim Cheung

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8208944/siu-tim-cheung-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

82
papers
4,489
citations
4,919
ext. papers
4,919
ext. citations
34
papers
66
g-index
7.2
avg, IF
L-index

| # | Paper | IF | Citations |
|----|--|---------------|-----------|
| 82 | Progranulin mediates immune evasion of pancreatic ductal adenocarcinoma through regulation of MHCI expression <i>Nature Communications</i> , 2022 , 13, 156 | 17.4 | 4 |
| 81 | Contaminated and misidentified cell lines commonly use in cancer research. <i>Molecular Carcinogenesis</i> , 2020 , 59, 573-574 | 5 | 4 |
| 80 | Genetic variation in ABCB5 associates with risk of hepatocellular carcinoma. <i>Journal of Cellular and Molecular Medicine</i> , 2020 , 24, 10705-10713 | 5.6 | 2 |
| 79 | Clinical application of Albumin-Bilirubin (ALBI) score: The current status. <i>Journal of the Royal College of Surgeons of Edinburgh</i> , 2020 , 18, 178-186 | 2.5 | 11 |
| 78 | The ATP-binding cassette transporter ABCF1 is a hepatic oncofetal protein that promotes chemoresistance, EMT and cancer stemness in hepatocellular carcinoma. <i>Cancer Letters</i> , 2019 , 457, 98- | 1 <i>6</i> 99 | 17 |
| 77 | STAT3: An Emerging Therapeutic Target for Hepatocellular Carcinoma. <i>Cancers</i> , 2019 , 11, | 6.6 | 40 |
| 76 | C-terminal truncated HBx protein activates caveolin-1/LRP6/Etatenin/FRMD5 axis in promoting hepatocarcinogenesis. <i>Cancer Letters</i> , 2019 , 444, 60-69 | 9.9 | 16 |
| 75 | A Brief Overview of Progranulin in Health and Disease. <i>Methods in Molecular Biology</i> , 2018 , 1806, 3-15 | 1.4 | 29 |
| 74 | Treatment of advanced hepatocellular carcinoma: immunotherapy from checkpoint blockade to potential of cellular treatment. <i>Translational Gastroenterology and Hepatology</i> , 2018 , 3, 89 | 5.2 | 20 |
| 73 | Establishment and characterization of new tumor xenografts and cancer cell lines from EBV-positive nasopharyngeal carcinoma. <i>Nature Communications</i> , 2018 , 9, 4663 | 17.4 | 63 |
| 72 | Granulin epithelin precursor promotes colorectal carcinogenesis by activating MARK/ERK pathway. Journal of Translational Medicine, 2018 , 16, 150 | 8.5 | 9 |
| 71 | Novel biomarkers GEP/ABCB5 regulate response to adjuvant transarterial chemoembolization after curative hepatectomy for hepatocellular carcinoma. <i>Hepatobiliary and Pancreatic Diseases International</i> , 2018 , 17, 524-530 | 2.1 | 3 |
| 70 | Mouse Monoclonal Antibodies Against Progranulin (PGRN/GEP) as Therapeutics in Preclinical Cancer Models. <i>Methods in Molecular Biology</i> , 2018 , 1806, 131-144 | 1.4 | 1 |
| 69 | Methods to Analyze the Role of Progranulin (PGRN/GEP) on Cancer Stem Cell Features. <i>Methods in Molecular Biology</i> , 2018 , 1806, 145-153 | 1.4 | 1 |
| 68 | Granulin-epithelin precursor interacts with 78-kDa glucose-regulated protein in hepatocellular carcinoma. <i>BMC Cancer</i> , 2017 , 17, 409 | 4.8 | 6 |
| 67 | Octamer 4/microRNA-1246 signaling axis drives Wnt/Etatenin activation in liver cancer stem cells. <i>Hepatology</i> , 2016 , 64, 2062-2076 | 11.2 | 122 |
| 66 | Comprehensive characterization of the patient-derived xenograft and the paralleled primary hepatocellular carcinoma cell line. <i>Cancer Cell International</i> , 2016 , 16, 41 | 6.4 | 12 |

(2011-2016)

| 65 | Hepatic cancer stem cell marker granulin-epithelin precursor and Ecatenin expression associate with recurrence in hepatocellular carcinoma. <i>Oncotarget</i> , 2016 , 7, 21644-57 | 3.3 | 21 |
|----|--|------|-----|
| 64 | Overexpression of PIN1 Enhances Cancer Growth and Aggressiveness with Cyclin D1 Induction in EBV-Associated Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2016 , 11, e0156833 | 3.7 | 23 |
| 63 | Pancreatic cancer risk variant in LINC00673 creates a miR-1231 binding site and interferes with PTPN11 degradation. <i>Nature Genetics</i> , 2016 , 48, 747-57 | 36.3 | 187 |
| 62 | Copy number gain of granulin-epithelin precursor (GEP) at chromosome 17q21 associates with overexpression in human liver cancer. <i>BMC Cancer</i> , 2015 , 15, 264 | 4.8 | 4 |
| 61 | Restoration of natural killer activity in hepatocellular carcinoma by treatment with antibody against granulin-epithelin precursor. <i>Oncolmmunology</i> , 2015 , 4, e1016706 | 7.2 | 13 |
| 60 | Glioma Association and Balancing Selection of ZFPM2. <i>PLoS ONE</i> , 2015 , 10, e0133003 | 3.7 | 5 |
| 59 | Establishment and characterization of a novel primary hepatocellular carcinoma cell line with metastatic ability in vivo. <i>Cancer Cell International</i> , 2014 , 14, 103 | 6.4 | 22 |
| 58 | Granulin-epithelin precursor interacts with heparan sulfate on liver cancer cells. <i>Carcinogenesis</i> , 2014 , 35, 2485-94 | 4.6 | 13 |
| 57 | MicroRNA-494 within an oncogenic microRNA megacluster regulates G1/S transition in liver tumorigenesis through suppression of mutated in colorectal cancer. <i>Hepatology</i> , 2014 , 59, 202-15 | 11.2 | 97 |
| 56 | Granulin-epithelin precursor renders hepatocellular carcinoma cells resistant to natural killer cytotoxicity. <i>Cancer Immunology Research</i> , 2014 , 2, 1209-19 | 12.5 | 28 |
| 55 | Antibody against granulin-epithelin precursor sensitizes hepatocellular carcinoma to chemotherapeutic agents. <i>Molecular Cancer Therapeutics</i> , 2014 , 13, 3001-12 | 6.1 | 23 |
| 54 | Cancer stem-like cells in Epstein-Barr virus-associated nasopharyngeal carcinoma. <i>Chinese Journal of Cancer</i> , 2014 , 33, 529-38 | | 18 |
| 53 | Complete genomic sequence of Epstein-Barr virus in nasopharyngeal carcinoma cell line C666-1. <i>Infectious Agents and Cancer</i> , 2013 , 8, 29 | 3.5 | 32 |
| 52 | Identification of common variants in BRCA2 and MAP2K4 for susceptibility to sporadic pancreatic cancer. <i>Carcinogenesis</i> , 2013 , 34, 1001-5 | 4.6 | 18 |
| 51 | Identification and characterization of tropomyosin 3 associated with granulin-epithelin precursor in human hepatocellular carcinoma. <i>PLoS ONE</i> , 2012 , 7, e40324 | 3.7 | 20 |
| 50 | CD44+ cancer stem-like cells in EBV-associated nasopharyngeal carcinoma. <i>PLoS ONE</i> , 2012 , 7, e52426 | 3.7 | 60 |
| 49 | Genome-wide association study identifies five loci associated with susceptibility to pancreatic cancer in Chinese populations. <i>Nature Genetics</i> , 2011 , 44, 62-6 | 36.3 | 141 |
| 48 | Granulin-epithelin precursor and ATP-dependent binding cassette (ABC)B5 regulate liver cancer cell chemoresistance. <i>Gastroenterology</i> , 2011 , 140, 344-55 | 13.3 | 114 |

| 47 | Granulin-epithelin precursor is an oncofetal protein defining hepatic cancer stem cells. <i>PLoS ONE</i> , 2011 , 6, e28246 | 3.7 | 42 |
|----|---|------|-----|
| 46 | Bmi1 functions as an oncogene independent of Ink4A/Arf repression in hepatic carcinogenesis. <i>Molecular Cancer Research</i> , 2009 , 7, 1937-45 | 6.6 | 55 |
| 45 | New tools for functional genomic analysis. <i>Drug Discovery Today</i> , 2009 , 14, 754-60 | 8.8 | 26 |
| 44 | Albumin mRNA in plasma predicts post-transplant recurrence of patients with hepatocellular carcinoma. <i>Transplantation</i> , 2008 , 85, 81-7 | 1.8 | 28 |
| 43 | Granulin-epithelin precursor as a therapeutic target for hepatocellular carcinoma. <i>Hepatology</i> , 2008 , 47, 1524-32 | 11.2 | 90 |
| 42 | Atypical localization of membrane type 1-matrix metalloproteinase in the nucleus is associated with aggressive features of hepatocellular carcinoma. <i>Molecular Carcinogenesis</i> , 2007 , 46, 225-30 | 5 | 51 |
| 41 | Overexpression of NDRG1 is an indicator of poor prognosis in hepatocellular carcinoma. <i>Modern Pathology</i> , 2007 , 20, 76-83 | 9.8 | 91 |
| 40 | Down-regulation of retinol binding protein 5 is associated with aggressive tumor features in hepatocellular carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2007 , 133, 929-36 | 4.9 | 7 |
| 39 | Distinct pathways of genomic progression to benign and malignant tumors of the liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 14771-6 | 11.5 | 165 |
| 38 | Genomic and proteomic biomarkers for diagnosis and prognosis of hepatocellular carcinoma. <i>Biomarkers in Medicine</i> , 2007 , 1, 273-84 | 2.3 | 17 |
| 37 | Inhibition of hepatocellular carcinoma invasion by suppression of claudin-10 in HLE cells. <i>Molecular Cancer Therapeutics</i> , 2007 , 6, 2858-67 | 6.1 | 38 |
| 36 | Increased expression of glycosyl-phosphatidylinositol anchor attachment protein 1 (GPAA1) is associated with gene amplification in hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2006 , 119, 1330-7 | 7.5 | 32 |
| 35 | Hedgehog signaling in human hepatocellular carcinoma. Cancer Biology and Therapy, 2006, 5, 111-7 | 4.6 | 87 |
| 34 | Liver intestine-cadherin (CDH17) haplotype is associated with increased risk of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2006 , 12, 5248-52 | 12.9 | 31 |
| 33 | Preoperative plasma transcript AA454543 level is an independent prognostic factor for hepatocellular carcinoma after partial hepatectomy. <i>Neoplasia</i> , 2006 , 8, 696-701 | 6.4 | 8 |
| 32 | GEP associates with wild-type p53 in hepatocellular carcinoma. <i>Oncology Reports</i> , 2006 , 15, 1507 | 3.5 | 1 |
| 31 | Anterior approach versus conventional approach right hepatic resection for large hepatocellular carcinoma: a prospective randomized controlled study. <i>Annals of Surgery</i> , 2006 , 244, 194-203 | 7.8 | 184 |
| 30 | SPARC and Hevin expression correlate with tumour angiogenesis in hepatocellular carcinoma. Journal of Pathology, 2006 , 210, 459-68 | 9.4 | 65 |

(2001-2006)

| 29 | GEP associates with wild-type p53 in hepatocellular carcinoma. Oncology Reports, 2006, 15, 1507-11 | 3.5 | 13 |
|----|---|------|-----|
| 28 | Transcript AA454543 is a novel prognostic marker for hepatocellular carcinoma after curative partial hepatectomy. <i>Neoplasia</i> , 2005 , 7, 91-8 | 6.4 | 9 |
| 27 | Cyclins and CDKs in Liver Diseases 2005 , 325-331 | | |
| 26 | An integrated data analysis approach to characterize genes highly expressed in hepatocellular carcinoma. <i>Oncogene</i> , 2005 , 24, 3737-47 | 9.2 | 107 |
| 25 | Array-based comparative genomic hybridization reveals recurrent chromosomal aberrations and Jab1 as a potential target for 8q gain in hepatocellular carcinoma. <i>Carcinogenesis</i> , 2005 , 26, 2050-7 | 4.6 | 113 |
| 24 | Mechanism of metastasis by membrane type 1-matrix metalloproteinase in hepatocellular carcinoma. World Journal of Gastroenterology, 2005, 11, 6269-76 | 5.6 | 18 |
| 23 | Claudin-10 expression level is associated with recurrence of primary hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2005 , 11, 551-6 | 12.9 | 80 |
| 22 | Granulin-epithelin precursor overexpression promotes growth and invasion of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2004 , 10, 7629-36 | 12.9 | 104 |
| 21 | Clinical significance of thrombospondin 1 expression in hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2004 , 10, 4150-7 | 12.9 | 86 |
| 20 | Novel endothelial cell markers in hepatocellular carcinoma. <i>Modern Pathology</i> , 2004 , 17, 1198-210 | 9.8 | 69 |
| 19 | Decreased expression of cytochrome P450 2E1 is associated with poor prognosis of hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2004 , 111, 494-500 | 7.5 | 27 |
| 18 | De novo sarcoma of donor origin in a liver allograft determined by microsatellite analysis: a short report. <i>Liver Transplantation</i> , 2004 , 10, 320-3 | 4.5 | 5 |
| 17 | Fibrosing cholestatic hepatitis secondary to precore/core promoter hepatitis B variant with lamivudine resistance: successful retransplantation with combination adefovir dipivoxil and hepatitis B immunoglobulin. <i>Liver Transplantation</i> , 2004 , 10, 557-63 | 4.5 | 26 |
| 16 | Development of antibody to hepatitis B surface antigen after liver transplantation for chronic hepatitis B. <i>Hepatology</i> , 2003 , 37, 36-43 | 11.2 | 59 |
| 15 | Liver as an ideal target for gene therapy: expression of CTLA4Ig by retroviral gene transfer. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2002 , 17, 1008-14 | 4 | 24 |
| 14 | Gene expression patterns in human liver cancers. <i>Molecular Biology of the Cell</i> , 2002 , 13, 1929-39 | 3.5 | 680 |
| 13 | Identify metastasis-associated genes in hepatocellular carcinoma through clonality delineation for multinodular tumor. <i>Cancer Research</i> , 2002 , 62, 4711-21 | 10.1 | 74 |
| 12 | Liver transplantation in Asian patients with chronic hepatitis B using lamivudine prophylaxis. <i>Annals of Surgery</i> , 2001 , 233, 276-81 | 7.8 | 110 |

| 11 | Novel intertypic recombinants of epstein-barr virus in the chinese population. <i>Journal of Virology</i> , 2000 , 74, 1544-8 | 6.6 | 55 |
|----|---|-----------------|-----|
| 10 | CTL control of EBV in nasopharyngeal carcinoma (NPC): EBV-specific CTL responses in the blood and tumors of NPC patients and the antigen-processing function of the tumor cells. <i>Journal of Immunology</i> , 2000 , 165, 573-82 | 5.3 | 96 |
| 9 | Evaluation of quantitative PCR and branched-chain DNA assay for detection of hepatitis B virus DNA in sera from hepatocellular carcinoma and liver transplant patients. <i>Journal of Clinical Microbiology</i> , 2000 , 38, 1977-80 | 9.7 | 9 |
| 8 | Indications for liver transplantation in patients with chronic hepatitis B and C virus infection and hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2000 , 15 Suppl, E181- | -6 ⁴ | 18 |
| 7 | Inhibiting tumorigenic potential by restoration of p16 in nasopharyngeal carcinoma. <i>British Journal of Cancer</i> , 1999 , 81, 1122-6 | 8.7 | 40 |
| 6 | Nasopharyngeal carcinoma cell line (C666-1) consistently harbouring Epstein-Barr virus. <i>International Journal of Cancer</i> , 1999 , 83, 121-6 | 7.5 | 304 |
| 5 | Nasopharyngeal carcinoma cell line (C666-1) consistently harbouring Epstein-Barr virus 1999 , 83, 121 | | 2 |
| 4 | Characterization of a new EBV-associated nasopharyngeal carcinoma cell line. <i>Cancer Genetics and Cytogenetics</i> , 1998 , 101, 83-8 | | 34 |
| 3 | Specific latent membrane protein 1 gene sequences in type 1 and type 2 Epstein-Barr virus from nasopharyngeal carcinoma in Hong Kong. <i>International Journal of Cancer</i> , 1998 , 76, 399-406 | 7.5 | 56 |
| 2 | Prevalence of LMP1 deletion variant of Epstein-Barr virus in nasopharyngeal carcinoma and gastric tumors in Hong Kong. <i>International Journal of Cancer</i> , 1996 , 66, 711-2 | 7.5 | 52 |
| 1 | Prevalence of LMP1 deletion variant of Epstein-Barr virus in nasopharyngeal carcinoma and gastric tumors in Hong Kong 1996 , 66, 711 | | 1 |