

# Taejun Yoon

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8208967/publications.pdf>

Version: 2024-02-01

21  
papers

79  
citations

1684188  
5  
h-index

1588992  
8  
g-index

21  
all docs

21  
docs citations

21  
times ranked

110  
citing authors

#	ARTICLE	IF	CITATIONS
1	GeTe Nanosheets as Theranostic Agents for Multimodal Imaging and Therapy of Inflammatory Bowel Disease. <i>Advanced Functional Materials</i> , 2022, 32, 2107433.	14.9	7
2	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021, 35, 1761-1768.	1.3	1
3	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , 2021, 23, 77.	3.5	8
4	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021, 40, 3703-3710.	2.2	1
5	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021, 134, 1168-1174.	2.3	0
6	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021, 2021, 1-8.	3.0	2
7	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021, 16, e0254226.	2.5	1
8	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021, 62, 1016.	2.2	3
9	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e24048.	2.1	1
10	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021, 69, 91-95.	1.6	0
11	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, . .	0.8	0
12	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4975.	4.1	14
13	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020, 2020, 1-9.	1.3	4
14	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020, 22, 73.	3.5	6
15	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.	1.2	7
16	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020, 61, 720.	2.2	2
17	Atializumab, a humanized anti-aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , 2019, 220, 119408.	11.4	7
18	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019, 2019, 1-6.	1.9	7

#	ARTICLE	IF	CITATIONS
19	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019, 38, 1685-1690.	2.2	5
20	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121.	0.8	2
21	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	1