
## Phui-Ly Liew

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8213409/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Epigenetic biomarker and drug development in gynecological cancers. , 2022, , 223-255.		0
2	Epigenomic Analysis Reveals the KCNK9 Potassium Channel as a Potential Therapeutic Target for Adenomyosis. International Journal of Molecular Sciences, 2022, 23, 5973.	4.1	3
3	BHLHE22 Expression Is Associated with a Proinflammatory Immune Microenvironment and Confers a Favorable Prognosis in Endometrial Cancer. International Journal of Molecular Sciences, 2022, 23, 7158.	4.1	2
4	Nerve growth factor interacts with CHRM4 and promotes neuroendocrine differentiation of prostate cancer and castration resistance. Communications Biology, 2021, 4, 22.	4.4	25
5	Complete remission of heavily treated ovarian clear cell carcinoma with ARID1A mutations after pembrolizumab and bevacizumab combination therapy: a case report. Journal of Ovarian Research, 2020, 13, 143.	3.0	9
6	Hypermethylation and decreased expression of TMEM240 are potential early-onset biomarkers for colorectal cancer detection, poor prognosis, and early recurrence prediction. Clinical Epigenetics, 2020, 12, 67.	4.1	12
7	Clinicopathologic and Prognostic Significance of Thymopoietin-α Overexpression in Gastric Cancer. Journal of Cancer, 2019, 10, 5099-5107.	2.5	5
8	Combined genetic mutations and DNA-methylated genes as biomarkers for endometrial cancer detection from cervical scrapings. Clinical Epigenetics, 2019, 11, 170.	4.1	28
9	Distinct methylation profile of mucinous ovarian carcinoma reveals susceptibility to proteasome inhibitors. International Journal of Cancer, 2018, 143, 355-367.	5.1	12
10	GATA3 as a master regulator and therapeutic target in ovarian highâ€grade serous carcinoma stem cells. International Journal of Cancer, 2018, 143, 3106-3119.	5.1	31
11	Epigenetic loss of heparan sulfate 3â€ <scp>O</scp> â€sulfation sensitizes ovarian carcinoma to oncogenic signals and predicts prognosis. International Journal of Cancer, 2018, 143, 1943-1953.	5.1	11
12	High HMGA2 Expression Correlates with Reduced Recurrence-free Survival and Poor Overall Survival in Oral Squamous Cell Carcinoma. Anticancer Research, 2017, 37, 1891-1899.	1.1	18
13	DEF6 expression in ovarian carcinoma correlates with poor patient survival. Diagnostic Pathology, 2016, 11, 68.	2.0	9
14	Nasal dissemination of a single-clone IgH-rearranged conjunctival MALT lymphoma through the nasolacrimal duct: A case report. Oncology Letters, 2016, 12, 1007-1010.	1.8	3
15	Prognostic and predictive values of Nrf2, Keap1, p16 and E-cadherin expression in ovarian epithelial carcinoma. International Journal of Clinical and Experimental Pathology, 2015, 8, 5642-9.	0.5	28
16	Hepatic tumor necrosis factor-î±, leptin and adiponectin expression in morbid obese patients: Clinicopathological correlations. Obesity Research and Clinical Practice, 2012, 6, e55-e62.	1.8	1
17	Roles of hepatic progenitor cells activation, ductular reaction proliferation and Notch signaling in morbid obesity. Hepato-Gastroenterology, 2012, 59, 1921-7.	0.5	3
18	Rapid Staining and Immediate Interpretation of Fine-Needle Aspiration Cytology for Palpable Breast Lesions: Diagnostic Accuracy, Mammographic, Ultrasonographic and Histopathologic Correlations. Acta Cytologica, 2011, 55, 30-37.	1.3	19

#	Article	IF	CITATIONS
19	Gallbladder Disease among Obese Patients in Taiwan. Obesity Surgery, 2007, 17, 383-390.	2.1	32