Toshi A Furukawa

List of Publications by Citations

Source: https://exaly.com/author-pdf/8214749/toshi-a-furukawa-publications-by-citations.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

656 21,877 69 129 h-index g-index citations papers 6.2 6.92 27,191 743 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
656	Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. <i>Lancet, The</i> , 2018 , 391, 1357-1366	40	1283
655	Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. <i>Lancet, The</i> , 2009 , 373, 746-58	40	1217
654	The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. <i>Psychological Medicine</i> , 2003 , 33, 357-62	6.9	1022
653	Imputing missing standard deviations in meta-analyses can provide accurate results. <i>Journal of Clinical Epidemiology</i> , 2006 , 59, 7-10	5.7	937
652	The performance of the Japanese version of the K6 and K10 in the World Mental Health Survey Japan. <i>International Journal of Methods in Psychiatric Research</i> , 2008 , 17, 152-8	4.3	709
651	Screening for serious mental illness in the general population with the K6 screening scale: results from the WHO World Mental Health (WMH) survey initiative. <i>International Journal of Methods in Psychiatric Research</i> , 2010 , 19 Suppl 1, 4-22	4.3	686
650	Relapse prevention with antidepressant drug treatment in depressive disorders: a systematic review. <i>Lancet, The</i> , 2003 , 361, 653-61	40	633
649	GRADE guidelines: 13. Preparing summary of findings tables and evidence profiles-continuous outcomes. <i>Journal of Clinical Epidemiology</i> , 2013 , 66, 173-83	5.7	371
648	Cross-cultural validation of the Beck Depression Inventory-II in Japan. <i>Psychiatry Research</i> , 2002 , 110, 291-9	9.9	296
647	Twelve-month prevalence, severity, and treatment of common mental disorders in communities in Japan: preliminary finding from the World Mental Health Japan Survey 2002-2003. <i>Psychiatry and Clinical Neurosciences</i> , 2005 , 59, 441-52	6.2	266
646	Waiting list may be a nocebo condition in psychotherapy trials: a contribution from network meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2014 , 130, 181-92	6.5	234
645	Advances in the GRADE approach to rate the certainty in estimates from a network meta-analysis. <i>Journal of Clinical Epidemiology</i> , 2018 , 93, 36-44	5.7	224
644	Can we individualize the 'number needed to treat'? An empirical study of summary effect measures in meta-analyses. <i>International Journal of Epidemiology</i> , 2002 , 31, 72-6	7.8	212
643	Comparative efficacy and tolerability of pharmacological treatments in the maintenance treatment of bipolar disorder: a systematic review and network meta-analysis. <i>Lancet Psychiatry,the</i> , 2014 , 1, 351-	9 ^{23.3}	207
642	Imputing response rates from means and standard deviations in meta-analyses. <i>International Clinical Psychopharmacology</i> , 2005 , 20, 49-52	2.2	199
641	Dose equivalents of antidepressants: Evidence-based recommendations from randomized controlled trials. <i>Journal of Affective Disorders</i> , 2015 , 180, 179-84	6.6	178
640	Which is more generalizable, powerful and interpretable in meta-analyses, mean difference or standardized mean difference?. <i>BMC Medical Research Methodology</i> , 2014 , 14, 30	4.7	146

(2021-2009)

639	Depression, inflammation, and pain in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 61, 1018-24		145
638	Systematic review and meta-analysis of cannabis treatment for chronic pain. <i>Pain Medicine</i> , 2009 , 10, 1353-68	2.8	137
637	Pooling health-related quality of life outcomes in meta-analysis-a tutorial and review of methods for enhancing interpretability. <i>Research Synthesis Methods</i> , 2011 , 2, 188-203	7.2	134
636	Effectiveness and Acceptability of Cognitive Behavior Therapy Delivery Formats in Adults With Depression: A Network Meta-analysis. <i>JAMA Psychiatry</i> , 2019 , 76, 700-707	14.5	128
635	Dose Equivalents for Second-Generation Antipsychotic Drugs: The Classical Mean Dose Method. <i>Schizophrenia Bulletin</i> , 2015 , 41, 1397-402	1.3	127
634	Multicentre prospective study of perinatal depression in Japan: incidence and correlates of antenatal and postnatal depression. <i>Archives of Womeng Mental Health</i> , 2006 , 9, 121-30	5	123
633	Psychotherapy plus antidepressant for panic disorder with or without agoraphobia: systematic review. <i>British Journal of Psychiatry</i> , 2006 , 188, 305-12	5.4	121
632	Placebo response rates in antidepressant trials: a systematic review of published and unpublished double-blind randomised controlled studies. <i>Lancet Psychiatry,the</i> , 2016 , 3, 1059-1066	23.3	112
631	Meta-analysis of effects and side effects of low dosage tricyclic antidepressants in depression: systematic review. <i>BMJ, The</i> , 2002 , 325, 991	5.9	111
630	Psychotherapy for depression among children and adolescents: a systematic review. <i>Acta Psychiatrica Scandinavica</i> , 2007 , 116, 84-95	6.5	109
629	Gradations of clinical severity and sensitivity to change assessed with the Beck Depression Inventory-II in Japanese patients with depression. <i>Psychiatry Research</i> , 2005 , 135, 229-35	9.9	109
628	Comparative efficacy and acceptability of first-generation and second-generation antidepressants in the acute treatment of major depression: protocol for a network meta-analysis. <i>BMJ Open</i> , 2016 , 6, e010919	3	105
627	Fluoxetine versus other types of pharmacotherapy for depression. <i>Cochrane Database of Systematic Reviews</i> , 2005 , CD004185		103
626	Hand-sewn versus mechanical esophagogastric anastomosis after esophagectomy: a systematic review and meta-analysis. <i>Annals of Surgery</i> , 2013 , 257, 238-48	7.8	102
625	Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan. <i>Psycho-Oncology</i> , 2011 , 20, 497-505	3.9	101
624	Assessment of mood: guides for clinicians. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 581-9	4.1	101
623	Optimal dose of selective serotonin reuptake inhibitors, venlafaxine, and mirtazapine in major depression: a systematic review and dose-response meta-analysis. <i>Lancet Psychiatry,the</i> , 2019 , 6, 601-60	3 3.3	99
622	Internet-Based Cognitive Behavioral Therapy for Depression: A Systematic Review and Individual Patient Data Network Meta-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 361-371	14.5	97

621	How to obtain NNT from Cohen's d: comparison of two methods. <i>PLoS ONE</i> , 2011 , 6, e19070	3.7	94
620	Association between unreported outcomes and effect size estimates in Cochrane meta-analyses. JAMA - Journal of the American Medical Association, 2007, 297, 468-70	27.4	92
619	Depression as a potential causal factor in subsequent miscarriage in recurrent spontaneous aborters. <i>Human Reproduction</i> , 2002 , 17, 2580-4	5.7	92
618	Time to recurrence after recovery from major depressive episodes and its predictors. <i>Psychological Medicine</i> , 2003 , 33, 839-45	6.9	91
617	Supported employment for adults with severe mental illness. <i>The Cochrane Library</i> , 2013 , CD008297	5.2	89
616	Psychotherapy for depression among incurable cancer patients. <i>The Cochrane Library</i> , 2008 , CD005537	5.2	89
615	A network meta-analysis of the effects of psychotherapies, pharmacotherapies and their combination in the treatment of adult depression. <i>World Psychiatry</i> , 2020 , 19, 92-107	14.4	89
614	Comparative Efficacy and Acceptability of 21 Antidepressant Drugs for the Acute Treatment of Adults With Major Depressive Disorder: A Systematic Review and Network Meta-Analysis. <i>Focus</i> (American Psychiatric Publishing), 2018 , 16, 420-429	1.1	89
613	Evidence-based guidelines for interpretation of the Panic Disorder Severity Scale. <i>Depression and Anxiety</i> , 2009 , 26, 922-9	8.4	88
612	Dismantling cognitive-behaviour therapy for panic disorder: a systematic review and component network meta-analysis. <i>Psychological Medicine</i> , 2018 , 48, 1945-1953	6.9	86
611	Brief behavioral therapy for refractory insomnia in residual depression: an assessor-blind, randomized controlled trial. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 1651-8	4.6	86
610	Is computerised CBT really helpful for adult depression?-A meta-analytic re-evaluation of CCBT for adult depression in terms of clinical implementation and methodological validity. <i>BMC Psychiatry</i> , 2013 , 13, 113	4.2	85
609	Minimally important difference estimates and methods: a protocol. <i>BMJ Open</i> , 2015 , 5, e007953	3	81
608	Escitalopram versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2009 , CD006532	5.2	81
607	Benzodiazepines in generalized anxiety disorder: heterogeneity of outcomes based on a systematic review and meta-analysis of clinical trials. <i>Journal of Psychopharmacology</i> , 2007 , 21, 774-82	4.6	81
606	Mirtazapine versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2011 , CD006528	5.2	80
605	Ketamine and other glutamate receptor modulators for depression in adults. <i>The Cochrane Library</i> , 2015 , CD011612	5.2	79
604	Certolizumab pegol for induction of remission in Crohn's disease. <i>The Cochrane Library</i> , 2017 ,	5.2	78

(2013-2011)

603	Strategic use of new generation antidepressants for depression, SUN(^_^)D: study design and rationale. <i>Trials</i> , 2011 , 12,	2.8	78	
602	Effectiveness of selective serotonin reuptake inhibitors. <i>Cmaj</i> , 2008 , 178, 1185-1186	3.5	78	
601	Social Support Questionnaire among psychiatric patients with various diagnoses and normal controls. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1999 , 34, 216-22	4.5	78	
600	Cognitive behavioural therapies versus other psychological therapies for depression. <i>The Cochrane Library</i> , 2018 ,	5.2	78	
599	Cognitive behavioural therapies versus treatment as usual for depression. <i>The Cochrane Library</i> , 2018 ,	5.2	78	
598	Roux-en-Y versus Billroth-I reconstruction after distal gastrectomy for gastric cancer. <i>The Cochrane Library</i> , 2018 ,	5.2	78	
597	Effectiveness of paroxetine in the treatment of acute major depression in adults: a systematic re-examination of published and unpublished data from randomized trials. <i>Cmaj</i> , 2008 , 178, 296-305	3.5	76	
596	Combined psychotherapy plus antidepressants for panic disorder with or without agoraphobia. <i>The Cochrane Library</i> , 2007 , CD004364	5.2	74	
595	Prognostic value of malnutrition assessed by Controlling Nutritional Status score for long-term mortality in patients with acute heart failure. <i>International Journal of Cardiology</i> , 2017 , 230, 529-536	3.2	73	
594	Evidence-based guidelines for interpretation of the Hamilton Rating Scale for Depression. <i>Journal of Clinical Psychopharmacology</i> , 2007 , 27, 531-4	1.7	73	
593	Preoperative inspiratory muscle training for postoperative pulmonary complications in adults undergoing cardiac and major abdominal surgery. <i>The Cochrane Library</i> , 2015 , CD010356	5.2	72	
592	Psychotherapy for depression among advanced, incurable cancer patients: A systematic review and meta-analysis. <i>Cancer Treatment Reviews</i> , 2017 , 56, 16-27	14.4	71	
591	From effect size into number needed to treat. <i>Lancet, The</i> , 1999 , 353, 1680	40	71	
590	Determinants of the quality of life in Alzheimer's disease patients as assessed by the Japanese version of the Quality of Life-Alzheimer's disease scale. <i>Dementia and Geriatric Cognitive Disorders</i> , 2006 , 21, 182-91	2.6	70	
589	Initial severity of schizophrenia and efficacy of antipsychotics: participant-level meta-analysis of 6 placebo-controlled studies. <i>JAMA Psychiatry</i> , 2015 , 72, 14-21	14.5	69	
588	Psychosocial factors, disease status, and quality of life in patients with rheumatoid arthritis. <i>Journal of Psychosomatic Research</i> , 2009 , 67, 425-31	4.1	66	
587	Smartphone Cognitive Behavioral Therapy as an Adjunct to Pharmacotherapy for Refractory Depression: Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2017 , 19, e373	7.6	65	
586	Validity of the Patient Health Questionnaire (PHQ)-9 and PHQ-2 in general internal medicine primary care at a Japanese rural hospital: a cross-sectional study. <i>General Hospital Psychiatry</i> , 2013 ,	5.6	64	

585	Psychological interventions to reduce positive symptoms in schizophrenia: systematic review and network meta-analysis. <i>World Psychiatry</i> , 2018 , 17, 316-329	14.4	64
584	'Third wave' cognitive and behavioural therapies versus other psychological therapies for depression. <i>The Cochrane Library</i> , 2013 , CD008704	5.2	63
583	Emotional distress and its correlates among parents of children with pervasive developmental disorders. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 651-7	6.2	63
582	Safety reporting and adverse-event profile of mirtazapine described in randomized controlled trials in comparison with other classes of antidepressants in the acute-phase treatment of adults with depression: systematic review and meta-analysis. <i>CNS Drugs</i> , 2010 , 24, 35-53	6.7	62
581	Patients' supportive care needs and psychological distress in advanced breast cancer patients in Japanese Journal of Clinical Oncology, 2011 , 41, 530-6	2.8	61
580	Neuropsychiatric symptoms predict change in quality of life of Alzheimer disease patients: a two-year follow-up study. <i>Psychiatry and Clinical Neurosciences</i> , 2009 , 63, 374-84	6.2	59
579	Alexithymia, depression and social support among Japanese workers. <i>Psychotherapy and Psychosomatics</i> , 2003 , 72, 307-14	9.4	59
578	Comparative efficacy and acceptability of antidepressants, psychotherapies, and their combination for acute treatment of children and adolescents with depressive disorder: a systematic review and network meta-analysis. <i>Lancet Psychiatry,the</i> , 2020 , 7, 581-601	23.3	58
577	Family psychoeducation for major depression: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2011 , 198, 385-90	5.4	58
576	Postictal suppression correlates with therapeutic efficacy for depression in bilateral sine and pulse wave electroconvulsive therapy. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 168-73	6.2	58
575	Psychological therapies for panic disorder with or without agoraphobia in adults: a network meta-analysis. <i>The Cochrane Library</i> , 2016 , 4, CD011004	5.2	57
574	Sertraline versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2010 , CD006117	5.2	56
573	Application of the Center for Epidemiologic Studies Depression Scale among first-visit psychiatric patients: a new approach to improve its performance. <i>Journal of Affective Disorders</i> , 1997 , 46, 1-13	6.6	56
572	The differential impact of executive attention dysfunction on episodic memory in obsessive-compulsive disorder patients with checking symptoms vs. those with washing symptoms. <i>Journal of Psychiatric Research</i> , 2007 , 41, 776-84	5.2	56
571	Screening for depression in terminally ill cancer patients in Japan. <i>Journal of Pain and Symptom Management</i> , 2006 , 31, 5-12	4.8	56
570	Is antidepressant-benzodiazepine combination therapy clinically more useful? A meta-analytic study. <i>Journal of Affective Disorders</i> , 2001 , 65, 173-7	6.6	56
569	Long-term Outcomes of Laparoscopic Versus Open Surgery for Clinical Stage I Gastric Cancer: The LOC-1 Study. <i>Annals of Surgery</i> , 2016 , 264, 214-22	7.8	56
568	Effects of an Internet-based cognitive behavioral therapy (iCBT) program in Manga format on improving subthreshold depressive symptoms among healthy workers: a randomized controlled trial. <i>PLoS ONE</i> , 2014 , 9, e97167	3.7	55

(2007-2000)

567	Time to recovery of an inception cohort with hitherto untreated unipolar major depressive episodes. <i>British Journal of Psychiatry</i> , 2000 , 177, 331-5	5.4	55
566	Early parental separation experiences among patients with bipolar disorder and major depression: a case-control study. <i>Journal of Affective Disorders</i> , 1999 , 52, 85-91	6.6	55
565	Sertraline versus other antidepressive agents for depression. <i>Cochrane Database of Systematic Reviews</i> , 2009 , CD006117		54
564	Behavioural therapies versus other psychological therapies for depression. <i>The Cochrane Library</i> , 2013 , CD008696	5.2	53
563	Cross-cultural equivalence in depression assessment: Japan-Europe-North American study. <i>Acta Psychiatrica Scandinavica</i> , 2005 , 112, 279-85	6.5	53
562	Publication bias is underreported in systematic reviews published in high-impact-factor journals: metaepidemiologic study. <i>Journal of Clinical Epidemiology</i> , 2014 , 67, 1320-6	5.7	52
561	Mirtazapine versus other antidepressants in the acute-phase treatment of adults with major depression: systematic review and meta-analysis. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1404-15	4.6	51
560	The epidemiology of chronic fatigue, physical illness, and symptoms of common mental disorders: a cross-sectional survey from the second British National Survey of Psychiatric Morbidity. <i>Journal of Psychosomatic Research</i> , 2008 , 64, 357-62	4.1	50
559	Living network meta-analysis compared with pairwise meta-analysis in comparative effectiveness research: empirical study. <i>BMJ, The</i> , 2018 , 360, k585	5.9	49
558	Changes after behavior therapy among responsive and nonresponsive patients with obsessive-compulsive disorder. <i>Psychiatry Research - Neuroimaging</i> , 2009 , 172, 242-50	2.9	49
557	Polymyxin B-immobilized hemoperfusion and mortality in critically ill adult patients with sepsis/septic shock: a systematic review with meta-analysis and trial sequential analysis. <i>Intensive Care Medicine</i> , 2018 , 44, 167-178	14.5	49
556	Does formulation matter? A systematic review and meta-analysis of oral versus long-acting antipsychotic studies. <i>Schizophrenia Research</i> , 2017 , 183, 10-21	3.6	48
555	Course of psychological distress and its predictors in advanced non-small cell lung cancer patients. <i>Psycho-Oncology</i> , 2006 , 15, 463-73	3.9	48
554	Direct oral anticoagulants versus warfarin for preventing stroke and systemic embolic events among atrial fibrillation patients with chronic kidney disease. <i>The Cochrane Library</i> , 2017 , 11, CD011373	5.2	47
553	Stratum-specific likelihood ratios of two versions of the general health questionnaire. <i>Psychological Medicine</i> , 2001 , 31, 519-29	6.9	47
552	Efficacy of cognitive bias modification interventions in anxiety and depressive disorders: a systematic review and network meta-analysis. <i>Lancet Psychiatry,the</i> , 2020 , 7, 506-514	23.3	46
551	Associated and predictive factors of sleep disturbance in advanced cancer patients. <i>Psycho-Oncology</i> , 2007 , 16, 888-94	3.9	46
550	The new GRID Hamilton Rating Scale for Depression demonstrates excellent inter-rater reliability for inexperienced and experienced raters before and after training. <i>Psychiatry Research</i> , 2007 , 153, 61-7	9.9	45

549	Weight changes and eating attitudes of Japanese adolescents under acculturative stresses: a prospective study. <i>International Journal of Eating Disorders</i> , 1994 , 15, 71-9	6.3	45
548	Telephone cognitive-behavioral therapy for subthreshold depression and presenteeism in workplace: a randomized controlled trial. <i>PLoS ONE</i> , 2012 , 7, e35330	3.7	45
547	Standardization of health outcomes assessment for depression and anxiety: recommendations from the ICHOM Depression and Anxiety Working Group. <i>Quality of Life Research</i> , 2017 , 26, 3211-3225	3.7	44
546	Distinct neuropsychological profiles of three major symptom dimensions in obsessive-compulsive disorder. <i>Psychiatry Research</i> , 2011 , 187, 166-73	9.9	44
545	Reliability and validity of the Japanese version of the Short-form Supportive Care Needs Survey questionnaire (SCNS-SF34-J). <i>Psycho-Oncology</i> , 2009 , 18, 1003-10	3.9	44
544	Choice of data extraction tools for systematic reviews depends on resources and review complexity. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 506-10	5.7	44
543	Metareview on short-term effectiveness and safety of antidepressants for depression: an evidence-based approach to inform clinical practice. <i>Canadian Journal of Psychiatry</i> , 2007 , 52, 553-62	4.8	44
542	Mood stabilizers and antipsychotics for acute mania: a systematic review and meta-analysis of combination/augmentation therapy versus monotherapy. <i>CNS Drugs</i> , 2014 , 28, 989-1003	6.7	43
541	Fluoxetine versus other types of pharmacotherapy for depression. <i>The Cochrane Library</i> , 2013 , CD0041	8 5 .2	43
540	Duloxetine versus other anti-depressive agents for depression. <i>The Cochrane Library</i> , 2012 , 10, CD0065	3 3 .2	43
539	Quality of life of parents raising children with pervasive developmental disorders. <i>BMC Psychiatry</i> , 2012 , 12, 119	4.2	43
538	Are all antidepressants really the same? The case of fluoxetine: a systematic review. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 850-64	4.6	41
537	Cognitive-Behavioral Analysis System of Psychotherapy, Drug, or Their Combination for Persistent Depressive Disorder: Personalizing the Treatment Choice Using Individual Participant Data Network Metaregression. <i>Psychotherapy and Psychosomatics</i> , 2018 , 87, 140-153	9.4	41
536	National Prescription Patterns of Antidepressants in the Treatment of Adults With Major Depression in the US Between 1996 and 2015: A Population Representative Survey Based Analysis. <i>Frontiers in Psychiatry</i> , 2020 , 11, 35	5	40
535	Symptom indicator of severity of depression in cancer patients: a comparison of the DSM-IV criteria with alternative diagnostic criteria. <i>General Hospital Psychiatry</i> , 2009 , 31, 225-32	5.6	39
534	Haplotype association between GABAA receptor gamma2 subunit gene (GABRG2) and methamphetamine use disorder. <i>Pharmacogenomics Journal</i> , 2005 , 5, 89-95	3.5	39
533	Social class inequalities in self-rated health and their gender and age group differences in Japan. Journal of Epidemiology, 2006 , 16, 223-32	3.4	38
532	Evaluating the credibility of anchor based estimates of minimal important differences for patient reported outcomes: instrument development and reliability study. <i>BMJ, The</i> , 2020 , 369, m1714	5.9	37

(2009-2016)

5	31	Continuation and discontinuation of benzodiazepine prescriptions: A cohort study based on a large claims database in Japan. <i>Psychiatry Research</i> , 2016 , 237, 201-7	9.9	37	
5	30	Citalopram versus other anti-depressive agents for depression. <i>The Cochrane Library</i> , 2012 , CD006534	5.2	37	
5	29	Antidepressant medications and other treatments of depressive disorders: a CINP Task Force report based on a review of evidence. <i>International Journal of Neuropsychopharmacology</i> , 2007 , 10 Suppl 1, S1-207	5.8	37	
5	28	Antidepressant plus benzodiazepine for major depression. <i>The Cochrane Library</i> , 2000 , CD001026	5.2	37	
5	27	Twelve-month use of mental health services in four areas in Japan: findings from the World Mental Health Japan Survey 2002-2003. <i>Psychiatry and Clinical Neurosciences</i> , 2006 , 60, 240-8	6.2	36	
5	26	Psychosocial predictors of successful delivery after unexplained recurrent spontaneous abortions: a cohort study. <i>Acta Psychiatrica Scandinavica</i> , 2004 , 109, 440-6	6.5	36	
5	25	Replication and contradiction of highly cited research papers in psychiatry: 10-year follow-up. British Journal of Psychiatry, 2015 , 207, 357-62	5.4	35	
5	24	'Third wave' cognitive and behavioural therapies versus treatment as usual for depression. <i>The Cochrane Library</i> , 2013 , CD008705	5.2	35	
5	23	Gender differences in factors associated with suicidal ideation in major depression among cancer patients. <i>Psycho-Oncology</i> , 2010 , 19, 384-9	3.9	35	
5	22	Cancer patients' reluctance to disclose their emotional distress to their physicians: a study of Japanese patients with lung cancer. <i>Psycho-Oncology</i> , 2008 , 17, 460-5	3.9	35	
5	21	An intervention to improve the interrater reliability of clinical EEG interpretations. <i>Psychiatry and Clinical Neurosciences</i> , 2003 , 57, 485-9	6.2	35	
5	20	Adjunctive Psychotherapy for Bipolar Disorder: A Systematic Review and Component Network Meta-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 141-150	14.5	35	
5	19	Initial severity of depression and efficacy of cognitive-behavioural therapy: individual-participant data meta-analysis of pill-placebo-controlled trials. <i>British Journal of Psychiatry</i> , 2017 , 210, 190-196	5.4	34	
5	18	Different Types and Acceptability of Psychotherapies for Acute Anxiety Disorders in Children and Adolescents: A Network Meta-analysis. <i>JAMA Psychiatry</i> , 2019 , 76, 41-50	14.5	34	
5	17	Impact of mental disorders on work performance in a community sample of workers in Japan: the World Mental Health Japan Survey 2002-2005. <i>Psychiatry Research</i> , 2012 , 198, 140-5	9.9	33	
5	16	Psychotic-like experiences are associated with violent behavior in adolescents. <i>Schizophrenia Research</i> , 2011 , 126, 245-51	3.6	33	
5	15	Cost of depression among adults in Japan in 2005. Psychiatry and Clinical Neurosciences, 2011, 65, 442-50	06.2	33	
5	14	Combined psychotherapy plus benzodiazepines for panic disorder. <i>The Cochrane Library</i> , 2009 , CD00533	3 5 .2	33	

513	Symptomatic recovery and social functioning in major depression. <i>Acta Psychiatrica Scandinavica</i> , 2001 , 103, 257-61	6.5	33
512	Discrepancies among megatrials. <i>Journal of Clinical Epidemiology</i> , 2000 , 53, 1193-9	5.7	33
511	Behavioral activation for late adolescents with subthreshold depression: a randomized controlled trial. <i>European Child and Adolescent Psychiatry</i> , 2016 , 25, 1171-1182	5.5	33
510	Social support and adjustment to a novel social environment. <i>International Journal of Social Psychiatry</i> , 1998 , 44, 56-70	8.5	32
509	Cultural invariance of likelihood ratios for the General Health Questionnaire. <i>Lancet, The</i> , 1999 , 353, 561-2	40	32
508	Long-term antipsychotic treatment in schizophrenia: systematic review and network meta-analysis of randomised controlled trials. <i>BJPsych Open</i> , 2016 , 2, 59-66	5	31
507	Group cognitive behavior therapy for Japanese patients with social anxiety disorder: preliminary outcomes and their predictors. <i>BMC Psychiatry</i> , 2007 , 7, 69	4.2	31
506	Reliability and validity of the Japanese version of the Frontal Assessment Battery in patients with the frontal variant of frontotemporal dementia. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 78-83	6.2	31
505	Does Internet-based cognitive behavioral therapy (iCBT) prevent major depressive episode for workers? A 12-month follow-up of a randomized controlled trial. <i>Psychological Medicine</i> , 2015 , 45, 1907	-69	30
504	Oncologists' recognition of supportive care needs and symptoms of their patients in a breast cancer outpatient consultation. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 1251-8	2.8	30
503	Prevalence of and risk factors for suicide-related outcomes in the World Health Organization World Mental Health Surveys Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2008 , 62, 442-9	6.2	30
502	Impairment of decision-making cognition in a case of frontotemporal lobar degeneration (FTLD) presenting with pathologic gambling and hoarding as the initial symptoms. <i>Cognitive and Behavioral Neurology</i> , 2007 , 20, 121-5	1.6	30
501	Initial severity of major depression and efficacy of new generation antidepressants: individual participant data meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 450-458	6.5	29
500	Optimising first- and second-line treatment strategies for untreated major depressive disorder - the SUN:)D study: a pragmatic, multi-centre, assessor-blinded randomised controlled trial. <i>BMC Medicine</i> , 2018 , 16, 103	11.4	29
499	Milnacipran versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2009 , CD006529	5.2	29
498	Cross-cultural evaluation of the Panic Disorder Severity Scale in Japan. <i>Depression and Anxiety</i> , 2004 , 20, 17-22	8.4	29
497	Childhood parental separation experiences and depressive symptomatology in acute major depression. <i>Psychiatry and Clinical Neurosciences</i> , 2003 , 57, 215-9	6.2	29
496	Review of dysthymia and persistent depressive disorder: history, correlates, and clinical implications. <i>Lancet Psychiatry,the</i> , 2020 , 7, 801-812	23.3	29

(2008-2018)

495	Distributional patterns of item responses and total scores on the PHQ-9 in the general population: data from the National Health and Nutrition Examination Survey. <i>BMC Psychiatry</i> , 2018 , 18, 108	4.2	28	
494	Etiologies of delirium and their relationship to reversibility and motor subtype in cancer patients. Japanese Journal of Clinical Oncology, 2009 , 39, 175-82	2.8	28	
493	Interoceptive hypersensitivity and interoceptive exposure in patients with panic disorder: specificity and effectiveness. <i>BMC Psychiatry</i> , 2006 , 6, 32	4.2	28	
492	Cohort Profile: The Tokyo Teen Cohort study (TTC). International Journal of Epidemiology, 2019 , 48, 141	4 7 18114	1 927	
491	Prevalence and correlates of illicit and non-medical use of psychotropic drugs in Japan: findings from the World Mental Health Japan Survey 2002-2004. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 777-83	4.5	27	
490	Course and outcome of depressive episodes: comparison between bipolar, unipolar and subthreshold depression. <i>Psychiatry Research</i> , 2000 , 96, 211-20	9.9	27	
489	Dismantling, optimising, and personalising internet cognitive behavioural therapy for depression: a systematic review and component network meta-analysis using individual participant data. <i>Lancet Psychiatry,the</i> , 2021 , 8, 500-511	23.3	27	
488	Help-seeking intention for depression in early adolescents: Associated factors and sex differences. Journal of Affective Disorders, 2018, 238, 359-365	6.6	26	
487	Fluvoxamine versus other anti-depressive agents for depression. <i>The Cochrane Library</i> , 2010 , CD006114	1 5.2	26	
486	Delirium Training Program for Nurses. <i>Psychosomatics</i> , 2010 , 51, 106-111	2.6	26	
485	Change in quality of life and their predictors in the long-term follow-up after group cognitive behavioral therapy for social anxiety disorder: a prospective cohort study. <i>BMC Psychiatry</i> , 2010 , 10, 81	4.2	26	
484	Factors correlated with fatigue in terminally ill cancer patients: a longitudinal study. <i>Journal of Pain and Symptom Management</i> , 2008 , 35, 515-23	4.8	26	
483	Combination of psychotherapy and benzodiazepines versus either therapy alone for panic disorder: a systematic review. <i>BMC Psychiatry</i> , 2007 , 7, 18	4.2	26	
482	Memory, attention, and executive functions before and after sine and pulse wave electroconvulsive therapies for treatment-resistant major depression. <i>Journal of ECT</i> , 2006 , 22, 107-12	2	26	
481	Psychosocial and psychological interventions for relapse prevention in schizophrenia: a systematic review and network meta-analysis. <i>Lancet Psychiatry,the</i> , 2021 , 8, 969-980	23.3	26	
480	Screening for depression among first-visit psychiatric patients: comparison of different scoring methods for the Center for Epidemiologic Studies Depression Scale using receiver operating characteristic analyses. <i>Psychiatry and Clinical Neurosciences</i> , 1997 , 51, 71-8	6.2	25	
479	Childhood parental loss and schizophrenia: evidence against pathogenic but for some pathoplastic effects. <i>Psychiatry Research</i> , 1998 , 81, 353-62	9.9	25	
478	Definitions of recovery and outcomes of major depression: results from a 10-year follow-up. <i>Acta Psychiatrica Scandinavica</i> , 2008 , 117, 35-40	6.5	25	

477	Depressive symptoms and depression in individuals with internet gaming disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2021 , 284, 136-142	6.6	25
476	Dissociation mediates the relationship between peer victimization and hallucinatory experiences among early adolescents. <i>Schizophrenia Research: Cognition</i> , 2016 , 4, 18-23	2.8	25
475	Stability of the Distribution of Patient Health Questionnaire-9 Scores Against Age in the General Population: Data From the National Health and Nutrition Examination Survey. <i>Frontiers in Psychiatry</i> , 2018 , 9, 390	5	25
474	Current status of research on cognitive therapy/cognitive behavior therapy in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2011 , 65, 121-9	6.2	24
473	Self-perceived burden in terminally ill cancer patients: a categorization of care strategies based on bereaved family members' perspectives. <i>Journal of Pain and Symptom Management</i> , 2010 , 40, 224-34	4.8	24
472	Regional cerebral blood flow changes in depression after electroconvulsive therapy. <i>Psychiatry Research - Neuroimaging</i> , 2006 , 147, 135-43	2.9	24
471	Psychotherapies for depression: a network meta-analysis covering efficacy, acceptability and long-term outcomes of all main treatment lypes. <i>World Psychiatry</i> , 2021 , 20, 283-293	14.4	24
470	Paroxetine versus other anti-depressive agents for depression. <i>The Cochrane Library</i> , 2014 , CD006531	5.2	23
469	Pre-post changes in psychosocial functioning among relatives of patients with depressive disorders after Brief Multifamily Psychoeducation: a pilot study. <i>BMC Psychiatry</i> , 2011 , 11, 56	4.2	23
468	Non-fearful vs. fearful panic attacks: a general population study from the National Comorbidity Survey. <i>Journal of Affective Disorders</i> , 2009 , 112, 273-8	6.6	23
467	Patient-perceived barriers to the psychological care of Japanese patients with lung cancer. Japanese Journal of Clinical Oncology, 2008 , 38, 653-60	2.8	23
466	Reliability and validity of the Japanese version of the Dysexecutive Questionnaire (DEX) in Alzheimer's disease: validation of a behavioral rating scale to assess dysexecutive symptoms in Japanese patients with Alzheimer's disease. <i>International Journal of Geriatric Psychiatry</i> , 2007 , 22, 951-6	3.9 5	23
465	Evaluation of diagnostic criteria for panic attack using item response theory: findings from the National Comorbidity Survey in USA. <i>Journal of Affective Disorders</i> , 2007 , 104, 197-201	6.6	23
464	Time for united action on depression: a Lancet-World Psychiatric Association Commission <i>Lancet, The,</i> 2022 , 399, 957-1022	40	23
463	Impact of placebo arms on outcomes in antidepressant trials: systematic review and meta-regression analysis. <i>International Journal of Epidemiology</i> , 2018 , 47, 1454-1464	7.8	22
462	Relative indices of treatment effect may be constant across different definitions of response in schizophrenia trials. <i>Schizophrenia Research</i> , 2011 , 126, 212-9	3.6	22
461	Depression, alexithymia and long-term mortality in chronic hemodialysis patients. <i>Psychotherapy and Psychosomatics</i> , 2010 , 79, 303-11	9.4	22
460	Low dosage tricyclic antidepressants for depression. <i>The Cochrane Library</i> , 2003 , CD003197	5.2	22

459	Clinimetric Criteria for Patient-Reported Outcome Measures. <i>Psychotherapy and Psychosomatics</i> , 2021 , 90, 222-232	9.4	22
458	Network Meta-analysis: Users' Guide for Surgeons: Part I - Credibility. <i>Clinical Orthopaedics and Related Research</i> , 2015 , 473, 2166-71	2.2	21
457	Prediction intervals for random-effects meta-analysis: A confidence distribution approach. <i>Statistical Methods in Medical Research</i> , 2019 , 28, 1689-1702	2.3	21
456	Psychotropic prescription patterns among patients diagnosed with depressive disorder based on claims database in Japan. <i>Clinical Drug Investigation</i> , 2013 , 33, 597-605	3.2	21
455	Mapping hypofrontality during letter fluency task in schizophrenia: a multi-channel near-infrared spectroscopy study. <i>Schizophrenia Research</i> , 2012 , 136, 63-9	3.6	21
454	Cognitive behavior therapy for psychological distress in patients with recurrent miscarriage. <i>Psychology Research and Behavior Management</i> , 2013 , 6, 37-43	3.8	21
453	Anticipatory nausea among ambulatory cancer patients undergoing chemotherapy: prevalence, associated factors, and impact on quality of life. <i>Cancer Science</i> , 2010 , 101, 2596-600	6.9	21
452	Efficacy and tolerability of milnacipran in the treatment of major depression in comparison with other antidepressants: a systematic review and meta-analysis. <i>CNS Drugs</i> , 2008 , 22, 587-602	6.7	21
451	Association between apathy/depression and executive function in patients with Alzheimer's disease. <i>International Psychogeriatrics</i> , 2008 , 20, 964-75	3.4	21
450	Mental health literacy in Japanese cancer patients: ability to recognize depression and preferences of treatments-comparison with Japanese lay public. <i>Psycho-Oncology</i> , 2007 , 16, 834-42	3.9	21
449	Greater impairment of ability in the divided attention task is seen in Alzheimer's disease patients with depression than in those without depression. <i>Dementia and Geriatric Cognitive Disorders</i> , 2007 , 23, 231-40	2.6	21
448	Is "relationship fear" unique to Japan? Symptom factors and patient clusters of social anxiety disorder among the Japanese clinical population. <i>Journal of Affective Disorders</i> , 2005 , 87, 131-7	6.6	21
447	Treatment received by depressed patients in Japan and its determinants: naturalistic observation from a multi-center collaborative follow-up study. <i>Journal of Affective Disorders</i> , 2000 , 60, 173-9	6.6	21
446	Antidepressants plus benzodiazepines for adults with major depression. <i>The Cochrane Library</i> , 2019 , 6, CD001026	5.2	21
445	Individual Differences in Response to Antidepressants: A Meta-analysis of Placebo-Controlled Randomized Clinical Trials. <i>JAMA Psychiatry</i> , 2020 , 77, 607-617	14.5	20
444	Is placebo response in antidepressant trials rising or not? A reanalysis of datasets to conclude this long-lasting controversy. <i>Evidence-Based Mental Health</i> , 2018 , 21, 1-3	11.1	20
443	Initial symptom severity of bipolar I disorder and the efficacy of olanzapine: a meta-analysis of individual participant data from five placebo-controlled studies. <i>Lancet Psychiatry,the</i> , 2017 , 4, 859-867	23.3	20
442	Effects of an internet-based cognitive behavioral therapy intervention on improving work engagement and other work-related outcomes: an analysis of secondary outcomes of a randomized controlled trial. <i>Journal of Occupational and Environmental Medicine</i> , 2015 , 57, 578-84	2	20

441	Cost-effectiveness of family psychoeducation to prevent relapse in major depression: results from a randomized controlled trial. <i>BMC Psychiatry</i> , 2012 , 12, 40	4.2	20
440	Help-seeking behavior among Japanese school students who self-harm: results from a self-report survey of 18,104 adolescents. <i>Neuropsychiatric Disease and Treatment</i> , 2012 , 8, 561-9	3.1	20
439	Strategic use of new generation antidepressants for depression: SUN(^_^)D study protocol. <i>Trials</i> , 2011 , 12, 116	2.8	20
438	Efficacy, tolerability and side-effect profile of fluvoxamine for major depression: meta-analysis. <i>Journal of Psychopharmacology</i> , 2009 , 23, 539-50	4.6	20
437	Psychosocial factors and survival after diagnosis of inoperable non-small cell lung cancer. <i>Psycho-Oncology</i> , 2009 , 18, 23-9	3.9	20
436	Evaluation of the DSM-IV and ICD-10 criteria for depressive disorders in a community population in Japan using item response theory. <i>International Journal of Methods in Psychiatric Research</i> , 2010 , 19, 211-22	4.3	20
435	An acidic fibroblast growth factor-like factor secreted into the brain cell culture medium upregulates apoE synthesis, HDL secretion and cholesterol metabolism in rat astrocytes. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2002 , 1589, 261-72	4.9	20
434	Does randomized evidence support sertraline as first-line antidepressant for adults with acute major depression? A systematic review and meta-analysis. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1732-4	1 2 .6	20
433	Sojourner readjustment: mental health of international students after one year's foreign sojourn and its psychosocial correlates. <i>Journal of Nervous and Mental Disease</i> , 1997 , 185, 263-8	1.8	20
432	Detailed course of depressive symptoms and risk for developing depression in late adolescents with subthreshold depression: a cohort study. <i>Neuropsychiatric Disease and Treatment</i> , 2017 , 13, 25-33	3.1	20
431	Translating the BDI and BDI-II into the HAMD and vice versa with equipercentile linking. <i>Epidemiology and Psychiatric Sciences</i> , 2019 , 29, e24	5.1	20
430	Distribution of psychological distress is stable in recent decades and follows an exponential pattern in the US population. <i>Scientific Reports</i> , 2019 , 9, 11982	4.9	19
429	A mindfulness-based stress management program and treatment with omega-3 fatty acids to maintain a healthy mental state in hospital nurses (Happy Nurse Project): study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 36	2.8	19
428	Protocol registration and selective outcome reporting in recent psychiatry trials: new antidepressants and cognitive behavioural therapies. <i>Acta Psychiatrica Scandinavica</i> , 2015 , 132, 489-98	6.5	19
427	Long-term treatment of depression with antidepressants: a systematic narrative review. <i>Canadian Journal of Psychiatry</i> , 2007 , 52, 545-52	4.8	19
426	Independent associations of alexithymia and social support with depression in hemodialysis patients. <i>Journal of Psychosomatic Research</i> , 2007 , 63, 349-56	4.1	19
425	Psychotherapy for depression in children and adolescents. <i>The Cochrane Library</i> , 2004 ,	5.2	19
424	Stratum-specific likelihood ratios of the general health questionnaire in the community: help-seeking and physical co-morbidity affect the test characteristics. <i>Psychological Medicine</i> , 2002 , 32, 743-8	6.9	19

(2015-2015)

423	Right tail of the distribution of depressive symptoms is stable and follows an exponential curve during middle adulthood. <i>PLoS ONE</i> , 2015 , 10, e0114624	3.7	19	
422	Effect of Recruitment Methods on Response Rate in a Web-Based Study for Primary Care Physicians: Factorial Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2018 , 20, e28	7.6	19	
421	A greater number of somatic pain sites is associated with poor mental health in adolescents: a cross-sectional study. <i>BMC Psychiatry</i> , 2013 , 13, 30	4.2	18	
420	Psychoeducation for major depressive disorders: a randomised controlled trial. <i>Psychiatry Research</i> , 2013 , 210, 134-9	9.9	18	
419	Intra-individual versus extra-individual components of social support. <i>Psychological Medicine</i> , 1997 , 27, 1183-91	6.9	18	
418	Cultural distance and its relationship to psychological adjustment of international exchange students. <i>Psychiatry and Clinical Neurosciences</i> , 1997 , 51, 87-91	6.2	18	
417	Multifaceted psychosocial intervention program for breast cancer patients after first recurrence: feasibility study. <i>Psycho-Oncology</i> , 2007 , 16, 517-24	3.9	18	
416	Enuresis and Hyperactivity-Inattention in Early Adolescence: Findings from a Population-Based Survey in Tokyo (Tokyo Early Adolescence Survey). <i>PLoS ONE</i> , 2016 , 11, e0158786	3.7	18	
415	Trends in antipsychotic prescriptions for Japanese outpatients during 2006-2012: a descriptive epidemiological study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 642-656	2.6	17	
414	Mechanisms of behavioral activation for late adolescents: Positive reinforcement mediate the relationship between activation and depressive symptoms from pre-treatment to post-treatment. <i>Journal of Affective Disorders</i> , 2016 , 204, 70-3	6.6	17	
413	Remifentanil in electroconvulsive therapy: a systematic review and meta-analysis of randomized controlled trials. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2016 , 266, 703-717	5.1	17	
412	A distribution model of the responses to each depressive symptom item in a general population: a cross-sectional study. <i>BMJ Open</i> , 2015 , 5, e008599	3	17	
411	Videotaped experiments to drop safety behaviors and self-focused attention for patients with social anxiety disorder: do they change subjective and objective evaluations of anxiety and performance?. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2009 , 40, 202-10	2.6	17	
410	Efficacy of olanzapine augmentation of paroxetine therapy in patients with severe body dysmorphic disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2008 , 62, 370	6.2	17	
409	SPECT-identified neuroanatomical predictor of the cognitive effects of donepezil treatment in patients with Alzheimer's disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2008 , 26, 556-66	2.6	17	
408	Lithium Levels in Tap Water and the Mental Health Problems of Adolescents: An Individual-Level Cross-Sectional Survey. <i>Journal of Clinical Psychiatry</i> , 2017 , 78, e252-e256	4.6	17	
407	Brief mindfulness-based stress management program for a better mental state in working populations - Happy Nurse Project: A randomized controlled trial. <i>Journal of Affective Disorders</i> , 2019 , 251, 186-194	6.6	16	
406	Cost-effectiveness of cognitive behavioral therapy for insomnia comorbid with depression: Analysis of a randomized controlled trial. <i>Psychiatry and Clinical Neurosciences</i> , 2015 , 69, 335-43	6.2	16	

405	Imputation of response rates from means and standard deviations in schizophrenia. <i>Schizophrenia Research</i> , 2013 , 151, 209-14	3.6	16
404	Adding smartphone-based cognitive-behavior therapy to pharmacotherapy for major depression (FLATT project): study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 293	2.8	16
403	Intertextuality, mediation, and members lategories in focus groups on humor. <i>Pragmatics and Society</i> , 2010 , 1, 257-283	0.5	16
402	Typus melancholicus in light of the five-factor model of personality. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1998 , 248, 64-9	5.1	16
401	Cross-cultural study of conviction subtype Taijin Kyofu: proposal and reliability of Nagoya-Osaka diagnostic criteria for social anxiety disorder. <i>Journal of Nervous and Mental Disease</i> , 2008 , 196, 307-13	1.8	16
400	Diagnostic agreement in psychiatry. A comparative study between ICD-9, ICD-10 and DSM-III-R. <i>British Journal of Psychiatry</i> , 1993 , 162, 621-6	5.4	16
399	Psychological interventions to prevent the onset of depressive disorders: A meta-analysis of randomized controlled trials. <i>Clinical Psychology Review</i> , 2021 , 83, 101955	10.8	16
398	The Need for Research on Treatments of Chronic Depression. <i>JAMA Psychiatry</i> , 2017 , 74, 242-243	14.5	15
397	Comparative efficacy and acceptability of antidepressants in the long-term treatment of major depression: protocol for a systematic review and networkmeta-analysis. <i>BMJ Open</i> , 2019 , 9, e027574	3	15
396	Exploratory analyses of effect modifiers in the antidepressant treatment of major depression: Individual-participant data meta-analysis of 2803 participants in seven placebo-controlled randomized trials. <i>Journal of Affective Disorders</i> , 2019 , 250, 419-424	6.6	15
395	Antidepressants and benzodiazepines for panic disorder in adults. <i>The Cochrane Library</i> , 2016 , 9, CD011	596.7	15
394	Second-generation antipsychotic drugs and short-term somatic serious adverse events: a systematic review and meta-analysis. <i>Lancet Psychiatry,the</i> , 2019 , 6, 753-765	23.3	15
393	Prescription patterns following first-line new generation antidepressants for depression in Japan: a naturalistic cohort study based on a large claims database. <i>Journal of Affective Disorders</i> , 2013 , 150, 916	;- <u>6.6</u> ;- 22	15
392	New methods can extend the use of minimal important difference units in meta-analyses of continuous outcome measures. <i>Journal of Clinical Epidemiology</i> , 2012 , 65, 817-26	5.7	15
391	Neuroanatomical abnormalities before onset of delusions in patients with Alzheimer's disease: a voxel-based morphometry study. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 1-8	3.1	15
390	Predicting maladjustment of exchange students in different cultures: a prospective study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1993 , 28, 142-6	4.5	15
389	Cognitive and Behavioral Skills Exercises Completed by Patients with Major Depression During Smartphone Cognitive Behavioral Therapy: Secondary Analysis of a Randomized Controlled Trial. <i>JMIR Mental Health</i> , 2018 , 5, e4	6	15
388	Pattern analysis of total item score and item response of the Kessler Screening Scale for Psychological Distress (K6) in a nationally representative sample of US adults. <i>PeerJ</i> , 2017 , 5, e2987	3.1	15

(2005-2018)

387	Smartphone problem-solving and behavioural activation therapy to reduce fear of recurrence among patients with breast cancer (SMartphone Intervention to LEssen fear of cancer recurrence: SMILE project): protocol for a randomised controlled trial. <i>BMJ Open</i> , 2018 , 8, e024794	3	15
386	Quality of randomized controlled trials of new generation antidepressants and antipsychotics identified in the China National Knowledge Infrastructure (CNKI): a literature and telephone interview study. <i>BMC Medical Research Methodology</i> , 2018 , 18, 96	4.7	15
385	Initial severity and efficacy of risperidone in autism: Results from the RUPP trial. <i>European Psychiatry</i> , 2016 , 32, 16-20	6	14
384	Lifetime comorbidities between phobic disorders and major depression in Japan: results from the World Mental Health Japan 2002-2004 Survey. <i>Depression and Anxiety</i> , 2009 , 26, 949-55	8.4	14
383	Cognitive-behavior therapy for Japanese patients with panic disorder: acute phase and one-year follow-up results. <i>Psychiatry and Clinical Neurosciences</i> , 2008 , 62, 313-21	6.2	14
382	Loss of fractal heart rate dynamics in depressive hemodialysis patients. <i>Psychosomatic Medicine</i> , 2008 , 70, 177-85	3.7	14
381	Validity of brief intelligence tests for patients with Alzheimer's disease. <i>Psychiatry and Clinical Neurosciences</i> , 2000 , 54, 435-9	6.2	14
3 80	Adipokines in anorexia nervosa: A systematic review and meta-analysis. <i>Psychoneuroendocrinology</i> , 2020 , 112, 104485	5	14
379	Measurement equivalence of the Social Interaction Anxiety Scale (SIAS) and Social Phobia Scale (SPS) across individuals with social anxiety disorder from Japanese and Australian sociocultural contexts. <i>Journal of Affective Disorders</i> , 2019 , 243, 165-174	6.6	14
378	Personalise antidepressant treatment for unipolar depression combining individual choices, risks and big data (PETRUSHKA): rationale and protocol. <i>Evidence-Based Mental Health</i> , 2020 , 23, 52-56	11.1	13
377	Specificity of CBT for Depression: A Contribution from Multiple Treatments Meta-analyses. <i>Cognitive Therapy and Research</i> , 2014 , 38, 249-260	2.7	13
376	Possible improvement of depression after systematic examination and explanation of live birth rates among women with recurrent miscarriage. <i>Journal of Obstetrics and Gynaecology</i> , 2013 , 33, 171-4	1.3	13
375	Ictal physiological characteristics of remitters during bilateral electroconvulsive therapy. <i>Psychiatry Research</i> , 2011 , 185, 462-4	9.9	13
374	Typus melancholicus is not the premorbid personality trait of unipolar (endogenous) depression. <i>Psychiatry and Clinical Neurosciences</i> , 1997 , 51, 197-202	6.2	13
373	Childhood parental loss and alcohol dependence among Japanese men: a case-control study. Group for Longitudinal Affective Disorders Study (GLADS). <i>Acta Psychiatrica Scandinavica</i> , 1998 , 97, 403-7	6.5	13
372	Problem-solving therapy for psychological distress in Japanese cancer patients: preliminary clinical experience from psychiatric consultations. <i>Japanese Journal of Clinical Oncology</i> , 2008 , 38, 867-70	2.8	13
371	Duloxetine versus other anti-depressive agents for depression 2007,		13
370	Quality of life and social role functioning in Japanese patients with panic disorder. <i>Journal of Nervous and Mental Disease</i> , 2005 , 193, 686-9	1.8	13

369	Relationship between EEG dimensional complexity and neuropsychological findings in Alzheimer's disease. <i>Psychiatry and Clinical Neurosciences</i> , 2000 , 54, 537-41	6.2	13
368	Perceived parental rearing, personality and mental status in Japanese adolescents. <i>Journal of Adolescence</i> , 1992 , 15, 317-22	3.4	13
367	Response rates in patients with schizophrenia and positive symptoms receiving cognitive behavioural therapy: a systematic review and single-group meta-analysis. <i>BMC Psychiatry</i> , 2018 , 18, 380	4.2	13
366	Although not consistently superior, the absolute approach to framing the minimally important difference has advantages over the relative approach. <i>Journal of Clinical Epidemiology</i> , 2015 , 68, 888-94	. 5.7	12
365	Exponential distribution of total depressive symptom scores in relation to exponential latent trait and item threshold distributions: a simulation study. <i>BMC Research Notes</i> , 2017 , 10, 614	2.3	12
364	Antidepressants might work for people with major depression: where do we go from here?. <i>Lancet Psychiatry,the</i> , 2018 , 5, 461-463	23.3	12
363	Video feedback with peer ratings in naturalistic anxiety-provoking situations for social anxiety disorder: Preliminary report. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2010 , 41, 6-10	2.6	12
362	Premorbid personality traits of patients with organic (ICD-10 F0), schizophrenic (F2), mood (F3), and neurotic (F4) disorders according to the five-factor model of personality. <i>Psychiatry Research</i> , 1998 , 78, 179-87	9.9	12
361	Can pill placebo augment cognitive-behavior therapy for panic disorder?. BMC Psychiatry, 2007, 7, 73	4.2	12
360	Two dimensions of anosognosia in patients with Alzheimer's disease: reliability and validity of the Japanese version of the Anosognosia Questionnaire for Dementia (AQ-D). <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 672-7	6.2	12
359	Relationship between quality of reports of antidepressant randomized controlled trials and treatment estimates: systematic review, meta-analysis, and meta-regression analysis. <i>Journal of Clinical Psychopharmacology</i> , 2007 , 27, 352-6	1.7	12
358	Distribution of Total Depressive Symptoms Scores and Each Depressive Symptom Item in a Sample of Japanese Employees. <i>PLoS ONE</i> , 2016 , 11, e0147577	3.7	12
357	Maternal diabetes in early pregnancy, and psychotic experiences and depressive symptoms in 10-year-old offspring: A population-based birth cohort study. <i>Schizophrenia Research</i> , 2019 , 206, 52-57	3.6	12
356	Network Meta-analysis: Users' Guide for Surgeons: Part II - Certainty. <i>Clinical Orthopaedics and Related Research</i> , 2015 , 473, 2172-8	2.2	11
355	Characteristic distribution of the total and individual item scores on the Kessler Screening Scale for Psychological Distress (K6) in US adults. <i>BMC Psychiatry</i> , 2017 , 17, 290	4.2	11
354	Bartlett-type corrections and bootstrap adjustments of likelihood-based inference methods for network meta-analysis. <i>Statistics in Medicine</i> , 2018 , 37, 1178-1190	2.3	11
353	Antidepressants versus placebo for panic disorder in adults. <i>The Cochrane Library</i> , 2018 , 4, CD010676	5.2	11
352	Psychological therapies versus pharmacological interventions for panic disorder with or without agoraphobia in adults. <i>The Cochrane Library</i> , 2016 , 10, CD011170	5.2	11

351	Using the contribution matrix to evaluate complex study limitations in a network meta-analysis: a case study of bipolar maintenance pharmacotherapy review. <i>BMC Research Notes</i> , 2016 , 9, 218	2.3	11
350	Healthy Campus Trial: a multiphase optimization strategy (MOST) fully factorial trial to optimize the smartphone cognitive behavioral therapy (CBT) app for mental health promotion among university students: study protocol for a randomized controlled trial. <i>Trials</i> , 2018 , 19, 353	2.8	11
349	Comparative efficacy and acceptability of antidepressants, psychological interventions, and their combination for depressive disorder in children and adolescents: protocol for a network meta-analysis. <i>BMJ Open</i> , 2017 , 7, e016608	3	11
348	Anxiety sensitivity as a predictor of broad dimensions of psychopathology after cognitive behavioral therapy for panic disorder. <i>Neuropsychiatric Disease and Treatment</i> , 2017 , 13, 1835-1840	3.1	11
347	Neural basis of three dimensions of agitated behaviors in patients with Alzheimer disease. Neuropsychiatric Disease and Treatment, 2014 , 10, 339-48	3.1	11
346	Change in quality of life after brief behavioral therapy for insomnia in concurrent depression: analysis of the effects of a randomized controlled trial. <i>Journal of Clinical Sleep Medicine</i> , 2014 , 10, 433-	.93.1	11
345	Ranking antidepressants [Authors' reply. Lancet, The, 2009, 373, 1761-1762	40	11
344	Sources of bias in diagnostic accuracy studies and the diagnostic process. <i>Cmaj</i> , 2006 , 174, 481-2	3.5	11
343	Validity of indirect comparisons in meta-analysis. <i>Lancet, The</i> , 2007 , 369, 270-1; author reply 271	40	11
342	Factors influencing adjustment of high school students in an international exchange program. Journal of Nervous and Mental Disease, 1994 , 182, 709-14	1.8	11
341	Relationship between Item Responses of Negative Affect Items and the Distribution of the Sum of the Item Scores in the General Population. <i>PLoS ONE</i> , 2016 , 11, e0165928	3.7	11
340	Cognitive-Behavioural Analysis System of Psychotherapy (CBASP), a drug, or their combination: differential therapeutics for persistent depressive disorder: a study protocol of an individual participant data network meta-analysis. <i>BMJ Open</i> , 2016 , 6, e011769	3	11
339	Dismantling, personalising and optimising internet cognitive-behavioural therapy for depression: a study protocol for individual participant data component network meta-analysis. <i>BMJ Open</i> , 2019 , 8, e026137	3	11
338	Comparative efficacy and tolerability of 32 oral and long-acting injectable antipsychotics for the maintenance treatment of adults with schizophrenia: a systematic review and network meta-analysis <i>Lancet, The</i> , 2022 , 399, 824-836	40	11
337	Overstatements in abstract conclusions claiming effectiveness of interventions in psychiatry: A meta-epidemiological investigation. <i>PLoS ONE</i> , 2017 , 12, e0184786	3.7	10
336	Omega-3 fatty acids for a better mental state in working populations - Happy Nurse Project: A 52-week randomized controlled trial. <i>Journal of Psychiatric Research</i> , 2018 , 102, 72-80	5.2	10
335	Maternal exposure to benzodiazepine and risk of preterm birth and low birth weight: A case-control study using a claims database in Japan. <i>Asia-Pacific Psychiatry</i> , 2018 , 10, e12309	3.2	10
334	Azapirones versus placebo for panic disorder in adults. <i>The Cochrane Library</i> , 2014 , CD010828	5.2	10

333	A failure to confirm the effectiveness of a brief group psychoeducational program for mothers of children with high-functioning pervasive developmental disorders: a randomized controlled pilot trial. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 1141-53	3.1	10
332	The first 100 patients in the SUN(^_^)D trial (strategic use of new generation antidepressants for depression): examination of feasibility and adherence during the pilot phase. <i>Trials</i> , 2012 , 13, 80	2.8	10
331	What is a Cochrane review?. Epidemiology and Psychiatric Sciences, 2011, 20, 231-3	5.1	10
330	Supported employment for adults with severe mental illness. <i>Cochrane Database of Systematic Reviews</i> , 2010 , 2010,		10
329	Postictal cardiovascular response predicts therapeutic efficacy of electroconvulsive therapy for depression. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 290-4	6.2	10
328	Mirtazapine versus other anti-depressive agents for depression 2007,		10
327	Citalopram versus other anti-depressive agents for depression 2007,		10
326	A case of late-onset obsessive compulsive disorder developing frontotemporal lobar degeneration. Journal of Neuropsychiatry and Clinical Neurosciences, 2007 , 19, 487-8	2.7	10
325	Remission rates with milnacipran 100 mg/day and 150 mg/day in the long-term treatment of major depression. <i>Clinical Drug Investigation</i> , 2006 , 26, 135-42	3.2	10
324	Efficacy of diazepam as an anti-anxiety agent: meta-analysis of double-blind, randomized controlled trials carried out in Japan. <i>Human Psychopharmacology</i> , 2003 , 18, 483-7	2.3	10
323	Effect of adjunctive vitamin C, glucocorticoids, and vitamin B1 on longer-term mortality in adults with sepsis or septic shock: a systematic review and a component network meta-analysis. <i>Intensive Care Medicine</i> , 2021 , 48, 16	14.5	10
322	Substance use improvement depends on Race/Ethnicity: Outpatient treatment disparities observed in a large US national sample. <i>Drug and Alcohol Dependence</i> , 2020 , 213, 108087	4.9	10
321	The effects of psychotherapies for depression on response, remission, reliable change, and deterioration: A meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 144, 288-299	6.5	10
320	Quantifying indirect evidence in network meta-analysis. <i>Statistics in Medicine</i> , 2017 , 36, 917-927	2.3	9
319	Side effect profile and comparative tolerability of 21 antidepressants in the acute treatment of major depression in adults: protocol for a network meta-analysis. <i>Evidence-Based Mental Health</i> , 2019 , 22, 61-66	11.1	9
318	Unguided Chatbot-Delivered Cognitive Behavioural Intervention for Problem Gamblers Through Messaging App: A Randomised Controlled Trial. <i>Journal of Gambling Studies</i> , 2020 , 36, 1391-1407	3	9
317	Distribution of item responses and total item scores for the Center for Epidemiologic Studies Depression Scale (CES-D): Data from the Irish Longitudinal Study on Ageing (TILDA). <i>PLoS ONE</i> , 2018 , 13, e0202607	3.7	9
316	Behavioral activation: Is it the expectation or achievement, of mastery or pleasure that contributes to improvement in depression?. <i>Journal of Affective Disorders</i> , 2018 , 238, 336-341	6.6	9

315	Benzodiazepines versus placebo for panic disorder in adults. <i>The Cochrane Library</i> , 2019 , 3, CD010677	5.2	9
314	Duration of untreated psychosis in a rural/suburban region of Japan. <i>Microbial Biotechnology</i> , 2012 , 6, 239-46	3.3	9
313	Prevalence of depression among outpatients visiting a general internal medicine polyclinic in rural Japan. <i>General Hospital Psychiatry</i> , 2013 , 35, 286-90	5.6	9
312	Brain structural abnormalities in behavior therapy-resistant obsessive-compulsive disorder revealed by voxel-based morphometry. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 1987-96	3.1	9
311	Deliberate self-harm in adolescents aged 12-18: a cross-sectional survey of 18,104 students. <i>Suicide and Life-Threatening Behavior</i> , 2012 , 42, 550-60	3.9	9
310	Decreased white matter integrity before the onset of delusions in patients with Alzheimer's disease: diffusion tensor imaging. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 25-9	3.1	9
309	Cancer patients' reluctance to discuss psychological distress with their physicians was not associated with underrecognition of depression by physicians: a preliminary study. <i>Palliative and Supportive Care</i> , 2009 , 7, 229-33	2.5	9
308	How many well vs. unwell days can you expect over 10 years, once you become depressed?. <i>Acta Psychiatrica Scandinavica</i> , 2009 , 119, 290-7	6.5	9
307	10-year course of social adjustment in major depression. <i>International Journal of Social Psychiatry</i> , 2011 , 57, 501-8	8.5	9
306	Panic disorder. <i>Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn</i> , 2012 , 106, 363-74	3	9
305	Cognitive behavioural therapies versus treatment as usual for depression. The Cochrane Library,		
	2010,	5.2	9
304		5.2 3.1	9
304	Absence status associated with focal activity and polydipsia-induced hyponatremia.		
	Absence status associated with focal activity and polydipsia-induced hyponatremia. Neuropsychiatric Disease and Treatment, 2008, 4, 495-8	3.1	9
303	Absence status associated with focal activity and polydipsia-induced hyponatremia. Neuropsychiatric Disease and Treatment, 2008, 4, 495-8 Coping behavior in depressed patients: a longitudinal study. Psychiatry Research, 2003, 121, 169-77 A cluster-analytically derived subtyping of chronic affective disorders. Acta Psychiatrica	3.1 9.9	9
303	Absence status associated with focal activity and polydipsia-induced hyponatremia. Neuropsychiatric Disease and Treatment, 2008, 4, 495-8 Coping behavior in depressed patients: a longitudinal study. Psychiatry Research, 2003, 121, 169-77 A cluster-analytically derived subtyping of chronic affective disorders. Acta Psychiatrica Scandinavica, 1992, 85, 177-82 Age-related changes in the distributions of depressive symptom items in the general population: a	3.19.96.5	9 9
303 302 301	Absence status associated with focal activity and polydipsia-induced hyponatremia. Neuropsychiatric Disease and Treatment, 2008, 4, 495-8 Coping behavior in depressed patients: a longitudinal study. Psychiatry Research, 2003, 121, 169-77 A cluster-analytically derived subtyping of chronic affective disorders. Acta Psychiatrica Scandinavica, 1992, 85, 177-82 Age-related changes in the distributions of depressive symptom items in the general population: a cross-sectional study using the exponential distribution model. PeerJ, 2016, 4, e1547 Association between antidepressant use during pregnancy and autism spectrum disorder in children: a retrospective cohort study based on Japanese claims data. Maternal Health, Neonatology	3.1 9.9 6.5 3.1	9 9 9

297	Different control conditions can produce different effect estimates in psychotherapy trials for depression. <i>Journal of Clinical Epidemiology</i> , 2021 , 132, 59-70	5.7	9
296	Psychological and Educational Interventions for Atopic Dermatitis in Adults: A Systematic Review and Meta-Analysis. <i>Behaviour Change</i> , 2017 , 34, 48-65	1.1	8
295	Predicting relapse in major depression after successful initial pharmacological treatment. <i>Journal of Affective Disorders</i> , 2019 , 250, 108-113	6.6	8
294	COVID-19 randomized controlled trials in medRxiv and PubMed. <i>European Journal of Internal Medicine</i> , 2020 , 81, 97-99	3.9	8
293	Differences in the placebo response in duloxetine and venlafaxine trials. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 472-480	6.5	8
292	Comparative efficacy and acceptability of antiepileptic drugs for classical trigeminal neuralgia: a Bayesian network meta-analysis protocol. <i>BMJ Open</i> , 2018 , 8, e017392	3	8
291	Influence of overstated abstract conclusions on clinicians: a web-based randomised controlled trial. <i>BMJ Open</i> , 2017 , 7, e018355	3	8
290	Ipsative imputation for a 15-item Geriatric Depression Scale in community-dwelling elderly people. <i>Psychogeriatrics</i> , 2014 , 14, 182-7	1.8	8
289	Physical pain associated with depression: results of a survey in Japanese patients and physicians. <i>Comprehensive Psychiatry</i> , 2012 , 53, 843-9	7.3	8
288	A semi-structured clinical interview for psychosis sub-groups (SCIPS): development and psychometric properties. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012 , 47, 563-80	4.5	8
287	Brief PANSS to assess and monitor the overall severity of schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2010 , 64, 262-7	6.2	8
286	Depressive symptoms among international exchange students, and their predictors. <i>Acta Psychiatrica Scandinavica</i> , 1997 , 96, 242-6	6.5	8
285	Spontaneous emission control by 17 layers of three-dimensional photonic crystals. <i>Electronics Letters</i> , 2008 , 44, 377	1.1	8
284	Impact of depression on insight into memory capacity in patients with Alzheimer disease. <i>Alzheimer Disease and Associated Disorders</i> , 2008 , 22, 369-74	2.5	8
283	Sensitization of catastrophic cognition in cognitive-behavioral therapy for panic disorder. <i>BMC Psychiatry</i> , 2007 , 7, 70	4.2	8
282	Ictal electroencephalographic correlates of posttreatment neuropsychological changes in electroconvulsive therapy: a hypothesis-generation study. <i>Journal of ECT</i> , 2007 , 23, 163-8	2	8
281	Sertraline versus other anti-depressive agents for depression. The Cochrane Library, 2006,	5.2	8
280	Antidepressants and suicide risk. British Journal of Psychiatry, 2001, 178, 477	5.4	8

279	Personality, social support, coping and emotional distress after repeated spontaneous abortions. Journal of Psychosomatic Obstetrics and Gynaecology, 2002 , 23, 133-40	3.6	8
278	Psychosocial factors regulating natural-killer cell activity in recurrent spontaneous abortions. <i>American Journal of Reproductive Immunology</i> , 2000 , 44, 299-302	3.8	8
277	Predictive validity of subtypes of chronic affective disorders derived by cluster analysis. <i>Acta Psychiatrica Scandinavica</i> , 1995 , 91, 379-85	6.5	8
276	All clinical trials must be reported in detail and made publicly available. <i>BMJ, The</i> , 2004 , 329, 626	5.9	8
275	Delirium training program for nurses. <i>Psychosomatics</i> , 2010 , 51, 106-11	2.6	8
274	Boundary curves of individual items in the distribution of total depressive symptom scores approximate an exponential pattern in a general population. <i>PeerJ</i> , 2016 , 4, e2566	3.1	8
273	Outlier detection and influence diagnostics in network meta-analysis. <i>Research Synthesis Methods</i> , 2020 , 11, 891-902	7.2	8
272	Individual Differences in Response to Antidepressants: A Meta-analysis of Placebo-Controlled Randomized Clinical Trials. <i>JAMA Psychiatry</i> , 2021 , 78, 490-497	14.5	8
271	Cognitive restructuring, behavioral activation and cognitive-behavioral therapy in the treatment of adult depression: A network meta-analysis. <i>Journal of Consulting and Clinical Psychology</i> , 2021 , 89, 563-5	5945	8
270	Predicting recurrence of depression using lifelog data: an explanatory feasibility study with a panel VAR approach. <i>BMC Psychiatry</i> , 2019 , 19, 391	4.2	8
269	Minimal important difference estimates for patient-reported outcomes: A systematic survey. Journal of Clinical Epidemiology, 2021 , 133, 61-71	5.7	8
268	Evidence synthesis, practice guidelines and real-world prescriptions of new generation antidepressants in the treatment of depression: a protocol for cumulative network meta-analyses and meta-epidemiological study. <i>BMJ Open</i> , 2018 , 8, e023222	3	8
267	Sex differences in auditory verbal hallucinations in early, middle and late adolescence: results from a survey of 17 451 Japanese students aged 12-18 years. <i>BMJ Open</i> , 2017 , 7, e015239	3	7
266	Quantity and quality of psychotherapy trials for depression in the past five decades. <i>Journal of Affective Disorders</i> , 2014 , 165, 190-5	6.6	7
265	Cognitive-behavioral therapy modifies the naturalistic course of social anxiety disorder: findings from an ABA design study in routine clinical practices. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 139-47	6.2	7
264	Item Response Patterns on the Patient Health Questionnaire-8 in a Nationally Representative Sample of US Adults. <i>Frontiers in Psychiatry</i> , 2017 , 8, 251	5	7
263	Retrospective propensity score matching and the selection of surgical procedures: how precise can a propensity estimate be?. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3200-1	2.2	7
262	Fear of negative evaluation is associated with delusional ideation in non-clinical population and patients with schizophrenia. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011 , 46, 703-10	4.5	7

261	Milnacipran versus other anti-depressive agents for depression 2007,		7
260	Japanese version of the Quality of Relationship Inventory: its reliability and validity among women with recurrent spontaneous abortion. <i>Psychiatry and Clinical Neurosciences</i> , 2002 , 56, 527-32	6.2	7
259	Parental loss in childhood and social support in adulthood among psychiatric patients. Group for Longitudinal Affective Disorders Study (GLADS). <i>Journal of Psychiatric Research</i> , 1999 , 33, 165-9	5.2	7
258	Longitudinal Analysis of Bidirectional Relationships between Nocturia and Depressive Symptoms: The Nagahama Study. <i>Journal of Urology</i> , 2020 , 203, 984-990	2.5	7
257	Impact of Nocturia on Mortality: The Nagahama Study. <i>Journal of Urology</i> , 2020 , 204, 996-1002	2.5	7
256	Patient reported outcome measures in clinical trials should be initially analyzed as continuous outcomes for statistical significance and responder analyses should be reserved as secondary analyses. <i>Journal of Clinical Epidemiology</i> , 2021 , 134, 95-102	5.7	7
255	Development and validation of a prediction model for the probability of responding to placebo in antidepressant trials: a pooled analysis of individual patient data. <i>Evidence-Based Mental Health</i> , 2019 , 22, 10-16	11.1	7
254	A review of methods for addressing components of interventions in meta-analysis. <i>PLoS ONE</i> , 2021 , 16, e0246631	3.7	7
253	Enduring effects of a 5-week behavioral activation program for subthreshold depression among late adolescents: an exploratory randomized controlled trial. <i>Neuropsychiatric Disease and Treatment</i> , 2018 , 14, 2633-2641	3.1	7
252	Initial treatment choices to achieve sustained response in major depression: a systematic review and network meta-analysis. <i>World Psychiatry</i> , 2021 , 20, 387-396	14.4	7
252 251		14.4	7
	and network meta-analysis. World Psychiatry, 2021, 20, 387-396 Reliability and validity of the Japanese version of the coping inventory for adults for stressful		
251	and network meta-analysis. World Psychiatry, 2021, 20, 387-396 Reliability and validity of the Japanese version of the coping inventory for adults for stressful situations in healthy people. Psychological Reports, 2015, 116, 447-69 Psychological interventions for positive symptoms in schizophrenia: protocol for a network		
251 250	and network meta-analysis. World Psychiatry, 2021, 20, 387-396 Reliability and validity of the Japanese version of the coping inventory for adults for stressful situations in healthy people. Psychological Reports, 2015, 116, 447-69 Psychological interventions for positive symptoms in schizophrenia: protocol for a network meta-analysis of randomised controlled trials. BMJ Open, 2018, 8, e019280 Disciplinary slapping is associated with bullying involvement regardless of warm parenting in early	1.6	6
251250249	and network meta-analysis. World Psychiatry, 2021, 20, 387-396 Reliability and validity of the Japanese version of the coping inventory for adults for stressful situations in healthy people. Psychological Reports, 2015, 116, 447-69 Psychological interventions for positive symptoms in schizophrenia: protocol for a network meta-analysis of randomised controlled trials. BMJ Open, 2018, 8, e019280 Disciplinary slapping is associated with bullying involvement regardless of warm parenting in early adolescence. Journal of Adolescence, 2018, 68, 207-216 In-situation safety behaviors among patients with panic disorder: descriptive and correlational	1.6 3 3.4	6 6
251250249248	Reliability and validity of the Japanese version of the coping inventory for adults for stressful situations in healthy people. <i>Psychological Reports</i> , 2015 , 116, 447-69 Psychological interventions for positive symptoms in schizophrenia: protocol for a network meta-analysis of randomised controlled trials. <i>BMJ Open</i> , 2018 , 8, e019280 Disciplinary slapping is associated with bullying involvement regardless of warm parenting in early adolescence. <i>Journal of Adolescence</i> , 2018 , 68, 207-216 In-situation safety behaviors among patients with panic disorder: descriptive and correlational study. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 332-9 Cognitive behavioural therapies versus other psychological therapies for depression. <i>The Cochrane</i>	1.6 3 3·4 6.2	6 6 6
251250249248247	Reliability and validity of the Japanese version of the coping inventory for adults for stressful situations in healthy people. <i>Psychological Reports</i> , 2015 , 116, 447-69 Psychological interventions for positive symptoms in schizophrenia: protocol for a network meta-analysis of randomised controlled trials. <i>BMJ Open</i> , 2018 , 8, e019280 Disciplinary slapping is associated with bullying involvement regardless of warm parenting in early adolescence. <i>Journal of Adolescence</i> , 2018 , 68, 207-216 In-situation safety behaviors among patients with panic disorder: descriptive and correlational study. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 332-9 Cognitive behavioural therapies versus other psychological therapies for depression. <i>The Cochrane Library</i> , 2010 ,	1.6 3 3·4 6.2	66666

243	Fluvoxamine versus other anti-depressive agents for depression 2006,		6	
242	Are the cochrane group registers comprehensive? A case study of Japanese psychiatry trials. <i>BMC Medical Research Methodology</i> , 2002 , 2, 6	4.7	6	
241	Parents, personality or partner? Correlates of marital relationships. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2002 , 37, 164-8	4.5	6	
240	Reliability, validity and standardization of the Japanese version of the Social Adjustment Scale-Self Report. <i>Psychiatry and Clinical Neurosciences</i> , 2003 , 57, 441-6	6.2	6	
239	A polydiagnostic study of depressive disorders according to DSM-IV and 23 classical diagnostic systems. <i>Psychiatry and Clinical Neurosciences</i> , 1999 , 53, 387-96	6.2	6	
238	No benefit from flexible titration above minimum licensed dose in prescribing antidepressants for major depression: systematic review. <i>Acta Psychiatrica Scandinavica</i> , 2020 , 141, 401-409	6.5	6	
237	Psychologic Treatment of Depression Compared With Pharmacotherapy and Combined Treatment in Primary Care: A Network Meta-Analysis. <i>Annals of Family Medicine</i> , 2021 , 19, 262-270	2.9	6	
236	Prediction of remission in pharmacotherapy of untreated major depression: development and validation of multivariable prediction models. <i>Psychological Medicine</i> , 2019 , 49, 2405-2413	6.9	6	
235	The Kilim plot: A tool for visualizing network meta-analysis results for multiple outcomes. <i>Research Synthesis Methods</i> , 2021 , 12, 86-95	7.2	6	
234	A Bayesian dose-response meta-analysis model: A simulations study and application. <i>Statistical Methods in Medical Research</i> , 2021 , 30, 1358-1372	2.3	6	
233	Applying GRADE to a network meta-analysis of antidepressants led to more conservative conclusions. <i>Journal of Clinical Epidemiology</i> , 2018 , 102, 87-98	5.7	6	
232	Comparing methods for estimating patient-specific treatment effects in individual patient data meta-analysis. <i>Statistics in Medicine</i> , 2021 , 40, 1553-1573	2.3	6	
231	Choto-san versus placebo for patients with dementia: systematic review and meta-analysis. <i>Psychogeriatrics</i> , 2017 , 17, 466-478	1.8	5	
230	Further evidence that item responses on the Kessler Psychological Distress Scale exhibit the characteristic pattern in the general population. <i>Heliyon</i> , 2019 , 5, e01387	3.6	5	
229	The impact of selective serotonin reuptake inhibitors on the thyroid function among patients with major depressive disorder: A systematic review and meta-analysis. <i>European Neuropsychopharmacology</i> , 2020 , 33, 139-145	1.2	5	
228	Identification of malignant early repolarization pattern by late QRS activity in high-resolution magnetocardiography. <i>Annals of Noninvasive Electrocardiology</i> , 2020 , 25, e12741	1.5	5	
227	Cumulative network meta-analyses, practice guidelines, and actual prescriptions for postmenopausal osteoporosis: a meta-epidemiological study. <i>Archives of Osteoporosis</i> , 2020 , 15, 21	2.9	5	
226	Measuring Clinical Importance in a Trial of Interventions for Mixed Urinary Incontinence. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 479	27.4	5	

225	Comparative efficacy and tolerability of new-generation antidepressants for major depressive disorder in children and adolescents: protocol of an individual patient data meta-analysis. <i>BMJ Open</i> , 2018 , 8, e018357	3	5
224	Development and validation of the Japanese version of the problem gambling severity index. <i>Addictive Behaviors</i> , 2019 , 98, 105987	4.2	5
223	Association between statistical significance and time to publication among systematic reviews: a study protocol for a meta-epidemiological investigation. <i>BMJ Open</i> , 2017 , 7, e018856	3	5
222	Effects of an internet-based cognitive behavioural therapy intervention on preventing major depressive episodes among workers: a protocol for a randomised controlled trial. <i>BMJ Open</i> , 2015 , 5, e007590	3	5
221	Brief psychoeducation for schizophrenia primarily intended to change the cognition of auditory hallucinations: an exploratory study. <i>Journal of Nervous and Mental Disease</i> , 2014 , 202, 35-9	1.8	5
220	Group cognitive behavioral therapy for patients with generalized social anxiety disorder in Japan: outcomes at 1-year follow up and outcome predictors. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 267-75	3.1	5
219	Not only body weight perception but also body mass index is relevant to suicidal ideation and self-harming behavior in Japanese adolescents. <i>Journal of Nervous and Mental Disease</i> , 2012 , 200, 305-9	1.8	5
218	Adequacy of continuation and maintenance treatments for major depression in Japan. <i>Journal of Psychopharmacology</i> , 2008 , 22, 153-6	4.6	5
217	Factor structure of social support and its relationship to minor psychiatric disorders among Japanese adolescents. <i>International Journal of Social Psychiatry</i> , 1995 , 41, 88-102	8.5	5
216	Characteristics of chronic outpatients with unipolar depression. <i>Acta Psychiatrica Scandinavica</i> , 1993 , 88, 273-7	6.5	5
215	Linking the Mini-Mental State Examination, the Alzheimer's Disease Assessment Scale-Cognitive Subscale and the Severe Impairment Battery: evidence from individual participant data from five randomised clinical trials of donepezil. <i>Evidence-Based Mental Health</i> , 2021 , 24, 56-61	11.1	5
214	Comparative efficacy and safety of bone-modifying agents for the treatment of bone metastases in patients with advanced renal cell carcinoma: a systematic review and meta-analysis. <i>Oncotarget</i> , 2017 , 8, 68890-68898	3.3	5
213	No consistent evidence of data availability bias existed in recent individual participant data meta-analyses: a meta-epidemiological study. <i>Journal of Clinical Epidemiology</i> , 2020 , 118, 107-114.e5	5.7	5
212	Vitamin C therapy for patients with sepsis or septic shock: a protocol for a systematic review and a network meta-analysis. <i>BMJ Open</i> , 2019 , 9, e033458	3	5
211	The association between role model presence and self-regulation in early adolescence: A cross-sectional study. <i>PLoS ONE</i> , 2019 , 14, e0222752	3.7	4
210	Predicting antidepressant response through early improvement of individual symptoms of depression incorporating baseline characteristics of patients: An individual patient data meta-analysis. <i>Journal of Psychiatric Research</i> , 2020 , 125, 85-90	5.2	4
209	The Characteristics and Social Functioning of Pathological Social Withdrawal, "Hikikomori," in a Secondary Care Setting: a One-Year Cohort Study. <i>BMC Psychiatry</i> , 2020 , 20, 352	4.2	4
208	Overstatements in abstract conclusions claiming effectiveness of interventions in psychiatry: a study protocol for a meta-epidemiological investigation. <i>BMJ Open</i> , 2016 , 6, e009832	3	4

207	Quality and impact of secondary information in promoting evidence-based clinical practice: a cross-sectional study about EBMH. <i>Evidence-Based Mental Health</i> , 2016 , 19, 82-5	11.1	4	
206	Brief multifamily Psychoeducation for family members of patients with chronic major depression: a randomized controlled trial. <i>BMC Psychiatry</i> , 2018 , 18, 207	4.2	4	
205	Comments on "Omentum-preserving gastrectomy for advanced gastric cancer: a propensity-matched retrospective cohort study". <i>Gastric Cancer</i> , 2014 , 17, 757-8	7.6	4	
204	Redundant systematic reviews on the same topic in surgery: a study protocol for a meta-epidemiological investigation. <i>BMJ Open</i> , 2017 , 7, e017411	3	4	
203	Enthusiasm for homework and improvement of psychological distress in subthreshold depression during behavior therapy: secondary analysis of data from a randomized controlled trial. <i>BMC Psychiatry</i> , 2015 , 15, 302	4.2	4	
202	Strategic use of new generation antidepressants for depression: SUN(^_^) D protocol update and statistical analysis plan. <i>Trials</i> , 2015 , 16, 459	2.8	4	
201	Effect of depression-related somatic pain on treatment satisfaction and daily living functions. <i>International Journal of Psychiatry in Medicine</i> , 2013 , 46, 27-38	1	4	
200	Interoceptive hypersensitivity as prognostic factor among patients with panic disorder who have received cognitive behavioral therapy. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2010 , 41, 325-9	2.6	4	
199	Making the best use of available evidence: the case of new generation antidepressants: a response to: are all antidepressants equal?. <i>Evidence-Based Mental Health</i> , 2009 , 12, 101-4	11.1	4	
198	Hallucinations in visually impaired individuals: an analysis of the National Comorbidity Survey Replication. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 104-8	4.5	4	
197	Neural correlates of memory in depression measured by brain perfusion SPECT at rest. <i>Psychiatry and Clinical Neurosciences</i> , 2009 , 63, 685-92	6.2	4	
196	Social anxiety disorder as a hidden psychiatric comorbidity among cancer patients. <i>Palliative and Supportive Care</i> , 2011 , 9, 103-5	2.5	4	
195	Reboxetine versus other antidepressive agents for depression. The Cochrane Library, 2009,	5.2	4	
194	Concordance between self- and observer-ratings on Kasahara's Inventory for the Melancholic Type Personality. <i>Psychiatry and Clinical Neurosciences</i> , 2002 , 56, 569-74	6.2	4	
193	Periodic psychosis associated with pseudo-pseudohypoparathyroidism. <i>Journal of Nervous and Mental Disease</i> , 1991 , 179, 637-8	1.8	4	
192	Family interventions for relapse prevention in schizophrenia: a systematic review and network meta-analysis <i>Lancet Psychiatry,the</i> , 2022 ,	23.3	4	
191	Acute interventions for aggression and agitation in psychosis: study protocol for a systematic review and network meta-analysis. <i>BMJ Open</i> , 2019 , 9, e032726	3	4	
190	Relationship between violent behavior and repeated weight-loss dieting among female adolescents in Japan. <i>PLoS ONE</i> , 2014 , 9, e107744	3.7	4	

189	The lack of statistical power of subgroup analyses in meta-analyses: a cautionary note. <i>Epidemiology and Psychiatric Sciences</i> , 2021 , 30, e78	5.1	4
188	The effects of psychological treatments of depression in children and adolescents on response, reliable change, and deterioration: a systematic review and meta-analysis. <i>European Child and Adolescent Psychiatry</i> , 2021 ,	5.5	4
187	Primary Endpoints in Current Phase II/III Trials for Alzheimer Disease: A Systematic Survey of Trials Registered at ClinicalTrials.gov. <i>Alzheimer Disease and Associated Disorders</i> , 2020 , 34, 97-100	2.5	4
186	Most at-risk individuals will not develop a mental disorder: the limited predictive strength of risk factors. <i>World Psychiatry</i> , 2021 , 20, 224-225	14.4	4
185	Polymyxin B-immobilised haemoperfusion and mortality in critically ill patients with sepsis/septic shock: a protocol for a systematic review and meta-analysis. <i>BMJ Open</i> , 2016 , 6, e012908	3	4
184	Guided or self-guided internet-based cognitive-behavioural therapy (iCBT) for depression? Study protocol of an individual participant data network meta-analysis. <i>BMJ Open</i> , 2019 , 9, e026820	3	4
183	Certolizumab pegol for induction of remission in Crohn's disease. <i>The Cochrane Library</i> , 2019 , 8, CD0128	3932	4
182	Permutation inference methods for multivariate meta-analysis. <i>Biometrics</i> , 2020 , 76, 337-347	1.8	4
181	How can we estimate QALYs based on PHQ-9 scores? Equipercentile linking analysis of PHQ-9 and EQ-5D. <i>Evidence-Based Mental Health</i> , 2021 ,	11.1	4
180	Efficient two-step multivariate random effects meta-analysis of individual participant data for longitudinal clinical trials using mixed effects models. <i>BMC Medical Research Methodology</i> , 2019 , 19, 33	4.7	3
179	Postcard intervention for depression in community-dwelling older adults: A randomised controlled trial. <i>Psychiatry Research</i> , 2015 , 229, 545-50	9.9	3
178	Can personalized treatment prediction improve the outcomes, compared with the group average approach, in a randomized trial? Developing and validating a multivariable prediction model in a pragmatic megatrial of acute treatment for major depression. <i>Journal of Affective Disorders</i> , 2020 ,	6.6	3
177	Cognitive-behavioural therapy for chronic subjective dizziness: Predictors of improvement in Dizziness Handicap Inventory at 6 months posttreatment. <i>Acta Oto-Laryngologica</i> , 2020 , 140, 827-832	1.6	3
176	Optimal sampling in derivation studies was associated with improved discrimination in external validation for heart failure prognostic models. <i>Journal of Clinical Epidemiology</i> , 2020 , 121, 71-80	5.7	3
175	Subcuticular sutures for skin closure in non-obstetric surgery. <i>The Cochrane Library</i> , 2020 , 4, CD012124	5.2	3
174	Statistical significance did not affect time to publication in non-Cochrane systematic reviews: a metaepidemiological study. <i>Journal of Clinical Epidemiology</i> , 2019 , 115, 25-34	5.7	3
173	Psychological therapies for panic disorder with or without agoraphobia in adults 2014 ,		3
172	Video-assisted thoracoscopic surgery lobectomy or open lobectomy for non-small-cell lung cancer? Minimizing selection bias in observational studies. <i>European Journal of Cardio-thoracic Surgery</i> , 2014 , 46, 150	3	3

171	New oral anticoagulants versus warfarin for preventing stroke and systemic embolic events among atrial fibrillation patients with chronic kidney disease. <i>The Cochrane Library</i> , 2014 ,	5.2	3
170	Near-infrared spectroscopy of bipolar disorder may be distinct from that of unipolar depression and of healthy controls. <i>Asia-Pacific Psychiatry</i> , 2012 , 4, 258-265	3.2	3
169	Can assessors in a psychotherapy trial be successfully blinded? Analysis of a randomized controlled trial on psychotherapy for refractory insomnia in residual depression. <i>Psychotherapy and Psychosomatics</i> , 2013 , 82, 401-3	9.4	3
168	Dissociations between reading responses and semantic priming effects in a dyslexic patient. <i>Cortex</i> , 1997 , 33, 753-61	3.8	3
167	Neuroleptic malignant syndrome-like state in an epileptic patient with organic brain comorbidity treated with zonisamide and carbamazepine. <i>Epilepsia</i> , 2007 , 48, 1999-2001	6.4	3
166	A case of frontotemporal lobar degeneration (FTLD) with panic attack as the first symptom. <i>Journal of Neuropsychiatry and Clinical Neurosciences</i> , 2007 , 19, 485-7	2.7	3
165	Meta-analyses and megatrials: neither is the infallible, universal standard. <i>Evidence-Based Mental Health</i> , 2004 , 7, 34-5	11.1	3
164	Review: placebo is better than no treatment for subjective continuous outcomes and for treatment of pain. <i>Evidence-Based Mental Health</i> , 2002 , 5, 15	11.1	3
163	Selective publication of antidepressant trials and its influence on apparent efficacy: Updated comparisons and meta-analyses of newer versus older trials <i>PLoS Medicine</i> , 2022 , 19, e1003886	11.6	3
162	Quality of COVID-19 research in preprints: a meta-epidemiological study protocol v1		3
161	Influence diagnostics and outlier detection for meta-analysis of diagnostic test accuracy. <i>Research Synthesis Methods</i> , 2020 , 11, 237-247	7.2	3
160	Optimal Dose of Selective Serotonin Reuptake Inhibitors, Venlafaxine, and Mirtazapine in Major Depression: A Systematic Review and Dose-Response Meta-Analysis. <i>Focus (American Psychiatric Publishing)</i> , 2020 , 18, 211-219	1.1	3
159	Responses to depressive symptom items exhibit a common mathematical pattern across the European populations. <i>Scientific Reports</i> , 2019 , 9, 14923	4.9	3
158	Factor structure of the Japanese version of the Quality of Life in Alzheimer's Disease Scale (QOL-AD). <i>Psychogeriatrics</i> , 2020 , 20, 79-86	1.8	3
157	Whose depression deteriorates during acute phase antidepressant treatment?. <i>Journal of Affective Disorders</i> , 2020 , 260, 342-348	6.6	3
156	Clinical feasibility and acceptability of adding cognitive behavioral therapy to pharmacotherapy for drug-resistant overactive bladder in women: A single-arm pilot study. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2021 , 13, 69-78	1.9	3
155	Melancholic features (DSM-IV) predict but do not moderate response to antidepressants in major depression: an individual participant data meta-analysis of 1219 patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 271, 521-526	5.1	3
154	Network meta-analysis of antidepressants - Authors' reply. <i>Lancet, The</i> , 2018 , 392, 1012-1013	40	3

153	Predictors of Broad Dimensions of Psychopathology among Patients with Panic Disorder after Cognitive-Behavioral Therapy. <i>Psychiatry Journal</i> , 2018 , 2018, 5183834	2.4	3
152	The characteristics of patients with severe social withdrawal 'Hikikomori' in two community psychiatry clinics in Japan. <i>Asian Journal of Psychiatry</i> , 2021 , 65, 102833	6.7	3
151	Publication statuses of clinical trials supporting FDA-approved immune checkpoint inhibitors: a meta-epidemiological investigation. <i>BMC Cancer</i> , 2019 , 19, 998	4.8	2
150	Smartphone problem-solving therapy to reduce fear of cancer recurrence among breast cancer survivors: an open single-arm pilot study. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 537-544	2.8	2
149	Depressed with cancer can respond to antidepressants, but further research is needed to confirm and expand on these findings. <i>Evidence-Based Mental Health</i> , 2015 , 18, 28	11.1	2
148	Group cognitive behavioural therapy (GCBT) versus treatment as usual (TAU) in the treatment of irritable bowel syndrome (IBS): a study protocol for a randomized controlled trial. <i>BMC Gastroenterology</i> , 2020 , 20, 29	3	2
147	Adolescent depression: from symptoms to individualised treatment?. Lancet Psychiatry, the, 2020, 7, 29	5 -2396	2
146	Acceptability of escitalopram versus duloxetine in outpatients with depression who did not respond to initial second-generation antidepressants: Study protocol for a randomized, parallel-group, non-inferiority trial. <i>Neuropsychopharmacology Reports</i> , 2019 , 39, 262-272	2.2	2
145	The postcard intervention against depression among community-dwelling older adults: study protocol for a randomized controlled trial. <i>Trials</i> , 2013 , 14, 202	2.8	2
144	Antidepressants, benzodiazepines and azapirones for panic disorder in adults: a network meta-analysis. <i>The Cochrane Library</i> , 2017 ,	5.2	2
143	Hyperbaric oxygen therapy for persistent post-concussion syndrome following mild traumatic brain injury. <i>The Cochrane Library</i> , 2017 ,	5.2	2
142	The relationship between depression and violent crime. Lancet Psychiatry, the, 2015, 2, 193-4	23.3	2
141	Public speaking fears and their correlates among 17,615 Japanese adolescents. <i>Asia-Pacific Psychiatry</i> , 2014 , 6, 99-104	3.2	2
140	Case of intrathecal baclofen-induced psychotic symptoms. <i>Psychiatry and Clinical Neurosciences</i> , 2011 , 65, 300-1	6.2	2
139	Reliability and validity of the Japanese version of the Agitated Behaviour in Dementia Scale in Alzheimer's disease: three dimensions of agitated behaviour in dementia. <i>Psychogeriatrics</i> , 2011 , 11, 212-20	1.8	2
138	Escitalopram versus other anti-depressive agents for depression 2007,		2
137	Cognitive impairment networks in Alzheimer's disease: Analysis of three double-blind randomized, placebo-controlled, clinical trials of donepezil <i>European Neuropsychopharmacology</i> , 2022 , 57, 50-58	1.2	2
136	Drug treatment for generalised anxiety disorder. <i>BMJ, The</i> , 2011 , 342, d1216	5.9	2

135	ROB-MEN: a tool to assess risk of bias due to missing evidence in network meta-analysis. <i>BMC Medicine</i> , 2021 , 19, 304	11.4	2
134	Differences in the Effect of Internet-Based Cognitive Behavioral Therapy for Improving Nonclinical Depressive Symptoms Among Workers by Time Preference: Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2018 , 20, e10231	7.6	2
133	Which psychotherapy is effective in panic disorder? And which delivery formats are supported by the evidence? Study protocol for two systematic reviews and network meta-analyses. <i>BMJ Open</i> , 2020 , 10, e038909	3	2
132	Use of social networking sites and desire for slimness among 10-year-old girls and boys: A population-based birth cohort study. <i>International Journal of Eating Disorders</i> , 2020 , 53, 288-295	6.3	2
131	Age-Related Changes in Item Responses to the Patient Health Questionnaire-9: Evidence From the National Health and Nutrition Examination Survey. <i>Frontiers in Psychiatry</i> , 2020 , 11, 723	5	2
130	An efficient way to assess the effect of COVID-19 on mental health in the general population. <i>Lancet Psychiatry,the</i> , 2021 , 8, e14-e15	23.3	2
129	Confidence intervals of prediction accuracy measures for multivariable prediction models based on the bootstrap-based optimism correction methods. <i>Statistics in Medicine</i> , 2021 , 40, 5691-5701	2.3	2
128	Subcuticular sutures for skin closure in non-obstetric surgery. The Cochrane Library, 2016,	5.2	2
127	Psychosocial treatments for relapse prevention in schizophrenia: study protocol for a systematic review and network meta-analysis of randomised evidence. <i>BMJ Open</i> , 2020 , 10, e035073	3	2
126	Visualizing the evolution of evidence: Cumulative network meta-analyses of new generation antidepressants in the last 40 years. <i>Research Synthesis Methods</i> , 2021 , 12, 74-85	7.2	2
125	Prognosis of delirium. Evidence-Based Mental Health, 2017, 20, 53-54	11.1	1
124	DSM-III-R change in definition might have affected placebo response to antidepressants - Authors' reply. <i>Lancet Psychiatry,the</i> , 2017 , 4, 22-23	23.3	1
123	Optimal dosing of antidepressant drugs - Authors' reply. Lancet Psychiatry, the, 2019, 6, 806-807	23.3	1
122	Physician characteristics associated with proper assessment of overstated conclusions in research abstracts: A secondary analysis of a randomized controlled trial. <i>PLoS ONE</i> , 2019 , 14, e0211206	3.7	1
121	Which drug is the best bet when your bipolar patient gets depressed?. <i>Evidence-Based Mental Health</i> , 2015 , 18, 80-1	11.1	1
120	Predicting the treatment response of certolizumab for individual adult patients with rheumatoid arthritis: protocol for an individual participant data meta-analysis. <i>Systematic Reviews</i> , 2020 , 9, 140	3	1
119	Note Added to Indicate Related Article Was Retracted. <i>JAMA Psychiatry</i> , 2020 , 77, 878	14.5	1
118	Evidence-Based Utilization of Prognostic Prediction Models in Cardiovascular Medicine. <i>Circulation Reports</i> , 2019 , 2, 10-16	0.7	1

117	Concerns About Selective Outcome Reporting. Journal of Clinical Oncology, 2017, 35, 688	2.2	1
116	Daikenchuto for reducing postoperative ileus in patients undergoing elective abdominal surgery. <i>The Cochrane Library</i> , 2018 , 4, CD012271	5.2	1
115	Efficacy of antidepressants over placebo is similar in two-armed versus three-armed or more-armed randomized placebo-controlled trials. <i>International Clinical Psychopharmacology</i> , 2018 , 33, 66-72	2.2	1
114	Dropouts in an Antidepressant Trial: How Do They Fare Afterwards?. <i>Psychotherapy and Psychosomatics</i> , 2018 , 87, 380-382	9.4	1
113	Why some depressive patients perform suicidal acts and others do not. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 660-661	6.2	1
112	Psychological therapies versus pharmacological interventions for panic disorder with or without agoraphobia in adults 2014 ,		1
111	Association between eicosapentaenoic acid (EPA) medication intake and new onset of depression among Japanese patients with hyperlipidemia: A 3-year follow-up study. <i>Journal of Nutrition & Intermediary Metabolism</i> , 2017 , 9, 12-16	2.8	1
110	Study protocol for a randomised controlled trial of cognitive processing therapy for post-traumatic stress disorder among Japanese patients: the Safety, Power, Intimacy, Esteem, Trust (SPINET) study. <i>BMJ Open</i> , 2017 , 7, e014292	3	1
109	The Mechanisms Underlying Changes in Broad Dimensions of Psychopathology During Cognitive Behavioral Therapy for Social Anxiety Disorder. <i>Journal of Clinical Medicine Research</i> , 2017 , 9, 1019-102	1 ^{2.9}	1
108	Ramelteon for insomnia. <i>The Cochrane Library</i> , 2014 ,	5.2	1
108	Ramelteon for insomnia. <i>The Cochrane Library</i> , 2014 , Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 ,	5.2 5.2	1
			1
107	Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 ,	5.2	1
107	Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 , Antidepressants during ECT. <i>Evidence-Based Mental Health</i> , 2014 , 17, 25-6 Behavior management approach for agitated behavior in Japanese patients with dementia: a pilot	5.2	1
107 106 105	Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 , Antidepressants during ECT. <i>Evidence-Based Mental Health</i> , 2014 , 17, 25-6 Behavior management approach for agitated behavior in Japanese patients with dementia: a pilot study. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 9-14 Preoperative inspiratory muscle training for postoperative pulmonary complications in adult	5.2	1 1
107 106 105	Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 , Antidepressants during ECT. <i>Evidence-Based Mental Health</i> , 2014 , 17, 25-6 Behavior management approach for agitated behavior in Japanese patients with dementia: a pilot study. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 9-14 Preoperative inspiratory muscle training for postoperative pulmonary complications in adult patients undergoing cardiac and major abdominal surgery 2013 , Panic disorder. When symptoms have no apparent physical cause. <i>Postgraduate Medicine</i> , 1997 ,	5.2 11.1 3.1	1 1 1
107 106 105 104	Bupropion versus other antidepressive agents for depression. <i>The Cochrane Library</i> , 2014 , Antidepressants during ECT. <i>Evidence-Based Mental Health</i> , 2014 , 17, 25-6 Behavior management approach for agitated behavior in Japanese patients with dementia: a pilot study. <i>Neuropsychiatric Disease and Treatment</i> , 2013 , 9, 9-14 Preoperative inspiratory muscle training for postoperative pulmonary complications in adult patients undergoing cardiac and major abdominal surgery 2013 , Panic disorder. When symptoms have no apparent physical cause. <i>Postgraduate Medicine</i> , 1997 , 101, 147-50, 153-4, 159-60 passim	5.2 11.1 3.1 3.7 4.9	1 1 1 1 1

99	A systematic survey identified methodological issues in studies estimating anchor-based minimal important differences in patient-reported outcomes. <i>Journal of Clinical Epidemiology</i> , 2021 , 142, 144-1	5∮· ⁷	1
98	Substance use disorder treatment success: assessing patient-reported use improvement and provider-evaluated treatment completion. <i>Addiction Research and Theory</i> ,1-8	2.6	1
97	A possible association between paternal longevity and major depression among elderly men. <i>Psychiatry and Clinical Neurosciences</i> , 1998 , 52, 577-9	6.2	1
96	Behavioral Activation Contributed to the Total Reduction of Depression Symptoms in the Smartphone-based Cognitive Behavioral Therapy: A Secondary Analysis of a Randomized, Controlled Trial. <i>Innovations in Clinical Neuroscience</i> , 2020 , 17, 21-25	1	1
95	Optimal dose of aripiprazole for augmentation therapy of antidepressant-refractory depression: preliminary findings based on a systematic review and dose-effect meta-analysis <i>British Journal of Psychiatry</i> , 2021 , 1-8	5.4	1
94	Comparative efficacy and acceptability of psychotherapies for panic disorder with or without agoraphobia: systematic review and network meta-analysis of randomised controlled trials <i>British Journal of Psychiatry</i> , 2021 , 1-13	5.4	1
93	State-of-the-Art Reporting 2016 , 189-202		1
92	Efficacy of brief intervention for drug misuse in primary care facilities: systematic review and meta-analysis protocol. <i>BMJ Open</i> , 2020 , 10, e036633	3	1
91	Cognitive behavioral therapy for overactive bladder in women: study protocol for a randomized controlled trial. <i>BMC Urology</i> , 2020 , 20, 129	2.2	1
90	The risk perception against COVID-19 and outpatients' anxiety of visiting the clinic during COVID-19 pandemic. <i>Psychiatry and Clinical Neurosciences</i> , 2021 , 75, 208-209	6.2	1
89	Antidepressant prescriptions have not fully reflected evolving evidence from cumulative network meta-analyses and guideline recommendations. <i>Journal of Clinical Epidemiology</i> , 2021 , 133, 14-23	5.7	1
88	Anxiety Impacts Consent Capacity to Treatment in Alzheimer's Disease. <i>Frontiers in Psychology</i> , 2021 , 12, 685430	3.4	1
87	Assessing recurrence of depression using a zero-inflated negative binomial model: A secondary analysis of lifelog data. <i>Psychiatry Research</i> , 2021 , 300, 113919	9.9	1
86	Psychological Impact of Masks and Infection-Prevention Plastic Partitions on Anxiety for Infection and Patient-Doctor Relationships During the COVID-19 Pandemic. <i>primary care companion for CNS disorders, The</i> , 2021 , 23,	1.2	1
85	Electronic health record nested pragmatic randomized controlled trial of a reminder system for serum lithium level monitoring in patients with mood disorder: KONOTORI study protocol. <i>Trials</i> , 2019 , 20, 706	2.8	1
84	Do newspapers preferentially cover biomedical studies involving national scientists?. <i>Public Understanding of Science</i> , 2019 , 28, 191-200	3.1	1
83	Response inhibition and anxiety in adolescents: Results from a population-based community sample. <i>Journal of Affective Disorders</i> , 2019 , 246, 89-95	6.6	1
82	Predictors of hypomanic and/or manic switch among patients initially diagnosed with unipolar major depression during acute-phase antidepressants treatment. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 90-91	6.2	1

81	Frequentist performances of Bayesian prediction intervals for random-effects meta-analysis. <i>Biometrical Journal</i> , 2021 , 63, 394-405	1.5	1
80	Quantifying the heterogeneity of cognitive functioning in Alzheimer's disease to extend the placebo-treatment dichotomy: Latent class analysis of individual-participant data from five pivotal randomized clinical trials of donepezil. <i>European Psychiatry</i> , 2021 , 64, e16	6	1
79	An individualized treatment rule to optimize probability of remission by continuation, switching, or combining antidepressant medications after failing a first-line antidepressant in a two-stage randomized trial. <i>Psychological Medicine</i> , 2021 , 1-10	6.9	1
78	Systematic differences in effect estimates between observational studies and randomized control trials in meta-analyses in nephrology. <i>Scientific Reports</i> , 2021 , 11, 6088	4.9	1
77	Cumulative network-meta-analyses, practice guidelines and actual prescriptions of drug treatments for postmenopausal osteoporosis: a study protocol for cumulative network meta-analyses and meta-epidemiological study. <i>BMJ Open</i> , 2018 , 8, e023218	3	1
76	The GAD-7 and the PHQ-8 exhibit the same mathematical pattern of item responses in the general population: analysis of data from the National Health Interview Survey. <i>BMC Psychology</i> , 2021 , 9, 149	2.8	1
75	Factors supporting substance use improvement for Black Americans: A population health observational study <i>Drug and Alcohol Dependence</i> , 2022 , 234, 109405	4.9	1
74	Trajectory of criterion symptoms of major depression under newly started antidepressant treatment: sleep disturbances and anergia linger on while suicidal ideas and psychomotor symptoms disappear early. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 140, 532-540	6.5	O
73	The relationships between symptoms and quality of life over the course of cognitive-behavioral therapy for panic disorder in Japan. <i>Asia-Pacific Psychiatry</i> , 2017 , 9, e12249	3.2	O
72	Early- and subsequent- response of cognitive functioning in Alzheimer's disease: Individual-participant data from five pivotal randomized clinical trials of donepezil <i>Journal of</i> Psychiatric Research, 2022 , 148, 159-164	5.2	O
71	About the issue of including or excluding studies from China in systematic reviews <i>Schizophrenia Research</i> , 2022 , 240, 162-163	3.6	0
70	A comprehensive validation of very early rule-out strategies for non-ST-segment elevation myocardial infarction in emergency departments: protocol for a multicentre prospective cohort study. <i>BMJ Open</i> , 2019 , 9, e026985	3	O
69	Acceptability of escitalopram versus duloxetine in outpatients with depression who did not respond to initial second-generation antidepressants: A randomized, parallel-group, non-inferiority trial. <i>Journal of Affective Disorders</i> , 2021 , 282, 1011-1020	6.6	0
68	Mathematical pattern of Kessler psychological distress distribution in the general population of the U.S. and Japan. <i>BMC Psychiatry</i> , 2021 , 21, 188	4.2	O
67	Protocol for development and validation of a prediction model for 5-year risk of incident overactive bladder in the general population: the Nagahama study. <i>BMC Urology</i> , 2021 , 21, 78	2.2	0
66	Impact of COVID-2019 on school attendance problems. <i>Journal of Global Health</i> , 2021 , 11, 03084	4.3	O
65	Network meta-analysis results against a fictional treatment of average performance: Treatment effects and ranking metric. <i>Research Synthesis Methods</i> , 2021 , 12, 161-175	7.2	0
64	High variability in results and methodological quality among overlapping systematic reviews on the same topics in surgery: a meta-epidemiological study. <i>British Journal of Surgery</i> , 2021 , 108, 1521-1529	5.3	O

63	Caregiver self-efficacy and associated factors among caregivers of patients with dementia with Lewy bodies and caregivers of patients with Alzheimer's disease. <i>Psychogeriatrics</i> , 2021 , 21, 783-794	1.8	0
62	Fingertip advanced glycation end products and psychotic symptoms among adolescents. <i>NPJ Schizophrenia</i> , 2021 , 7, 37	5.5	Ο
61	Around ten percent of most recent Cochrane reviews included outcomes in their literature search strategy and were associated with potentially exaggerated results: A research-on-research study. <i>Journal of Clinical Epidemiology</i> , 2021 , 141, 74-81	5.7	0
60	Roux-en-Y versus Billroth-I reconstruction after distal gastrectomy for gastric cancer. <i>The Cochrane Library</i> , 2021 , 9, CD012998	5.2	O
59	Development and validation of the Cognitive Behavioural Therapy Skills Scale among college students. <i>Evidence-Based Mental Health</i> , 2021 , 24, 70-76	11.1	O
58	A dose-effect network meta-analysis model with application in antidepressants using restricted cubic splines <i>Statistical Methods in Medical Research</i> , 2022 , 9622802211070256	2.3	Ο
57	Trends in Prescribing Antipsychotics for Children and Adolescents in Japan: A Descriptive Epidemiological Study Using a Large-Scale Pharmacy Dataset <i>Child Psychiatry and Human Development</i> , 2022 , 1	3.3	0
56	Unguided Computer-Assisted Self-Help Interventions Without Human Contact in Patients With Obsessive-Compulsive Disorder: Systematic Review and Meta-analysis <i>Journal of Medical Internet Research</i> , 2022 , 24, e35940	7.6	Ο
55	Association between patients' feedback comments and depressive mood, satisfaction, homework conducted, and dropouts during self-guided smartphone cognitive behavioral therapy. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 349-350	6.2	
54	Helping people with schizophrenia to quit smoking. Evidence-Based Mental Health, 2015, 18, 14-5	11.1	
53	Qualitative treatment-subgroup interactions in the antidepressant treatment of major depression: Application of QUINT to individual participant data from seven placebo-controlled randomized controlled trials. <i>Personalized Medicine in Psychiatry</i> , 2020 , 21-22, 100054	1.1	
52	Daikenchuto for reducing postoperative ileus in patients undergoing elective abdominal surgery. <i>The Cochrane Library</i> , 2020 , 3, CD012271	5.2	
51	Choosing the optimum strategy for rTMS. Evidence-Based Mental Health, 2018, 21, 77-78	11.1	
50	Response to letter regarding article by Formiga et al., "Utility of the Controlling Nutritional Status (CONUT) score in patients admitted due to acute heart failure". <i>International Journal of Cardiology</i> , 2018 , 256, 25	3.2	
49	Which psychotherapy for PTSD?. Evidence-Based Mental Health, 2016, 19, 118-119	11.1	
48	A guideline for whom?. <i>Epidemiology and Psychiatric Sciences</i> , 2016 , 25, 439-440	5.1	
47	Baseline severity and efficacy of antidepressants: into the third generation of research. <i>Lancet Psychiatry,the</i> , 2019 , 6, 715-716	23.3	
46	Bipolar treatment efficacy - Authors' reply. <i>Lancet Psychiatry,the</i> , 2014 , 1, 418-9	23.3	

45 Anxiety Disorders in Non-Western Countries **2014**, 612-621

44	Can we inflate effect size and thus increase chances of producing "positive" results if we raise the baseline threshold in schizophrenia trials?. <i>Schizophrenia Research</i> , 2013 , 144, 105-8	3.6
43	No significant difference in long-term remission between psychological therapies and antidepressants in panic disorder. <i>Evidence-Based Mental Health</i> , 2017 , 20, e18	11.1
42	What course are we to expect for a child with her first manic episode?. <i>Evidence-Based Mental Health</i> , 2015 , 18, 111-2	11.1
41	Reply to Letter: "Hand-sewn Versus Mechanical Esophagogastric Anastomosis After Esophagectomy: A Systematic Review and Meta-analysis". <i>Annals of Surgery</i> , 2015 , 261, e157	7.8
40	Reply to Letter: "Is There No Significant Difference in the Incidence of Anastomotic Leakage Between the Hand-sewn and Mechanical Esophagogastric Anastomosis?". <i>Annals of Surgery</i> , 2015 , 262, e81-2	7.8
39	Adverse effects of antidepressants during pregnancy. Evidence-Based Mental Health, 2014, 17, 103-4	11.1
38	Diagnostic test for dementias. <i>Evidence-Based Mental Health</i> , 2014 , 17, 45-6	11.1
37	Prognosis of a brief psychotic episode. Evidence-Based Mental Health, 2014, 17, 81-2	11.1
36	Mathematical model for the distribution of major depressive episode durations. <i>BMC Research Notes</i> , 2014 , 7, 636	2.3
35	A case with Hodgkin lymphoma and fronto-temporal lobular degeneration (FTLD)-like dementia facilitated by chemotherapy. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 365-8	2.8
34	A stepped-care prevention programme decreased depression and anxiety disorders in elderly patients. <i>Evidence-Based Medicine</i> , 2009 , 14, 146	
33	Exploring the link between depression and rheumatoid arthritis: prospects for optimal therapeutic success. <i>International Journal of Clinical Rheumatology</i> , 2010 , 5, 273-275	1.5
32	Psychosocial factors in recurrent miscarriages. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 1998 , 77, 572-573	3.8
31	Cognitive Patterns of Good and Poor Namers in Old Age. <i>Psychogeriatrics</i> , 2002 , 2, 35-39	1.8
30	The Influence of the Stroop Interference Effect on an Event-based Prospective Memory Task in Alzheimer's Disease Patients. <i>Psychogeriatrics</i> , 2002 , 2, 120-126	1.8
29	Cognitive therapy and social functioning in chronic depression. <i>British Journal of Psychiatry</i> , 2001 , 178, 477-8	5.4
28	Prognosis of depression and generalised anxiety in primary care. <i>British Journal of Psychiatry</i> , 2001 , 179, 177-8	5.4

27	Persistence of chronic mood disorders: a 2 year follow-up. <i>Psychiatry and Clinical Neurosciences</i> , 1995 , 49, 19-24	6.2
26	Reply by Authors. <i>Journal of Urology</i> , 2020 , 204, 1002	2.5
25	The inclusion of outcomes in search strategies for Cochrane Reviews: authors' reply. <i>Journal of Clinical Epidemiology</i> , 2021 ,	5-7
24	Clinical Relevance of the Behavioral Stages of Change with the Efficacy of Multifaceted Interventions on the Drop-Out from Regular Medical Care in Patients with Diabetes The Japan Diabetes Outcome Intervention Trial-2 (J-DOIT2). Diabetes, 2018, 67, 808-P	0.9
23	Publication statuses of clinical trials supporting immune checkpoint inhibitors recently approved by the United States Food and Drug Administration: A meta-epidemiological investigation <i>Journal of Clinical Oncology</i> , 2019 , 37, 89-89	2.2
22	Background styles in systematic review articles are not related to the publication in high-impact-factor journals: A meta-epidemiological study. <i>Medicine (United States)</i> , 2020 , 99, e23801	1.8
21	Comparative efficacy and safety of bone-modifying agents for the treatment of bone metastases in patients with advanced renal cell carcinoma: A systematic review and meta-analysis <i>Journal of Clinical Oncology</i> , 2017 , 35, e16067-e16067	2.2
20	The Relationship between Symptoms and Social Functioning over the Course of Cognitive Behavioral Therapy for Social Anxiety Disorder. <i>Psychiatry Journal</i> , 2020 , 2020, 3186450	2.4
19	Challenges in Systematic Reviews That Include Observational Studies: Comment on the Article by Karmacharya et al. <i>Arthritis and Rheumatology</i> , 2020 , 72, 2163-2164	9.5
18	Interpreting the Findings of a Meta-analysis of Psychosocial Interventions in Bipolar Disorder. <i>JAMA Psychiatry</i> , 2021 , 78, 448-449	14.5
17	Response to Jordan, Nich, Babuscio, Quainoo, and Carroll. <i>Drug and Alcohol Dependence</i> , 2021 , 223, 108	36 7.6
16	Diagnosing autism spectrum disorder (ASD) among adults. Evidence-Based Mental Health, 2016, 19, 46-	8 11.1
15	Treatment of depression in children and adolescents - Authors' reply. Lancet Psychiatry, the, 2021, 8, 97	-9283.3
14	Considerations Regarding a Network Meta-analysis of Systemic Treatments for Metastatic Castration-Sensitive Prostate Cancer. <i>JAMA Oncology</i> , 2021 , 7, 1068-1069	13.4
13	Increasing the clinical interpretability of PHQ-9 through equipercentile linking with health utility values by EQ-5D-3L. <i>Evidence-Based Mental Health</i> , 2021 , 24, e6	11.1
12	The Methodological Quality Score of COVID-19 Systematic Reviews is Low, Except for Cochrane Reviews: A Meta-epidemiological Study. <i>Annals of Clinical Epidemiology</i> , 2021 , 3, 46-55	2.5
11	State population indices predict Asian American and Pacific Islander successful substance use treatment completion: An exploratory observational study. <i>Journal of Ethnicity in Substance Abuse</i> , 2021 , 1-16	1.1
10	Fear of Fear and Broad Dimensions of Psychopathology over the Course of Cognitive Behavioural Therapy for Panic Disorder with Agoraphobia in Japan. <i>East Asian Archives of Psychiatry</i> , 2017 , 27, 150-5	5 1.1

9	Efficacy of Brief Intervention for Unhealthy Drug Use in Outpatient Medical Care: a Systematic Review and Meta-analysis <i>Journal of General Internal Medicine</i> , 2022 ,	4
8	Efficacy and acceptability of next step treatment strategies in adults with treatment-resistant major depressive disorder: protocol for systematic review and network meta-analysis <i>BMJ Open</i> , 2022 , 12, e056777	3
7	The association between role model presence and self-regulation in early adolescence: A cross-sectional study 2019 , 14, e0222752	
6	The association between role model presence and self-regulation in early adolescence: A cross-sectional study 2019 , 14, e0222752	
5	The association between role model presence and self-regulation in early adolescence: A cross-sectional study 2019 , 14, e0222752	
4	The association between role model presence and self-regulation in early adolescence: A cross-sectional study 2019 , 14, e0222752	
3	Rapid discontinuation of antidepressants reduces time to recurrence of depression and panic. <i>Evidence-Based Mental Health</i> , 2011 , 14, 13	11.1
2	What Is the Minimum Clinically Important Change in Negative Symptoms of Schizophrenia? PANSS Based Analyses of a Phase III Clinical Trial <i>Frontiers in Psychiatry</i> , 2022 , 13, 816339	5
1	Digital aids for relapse prevention in schizophrenia <i>Lancet Psychiatry,the</i> , 2022 , 9, 424-425	23.3