

# Vineesh Indira Chandran

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8215513/publications.pdf>

Version: 2024-02-01

11  
papers

307  
citations

933447

10  
h-index

1281871

11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

635  
citing authors

#	ARTICLE	IF	CITATIONS
1	Circulating TREM2 as a noninvasive diagnostic biomarker for NASH in patients with elevated liver stiffness. <i>Hepatology</i> , 2023, 77, 558-572.	7.3	17
2	Interplay between Caveolin-1 and body and tumor size affects clinical outcomes in breast cancer. <i>Translational Oncology</i> , 2022, 22, 101464.	3.7	3
3	Hypoxia Attenuates Trastuzumab Uptake and Trastuzumab-Emtansine (T-DM1) Cytotoxicity through Redistribution of Phosphorylated Caveolin-1. <i>Molecular Cancer Research</i> , 2020, 18, 644-656.	3.4	17
4	Global extracellular vesicle proteomic signature defines U87-MG glioma cell hypoxic status with potential implications for non-invasive diagnostics. <i>Journal of Neuro-Oncology</i> , 2019, 144, 477-488.	2.9	24
5	Ultrasensitive Immunoprofiling of Plasma Extracellular Vesicles Identifies Syndecan-1 as a Potential Tool for Minimally Invasive Diagnosis of Glioma. <i>Clinical Cancer Research</i> , 2019, 25, 3115-3127.	7.0	72
6	Extracellular lipid loading augments hypoxic paracrine signaling and promotes glioma angiogenesis and macrophage infiltration. <i>Journal of Experimental and Clinical Cancer Research</i> , 2019, 38, 241.	8.6	21
7	Tumor antigen glycosaminoglycan modification regulates antibody-drug conjugate delivery and cytotoxicity. <i>Oncotarget</i> , 2017, 8, 66960-66974.	1.8	17
8	Metastasis Stimulation by Hypoxia and Acidosis-Induced Extracellular Lipid Uptake Is Mediated by Proteoglycan-Dependent Endocytosis. <i>Cancer Research</i> , 2016, 76, 4828-4840.	0.9	79
9	Improved Pharmacokinetic and Biodistribution Properties of the Selective Urokinase Inhibitor PAI-2 (SerpinB2) by Site-Specific PEGylation: Implications for Drug Delivery. <i>Pharmaceutical Research</i> , 2015, 32, 1045-1054.	3.5	10
10	HER2 and uPAR cooperativity contribute to metastatic phenotype of HER2-positive breast cancer. <i>Oncoscience</i> , 2015, 2, 207-224.	2.2	21
11	Anti-cancer activity of an acid-labile N-alkylisatin conjugate targeting the transferrin receptor. <i>Cancer Letters</i> , 2012, 316, 151-156.	7.2	26