

Yishuo Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8223240/publications.pdf>

Version: 2024-02-01

30
papers

813
citations

687363

13
h-index

501196

28
g-index

31
all docs

31
docs citations

31
times ranked

1452
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic polymorphisms at 19q13.33 are associated with [math>\alpha^2]proPSA (p2PSA) levels and provide additional predictive value to prostate health index for prostate cancer. <i>Prostate</i> , 2021, 81, 971-982.	2.3	4
2	The Clinical Implications and Molecular Mechanism of CX3CL1 Expression in Urothelial Bladder Cancer. <i>Frontiers in Oncology</i> , 2021, 11, 752860.	2.8	6
3	A Germline Variant at 8q24 Contributes to the Serum p2PSA Level in a Chinese Prostate Biopsy Cohort. <i>Frontiers in Oncology</i> , 2021, 11, 753920.	2.8	2
4	HOXA9, PCDH17, POU4F2, and ONECUT2 as a Urinary Biomarker Combination for the Detection of Bladder Cancer in Chinese Patients with Hematuria. <i>European Urology Focus</i> , 2020, 6, 284-291.	3.1	27
5	The study on copy number alteration of clear cell renal cancer in Chinese population. <i>Journal of Cancer</i> , 2020, 11, 16-24.	2.5	2
6	Cost-Effectiveness Analysis of Prostate Health Index in Decision Making for Initial Prostate Biopsy. <i>Frontiers in Oncology</i> , 2020, 10, 565382.	2.8	10
7	Rare Germline Pathogenic Mutations of DNA Repair Genes Are Most Strongly Associated with Grade Group 5 Prostate Cancer. <i>European Urology Oncology</i> , 2020, 3, 224-230.	5.4	41
8	Family history is significantly associated with prostate cancer and its early onset in Chinese population. <i>Prostate</i> , 2019, 79, 1762-1766.	2.3	6
9	Concept and benchmarks for assessing narrow-sense validity of genetic risk score values. <i>Prostate</i> , 2019, 79, 1099-1105.	2.3	18
10	Systematic evaluation of cancer-specific genetic risk score for 11 types of cancer in The Cancer Genome Atlas and Electronic Medical Records and Genomics cohorts. <i>Cancer Medicine</i> , 2019, 8, 3196-3205.	2.8	22
11	Identification of Cancer-Specific Methylation of Gene Combination for the Diagnosis of Bladder Cancer. <i>Journal of Cancer</i> , 2019, 10, 6761-6766.	2.5	37
12	Germline Mutations in ATM and BRCA1/2 Are Associated with Grade Reclassification in Men on Active Surveillance for Prostate Cancer. <i>European Urology</i> , 2019, 75, 743-749.	1.9	138
13	A comprehensive evaluation of CHEK2 germline mutations in men with prostate cancer. <i>Prostate</i> , 2018, 78, 607-615.	2.3	57
14	Genome-wide Association Study (GWAS) of Germline Copy Number Variations (CNVs) Reveal Genetic Risks of Prostate Cancer in Chinese population. <i>Journal of Cancer</i> , 2018, 9, 923-928.	2.5	13
15	Germline mutations in PPF1B are associated with lethal prostate cancer. <i>Prostate</i> , 2018, 78, 1222-1228.	2.3	12
16	Germline mutations in DNA repair genes are associated with bladder cancer risk and unfavourable prognosis. <i>BJU International</i> , 2018, 122, 808-813.	2.5	15
17	A Cumulative Analysis of Current Evidence for Association between Expression of Epithelial-Mesenchymal Transition Markers and Clinicopathological Outcomes in Patients after Radical Prostatectomy. <i>Annals of Clinical and Laboratory Science</i> , 2018, 48, 18-28.	0.2	1
18	Germline Mutations in ATM and BRCA1/2 Distinguish Risk for Lethal and Indolent Prostate Cancer and are Associated with Early Age at Death. <i>European Urology</i> , 2017, 71, 740-747.	1.9	256

#	ARTICLE	IF	CITATIONS
19	Elevated hardness of peripheral gland on real-time elastography is an independent marker for high-risk prostate cancers. <i>Radiologia Medica</i> , 2017, 122, 944-951.	7.7	2
20	Prostate health index significantly reduced unnecessary prostate biopsies in patients with PSA 2-10 ng/mL and PSA >10 ng/mL: Results from a Multicenter Study in China. <i>Prostate</i> , 2017, 77, 1221-1229.	2.3	26
21	Validation of the novel susceptibility loci for prostate cancer in a Chinese population. <i>Oncology Letters</i> , 2017, 15, 2567-2573.	1.8	3
22	The preclinical assessment of XL388, a mTOR kinase inhibitor, as a promising anti-renal cell carcinoma agent. <i>Oncotarget</i> , 2017, 8, 30151-30161.	1.8	10
23	Germline genetic variations in PDZD2 and ITPR2 genes are associated with clear cell renal cell carcinoma in Chinese population. <i>Oncotarget</i> , 2017, 8, 24196-24201.	1.8	5
24	Race-specific genetic risk score is more accurate than nonrace-specific genetic risk score for predicting prostate cancer and high-grade diseases. <i>Asian Journal of Andrology</i> , 2016, 18, 525.	1.6	11
25	The effect of discrepancy between radiologic size and pathologic tumor size in renal cell cancer. <i>SpringerPlus</i> , 2016, 5, 899.	1.2	10
26	Genetic scores based on risk-associated single nucleotide polymorphisms (SNPs) can reveal inherited risk of renal cell carcinoma. <i>Oncotarget</i> , 2016, 7, 18631-18637.	1.8	7
27	Coexistence of YWHAZ amplification predicts better prognosis in muscle-invasive bladder cancer with CDKN2A or TP53 loss. <i>Oncotarget</i> , 2016, 7, 34752-34758.	1.8	6
28	Clinically available RNA profiling tests of prostate tumors: utility and comparison. <i>Asian Journal of Andrology</i> , 2016, 18, 575.	1.6	14
29	The Evaluation of the Risk Factors for Non-Muscle Invasive Bladder Cancer (NMIBC) Recurrence after Transurethral Resection (TURBt) in Chinese Population. <i>PLoS ONE</i> , 2015, 10, e0123617.	2.5	28
30	Plasma genistein and risk of prostate cancer in Chinese population. <i>International Urology and Nephrology</i> , 2015, 47, 965-970.	1.4	24