Mio Takeuchi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8226580/publications.pdf Version: 2024-02-01



Μιο Τλκειισμι

#	Article	IF	CITATIONS
1	Complete dominant inheritance of intracellular leucine accumulation traits in polyploid yeasts. Yeast, 2022, 39, 272-282.	1.7	2
2	Skin bacteria of rainbow trout antagonistic to the fish pathogen Flavobacterium psychrophilum. Scientific Reports, 2021, 11, 7518.	3.3	18
3	Acetate excretion by a methanotroph, <i>Methylocaldum marinum</i> S8, under aerobic conditions. Bioscience, Biotechnology and Biochemistry, 2021, 85, 2326-2333.	1.3	6
4	Microbial methane production and oxidation in the Holocene mud beneath the Kanto Plain of central Japan. Geochemical Journal, 2020, 54, 243-254.	1.0	3
5	Microbial community structure in deep natural gas-bearing aquifers subjected to sulfate-containing fluid injection. Journal of Bioscience and Bioengineering, 2019, 127, 45-51.	2.2	1
6	Possible cross-feeding pathway of facultative methylotroph Methyloceanibacter caenitepidi Gela4 on methanotroph Methylocaldum marinum S8. PLoS ONE, 2019, 14, e0213535.	2.5	22
7	Distribution of Dehalococcoides 16S rRNA and Dehalogenase Genes in Contaminated Sites. Environment and Natural Resources Research, 2017, 7, 37.	0.1	3
8	Tepidicaulis marinus gen. nov., sp. nov., a marine bacterium that reduces nitrate to nitrous oxide under strictly microaerobic conditions. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 1749-1754.	1.7	16
9	Methyloceanibacter caenitepidi gen. nov., sp. nov., a facultatively methylotrophic bacterium isolated from marine sediments near a hydrothermal vent. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 462-468.	1.7	54
10	Methylocaldum marinum sp. nov., a thermotolerant, methane-oxidizing bacterium isolated from marine sediments, and emended description of the genus Methylocaldum. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 3240-3246.	1.7	63
11	Estimating the viability of Chlorella exposed to oxidative stresses based around photocatalysis. International Biodeterioration and Biodegradation, 2013, 78, 1-6.	3.9	18
12	Carbon-14 age and chemical evolution of Ca(HCO3)2-type groundwater of age less than 8,000 years in a confined sandy and muddy Pleistocene aquifer, Japan. Hydrogeology Journal, 2013, 21, 1289-1305.	2.1	5
13	Carbon dioxide concentration dictates alternative methanogenic pathways in oil reservoirs. Nature Communications, 2013, 4, 1998.	12.8	98
14	Distribution and characterization of anammox in a swine wastewater activated sludge facility. Water Science and Technology, 2013, 67, 2330-2336.	2.5	8
15	Evidence for syntrophic acetate oxidation coupled to hydrogenotrophic methanogenesis in the highâ€ŧemperature petroleum reservoir of Yabase oil field (Japan). Environmental Microbiology, 2011, 13, 1995-2006.	3.8	114
16	A distinct freshwaterâ€edapted subgroup of ANMEâ€1 dominates active archaeal communities in terrestrial subsurfaces in Japan. Environmental Microbiology, 2011, 13, 3206-3218.	3.8	32
17	Comparative study of microbial dechlorination of chlorinated ethenes in an aquifer and a clayey aquitard. Journal of Contaminant Hydrology, 2011, 124, 14-24.	3.3	35
18	Characterization of humic acids in sediments from dam reservoirs by pyrolysis-gas chromatography/mass spectrometry using tetramethylammonium hydroxide: Influence of the structural features of humic acids on iron(II) binding capacity. Journal of Analytical and Applied Pyrolysis, 2011, 91, 323-331.	5.5	26

Μιο Τακευςηι

#	Article	IF	CITATIONS
19	Rate determination and distribution of anammox activity in activated sludge treating swine wastewater. Bioresource Technology, 2010, 101, 2685-2690.	9.6	42
20	Chemistry of fly ash and cyclone ash leachate from waste materials and effects of ash leachates on bacterial growth, nitrogen-transformation activity, and metal accumulation. Journal of Hazardous Materials, 2009, 165, 967-973.	12.4	7
21	Bacterial and Archaeal 16S rRNA Genes in Late Pleistocene to Holocene Muddy Sediments from the Kanto Plain of Japan. Geomicrobiology Journal, 2009, 26, 104-118.	2.0	21
22	Prevention of Phormidium tenue Biofilm Formation by TiO2 Photocatalysis. Microbes and Environments, 2009, 24, 241-245.	1.6	16
23	Role of Aquitard in Subsurface Microbial Activity. Journal of Geography (Chigaku Zasshi), 2007, 116, 915-921.	0.3	1
24	Arsenic resistance and removal by marine and non-marine bacteria. Journal of Biotechnology, 2007, 127, 434-442.	3.8	106
25	Impact of Changes in Redox Conditions on Leaching of Some Elements from MSW Fly Ash. Resource Geology, 2006, 56, 191-196.	0.8	2
26	Temperature and pH Dependence of Some Metals Leaching from Fly Ash of Municipal Solid Waste. Resource Geology, 2005, 55, 357-372.	0.8	12
27	Distribution and fate of biologically formed organoarsenicals in coastal marine sediment. Applied Organometallic Chemistry, 2005, 19, 945-951.	3.5	18
28	In situ bioremediation of a cis-dichloroethylene-contaminated aquifer utilizing methane-rich groundwater from an uncontaminated aquifer. Water Research, 2005, 39, 2438-2444.	11.3	18
29	Natural groundwater of a gas field utilizable for a bioremediation of trichloroethylene-contamination. Environmental Geology, 2004, 45, 891-898.	1.2	9
30	Distribution of Methanotrophs in Trichloroethylene-Contaminated Aquifers in a Natural Gas Field. Geomicrobiology Journal, 2001, 18, 387-399.	2.0	5