Andor Pivarcsi

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8232443/andor-pivarcsi-publications-by-year.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79
papers
6,587
citations
42
h-index
81
g-index

87
ext. papers
ext. citations
4
avg, IF
L-index

#	Paper	IF	Citations
79	Cross-talk between IFN-🗈nd TWEAK through miR-149 amplifies skin inflammation in psoriasis. Journal of Allergy and Clinical Immunology, 2021 , 147, 2225-2235	11.5	7
78	miR-19a/b and miR-20a Promote Wound Healing by Regulating the Inflammatory Response of Keratinocytes. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 659-671	4.3	19
77	Chromatin interactions in differentiating keratinocytes reveal novel atopic dermatitis- and psoriasis-associated genes. <i>Journal of Allergy and Clinical Immunology</i> , 2021 , 147, 1742-1752	11.5	1
76	MiR-130a Acts as a Tumor Suppressor MicroRNA in Cutaneous Squamous Cell Carcinoma and Regulates the Activity of the BMP/SMAD Pathway by Suppressing ACVR1. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 1922-1931	4.3	5
75	EGFR/Ras-induced CCL20 production modulates the tumour microenvironment. <i>British Journal of Cancer</i> , 2020 , 123, 942-954	8.7	12
74	A comprehensive analysis of coding and non-coding transcriptomic changes in cutaneous squamous cell carcinoma. <i>Scientific Reports</i> , 2020 , 10, 3637	4.9	35
73	Circulating microRNAs in extracellular vesicles as potential biomarkers for psoriatic arthritis in patients with psoriasis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020 , 34, 12	24 8 -125	66 ¹⁷
72	Next-Generation Sequencing Identifies the Keratinocyte-Specific miRNA Signature of Psoriasis. Journal of Investigative Dermatology, 2019 , 139, 2547-2550.e12	4.3	8
71	The Keratinocyte Transcriptome in Psoriasis: Pathways Related to Immune Responses, Cell Cycle and Keratinization. <i>Acta Dermato-Venereologica</i> , 2019 , 99, 196-205	2.2	23
70	Genome-Wide Screen for MicroRNAs Reveals a Role for miR-203 in Melanoma Metastasis. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 882-892	4.3	24
69	Extracellular microvesicle microRNAs as predictive biomarkers for targeted therapy in metastastic cutaneous malignant melanoma. <i>PLoS ONE</i> , 2018 , 13, e0206942	3.7	19
68	Tofacitinib Represses the Janus Kinase-Signal Transducer and Activators of Transcription Signalling Pathway in Keratinocytes. <i>Acta Dermato-Venereologica</i> , 2018 , 98, 772-775	2.2	6
67	Identification of chronological and photoageing-associated microRNAs in human skin. <i>Scientific Reports</i> , 2018 , 8, 12990	4.9	9
66	MicroRNA-132 promotes fibroblast migration via regulating RAS p21 protein activator 1 in skin wound healing. <i>Scientific Reports</i> , 2017 , 7, 7797	4.9	29
65	MicroRNA-132 with Therapeutic Potential in Chronic Wounds. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 2630-2638	4.3	49
64	MicroRNA-146a suppresses IL-17-mediated skin inflammation and is genetically associated with psoriasis. <i>Journal of Allergy and Clinical Immunology</i> , 2017 , 139, 550-561	11.5	79
63	MicroRNA-203 Inversely Correlates with Differentiation Grade, Targets c-MYC, and Functions as a Tumor Suppressor in cSCC. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 2485-2494	4.3	32

62	Exosomal microRNAs as putative predictive biomarkers for targeted therapy in stage IV cutaneous malignant melanoma (CMM) <i>Journal of Clinical Oncology</i> , 2016 , 34, 9579-9579	2.2	
61	MicroRNA-132 enhances transition from inflammation to proliferation during wound healing. <i>Journal of Clinical Investigation</i> , 2015 , 125, 3008-26	15.9	116
60	MicroRNA-31 Promotes Skin Wound Healing by Enhancing Keratinocyte Proliferation and Migration. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 1676-1685	4.3	101
59	Identification of novel non-coding RNA-based negative feedback regulating the expression of the oncogenic transcription factor GLI1. <i>Molecular Oncology</i> , 2014 , 8, 912-26	7.9	23
58	Activation of toll-like receptors alters the microRNA expression profile of keratinocytes. <i>Experimental Dermatology</i> , 2014 , 23, 281-3	4	22
57	miR-193b/365a cluster controls progression of epidermal squamous cell carcinoma. <i>Carcinogenesis</i> , 2014 , 35, 1110-20	4.6	60
56	MiR-146a negatively regulates TLR2-induced inflammatory responses in keratinocytes. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 1931-1940	4.3	75
55	MicroRNA-31 is overexpressed in cutaneous squamous cell carcinoma and regulates cell motility and colony formation ability of tumor cells. <i>PLoS ONE</i> , 2014 , 9, e103206	3.7	48
54	Genetic polymorphisms altering microRNA activity in psoriasisa key to solve the puzzle of missing heritability?. <i>Experimental Dermatology</i> , 2014 , 23, 620-4	4	26
53	Changes in the level of serum microRNAs in patients with psoriasis after antitumour necrosis factor-Therapy. <i>British Journal of Dermatology</i> , 2013 , 169, 563-70	4	65
52	MicroRNA-31 is overexpressed in psoriasis and modulates inflammatory cytokine and chemokine production in keratinocytes via targeting serine/threonine kinase 40. <i>Journal of Immunology</i> , 2013 , 190, 678-88	5.3	145
51	Are BIC (miR-155) polymorphisms associated with eczema susceptibility?. <i>Acta Dermato-Venereologica</i> , 2013 , 93, 366-7	2.2	5
50	Next-generation sequencing identifies microRNAs that associate with pathogenic autoimmune neuroinflammation in rats. <i>Journal of Immunology</i> , 2013 , 190, 4066-75	5.3	31
49	RNA editing of the GLI1 transcription factor modulates the output of Hedgehog signaling. <i>RNA Biology</i> , 2013 , 10, 321-33	4.8	56
48	Interleukin-8 is regulated by miR-203 at the posttranscriptional level in primary human keratinocytes. <i>European Journal of Dermatology</i> , 2013 ,	0.8	11
47	Constraints for monocyte-derived dendritic cell functions under inflammatory conditions. <i>European Journal of Immunology</i> , 2012 , 42, 458-69	6.1	13
46	Characterization of EGFR and ErbB2 expression in atopic dermatitis patients. <i>Archives of Dermatological Research</i> , 2012 , 304, 773-80	3.3	15
45	MiR-21 is up-regulated in psoriasis and suppresses T cell apoptosis. <i>Experimental Dermatology</i> , 2012 , 21, 312-4	4	116

44	MicroRNA-203 functions as a tumor suppressor in basal cell carcinoma. <i>Oncogenesis</i> , 2012 , 1, e3	6.6	71
43	MicroRNA-125b down-regulates matrix metallopeptidase 13 and inhibits cutaneous squamous cell carcinoma cell proliferation, migration, and invasion. <i>Journal of Biological Chemistry</i> , 2012 , 287, 29899-9	90 ⁵ 8 ⁴	141
42	MicroRNAs in inflammation and response to injuries induced by environmental pollution. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2011 , 717, 46-53	3.3	46
41	MiR-125b, a microRNA downregulated in psoriasis, modulates keratinocyte proliferation by targeting FGFR2. <i>Journal of Investigative Dermatology</i> , 2011 , 131, 1521-9	4.3	158
40	The expression of microRNA-203 during human skin morphogenesis. <i>Experimental Dermatology</i> , 2010 , 19, 854-6	4	50
39	MYCN-regulated microRNAs repress estrogen receptor-alpha (ESR1) expression and neuronal differentiation in human neuroblastoma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 1553-8	11.5	106
38	Protein kinase C-dependent upregulation of miR-203 induces the differentiation of human keratinocytes. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 124-34	4.3	98
37	MiR-155 is overexpressed in patients with atopic dermatitis and modulates T-cell proliferative responses by targeting cytotoxic T lymphocyte-associated antigen 4. <i>Journal of Allergy and Clinical Immunology</i> , 2010 , 126, 581-9.e1-20	11.5	208
36	A novel mechanism for anti-EGFR antibody action involves chemokine-mediated leukocyte infiltration. <i>International Journal of Cancer</i> , 2009 , 124, 2589-96	7.5	42
35	The human antimicrobial peptide LL-37 suppresses apoptosis in keratinocytes. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 937-44	4.3	63
34	microRNAs in inflammation. International Reviews of Immunology, 2009, 28, 535-61	4.6	170
33	Human antimicrobial protein hCAP18/LL-37 promotes a metastatic phenotype in breast cancer. Breast Cancer Research, 2009 , 11, R6	8.3	64
32	Advances in microRNAs: implications for immunity and inflammatory diseases. <i>Journal of Cellular and Molecular Medicine</i> , 2009 , 13, 24-38	5.6	127
31	Differential expression of D-type cyclins in HaCaT keratinocytes and in psoriasis. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 634-42	4.3	19
30	MicroRNAs: novel regulators in skin inflammation. Clinical and Experimental Dermatology, 2008, 33, 312	-5 .8	160
29	MicroRNAs and immunity: novel players in the regulation of normal immune function and inflammation. <i>Seminars in Cancer Biology</i> , 2008 , 18, 131-40	12.7	423
28	Toll-like receptor 9-independent suppression of skin inflammation by oligonucleotides. <i>Journal of Investigative Dermatology</i> , 2007 , 127, 746-8	4.3	6
27	Human adult epidermal melanocytes cultured without chemical mitogens express the EGF receptor and respond to EGF. <i>Archives of Dermatological Research</i> , 2007 , 299, 191-200	3.3	14

26	MicroRNAs: novel regulators involved in the pathogenesis of psoriasis?. <i>PLoS ONE</i> , 2007 , 2, e610	3.7	540
25	Tumor immune escape by the loss of homeostatic chemokine expression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 19055-60	11.5	109
24	IL-31: a new link between T cells and pruritus in atopic skin inflammation. <i>Journal of Allergy and Clinical Immunology</i> , 2006 , 117, 411-7	11.5	668
23	Budesonide, but not tacrolimus, affects the immune functions of normal human keratinocytes. <i>International Immunopharmacology</i> , 2006 , 6, 358-68	5.8	14
22	Chemokines Regulate Leukocyte Trafficking and Organ-specific Metastasis 2006 , 153-166		
21	The expression of keratinocyte growth factor receptor (FGFR2-IIIb) correlates with the high proliferative rate of HaCaT keratinocytes. <i>Experimental Dermatology</i> , 2006 , 15, 596-605	4	10
20	Propionibacterium acnes and lipopolysaccharide induce the expression of antimicrobial peptides and proinflammatory cytokines/chemokines in human sebocytes. <i>Microbes and Infection</i> , 2006 , 8, 2195-2	283	273
19	Microbial compounds induce the expression of pro-inflammatory cytokines, chemokines and human beta-defensin-2 in vaginal epithelial cells. <i>Microbes and Infection</i> , 2005 , 7, 1117-27	9.3	116
18	Distinct strains of Propionibacterium acnes induce selective human beta-defensin-2 and interleukin-8 expression in human keratinocytes through toll-like receptors. <i>Journal of Investigative Dermatology</i> , 2005 , 124, 931-8	4.3	241
17	Chemokine networks in atopic dermatitis: traffic signals of disease. <i>Current Allergy and Asthma Reports</i> , 2005 , 5, 284-90	5.6	59
16	Identification and characterization of a novel, psoriasis susceptibility-related noncoding RNA gene, PRINS. <i>Journal of Biological Chemistry</i> , 2005 , 280, 24159-67	5.4	152
15	Innate Immunity in the Skin: How Keratinocytes Fight Against Pathogens. <i>Current Immunology Reviews</i> , 2005 , 1, 29-42	1.3	55
14	CCL1-CCR8 interactions: an axis mediating the recruitment of T cells and Langerhans-type dendritic cells to sites of atopic skin inflammation. <i>Journal of Immunology</i> , 2005 , 174, 5082-91	5.3	162
13	CC chemokine ligand 18, an atopic dermatitis-associated and dendritic cell-derived chemokine, is regulated by staphylococcal products and allergen exposure. <i>Journal of Immunology</i> , 2004 , 173, 5810-7	5.3	101
12	Negative regulatory effect of histamine in DNFB-induced contact hypersensitivity. <i>International Immunology</i> , 2004 , 16, 1781-8	4.9	14
11	Proliferating keratinocytes are putative sources of the psoriasis susceptibility-related EDA+ (extra domain A of fibronectin) oncofetal fibronectin. <i>Journal of Investigative Dermatology</i> , 2004 , 123, 537-46	4.3	34
10	Differentiation-regulated expression of Toll-like receptors 2 and 4 in HaCaT keratinocytes. <i>Archives of Dermatological Research</i> , 2004 , 296, 120-4	3.3	38
9	Innate immune functions of the keratinocytes. A review. <i>Acta Microbiologica Et Immunologica Hungarica</i> , 2004 , 51, 303-10	1.8	50

8	The role of innate immunity in the pathogenesis of acne. <i>Dermatology</i> , 2003 , 206, 96-105	4.4	112
7	Expression and function of Toll-like receptors 2 and 4 in human keratinocytes. <i>International Immunology</i> , 2003 , 15, 721-30	4.9	248
6	Hemese, a hemocyte-specific transmembrane protein, affects the cellular immune response in Drosophila. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 2622-7	11.5	131
5	A MANNOSE-BINDING RECEPTOR IS EXPRESSED ON HUMAN KERATINOCYTES AND MEDIATES KILLING OF CANDIDA ALBICANS. <i>Mycoses</i> , 2002 , 45, 30-31	5.2	
4	Dithranol upregulates IL-10 receptors on the cultured human keratinocyte cell line HaCaT. <i>Inflammation Research</i> , 2001 , 50, 44-9	7.2	41
3	Serum factors regulate the expression of the proliferation-related genes alpha5 integrin and keratin 1, but not keratin 10, in HaCaT keratinocytes. <i>Archives of Dermatological Research</i> , 2001 , 293, 206-13	3.3	34
2	A mannose-binding receptor is expressed on human keratinocytes and mediates killing of Candida albicans. <i>Journal of Investigative Dermatology</i> , 2001 , 117, 205-13	4.3	69
1	Histidine decarboxylase expression in human melanoma. <i>Journal of Investigative Dermatology</i> , 2000 , 115, 345-52	4.3	45