
## Nunzio Pomara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8236223/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Drug-induced reductions in brain amyloid-β levels may adversely affect cognition and behavior by a disruption of functional connectivity homeostasis. Neurodegenerative Disease Management, 2019, 9, 189-191.	1.2	1
2	Basal Forebrain Resting Functional Activity Associated with Brain Amyloid-β Burden in ε4 Carriers and Older Women: Possible Implications for Baseline Cognition and Response to Amyloid-β-based Preventive Trials. Radiology, 2019, 290, 850-851.	3.6	2
3	The recency ratio as predictor of early MCI. International Psychogeriatrics, 2018, 30, 1883-1888.	0.6	23
4	Pathological Increases in Neuronal Hyperactivity in Selective Cholinergic and Noradrenergic Pathways May Limit the Efficacy of Amyloid-β-Based Interventions in Mild Cognitive Impairment and Alzheimer's Disease. Journal of Alzheimer's Disease Reports, 2018, 2, 165-167.	1.2	3
5	Basal forebrain mediated increase in brain CRF is associated with increased cholinergic tone and depression. Psychiatry Research - Neuroimaging, 2017, 264, 76-81.	0.9	0
6	The recency ratio is associated with reduced CSF glutamate in late-life depression. Neurobiology of Learning and Memory, 2017, 141, 14-18.	1.0	10
7	The Primacy Effect in Amnestic Mild Cognitive Impairment: Associations with Hippocampal Functional Connectivity. Frontiers in Aging Neuroscience, 2016, 8, 244.	1.7	18
8	Output order and variability in free recall are linked to cognitive ability and hippocampal volume in elderly individuals. Neuropsychologia, 2016, 80, 126-132.	0.7	9
9	Neuronal correlates of serial position performance in amnestic mild cognitive impairment Neuropsychology, 2016, 30, 906-914.	1.0	15
10	A study on the specificity of the association between hippocampal volume and delayed primacy performance in cognitively intact elderly individuals. Neuropsychologia, 2015, 69, 1-8.	0.7	25
11	The relationship between cerebrospinal fluid tau markers, hippocampal volume, and delayed primacy performance in cognitively intact elderly individuals. Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring, 2015, 1, 81-86.	1.2	7
12	Hippocampus and basal forebrain volumes modulate effects ofÂanticholinergic treatment on delayed recall in healthy older adults. Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring, 2015, 1, 216-219.	1.2	4
13	A review of the relationship between proinflammatory cytokines and major depressive disorder. Journal of Affective Disorders, 2014, 169, 15-20.	2.0	329
14	Lower plasma Â-amyloid levels are associated with moderately greater rate of cognitive decline among older people without dementia. Evidence-Based Mental Health, 2011, 14, 41-41.	2.2	2
15	Prospective strategies for cholinergic interventions in Alzheimer's disease. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 1986, 10, 553-569.	2.5	8