

Mohammad Aminur Rahman

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8237903/publications.pdf>

Version: 2024-02-01

18
papers

429
citations

933447

10
h-index

1125743

13
g-index

20
all docs

20
docs citations

20
times ranked

780
citing authors

#	ARTICLE	IF	CITATIONS
1	Platelet-derived growth factor receptor $\hat{\pm}$ /glial fibrillary acidic protein expressing peritumoral astrocytes associate with shorter median overall survival in glioblastoma patients. <i>Glia</i> , 2020, 68, 979-988.	4.9	4
2	Sequential bortezomib and temozolomide treatment promotes immunological responses in glioblastoma patients with positive clinical outcomes: A phase 1B study. <i>Immunity, Inflammation and Disease</i> , 2020, 8, 342-359.	2.7	19
3	Bortezomib administered prior to temozolomide depletes MGMT, chemosensitizes glioblastoma with unmethylated MGMT promoter and prolongs animal survival. <i>British Journal of Cancer</i> , 2019, 121, 545-555.	6.4	49
4	Pretreatment of Glioblastoma with Bortezomib Potentiates Natural Killer Cell Cytotoxicity through TRAIL/DR5 Mediated Apoptosis and Prolongs Animal Survival. <i>Cancers</i> , 2019, 11, 996.	3.7	28
5	Glioblastoma Stem-Like Cells Are More Susceptible Than Differentiated Cells to Natural Killer Cell Lysis Mediated Through Killer Immunoglobulin-Like Receptorsâ€™Human Leukocyte Antigen Ligand Mismatch and Activation Receptorâ€™Ligand Interactions. <i>Frontiers in Immunology</i> , 2018, 9, 1345.	4.8	52
6	Tumour-associated glial host cells display a stem-like phenotype with a distinct gene expression profile and promote growth of GBM xenografts. <i>BMC Cancer</i> , 2017, 17, 108.	2.6	11
7	Tumor-Host Interactions in Malignant Gliomas. , 2017, , 465-479.		0
8	Increased infiltration and tolerised antigen-specific CD8+ TEM cells in tumor but not peripheral blood have no impact on survival of HCMV+ glioblastoma patients. <i>Oncolimmunology</i> , 2017, 6, e1336272.	4.6	17
9	Identification of a Natural Killer Cell Receptor Allele That Prolongs Survival of Cytomegalovirus-Positive Glioblastoma Patients. <i>Cancer Research</i> , 2016, 76, 5326-5336.	0.9	26
10	Treatment with the PI3K inhibitor buparlisib (NVP-BKM120) suppresses the growth of established patient-derived GBM xenografts and prolongs survival in nude rats. <i>Journal of Neuro-Oncology</i> , 2016, 129, 57-66.	2.9	25
11	Abstract 1560: FGFR4 is expressed in the tumor and stromal compartments of human gliomas of all grades and histologies. , 2015, , .		0
12	Abstract 2101: The transcription factor POU3F2 is expressed in human gliomas and promotes tumorigenesis in vivo. , 2015, , .		2
13	The ASH1-RELATED3 SET-Domain Protein Controls Cell Division Competence of the Meristem and the Quiescent Center of the Arabidopsis Primary Root $\hat{\text{A}} \hat{\text{A}}$. <i>Plant Physiology</i> , 2014, 166, 632-643.	4.8	35
14	The <i>Arabidopsis</i> Histone Methyltransferase SUVR4 Binds Ubiquitin via a Domain with a Four-Helix Bundle Structure. <i>Biochemistry</i> , 2014, 53, 2091-2100.	2.5	7
15	Abstract LB-75: Oct7 is expressed in human gliomas and correlates with malignancy grade. , 2014, , .		0
16	The SUVR4 Histone Lysine Methyltransferase Binds Ubiquitin and Converts H3K9me1 to H3K9me3 on Transposon Chromatin in Arabidopsis. <i>PLoS Genetics</i> , 2011, 7, e1001325.	3.5	49
17	The CW domain, a new histone recognition module in chromatin proteins. <i>EMBO Journal</i> , 2011, 30, 1939-1952.	7.8	105
18	Arabidopsis SET-domain proteins with different histone methyltransferase activity. <i>Acta Crystallographica Section A: Foundations and Advances</i> , 2009, 65, s157-s158.	0.3	0