

# Mårten FernÅ

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8243998/publications.pdf>

Version: 2024-02-01

47  
papers

1,720  
citations

331259

21  
h-index

288905

40  
g-index

49  
all docs

49  
docs citations

49  
times ranked

3312  
citing authors

#	ARTICLE	IF	CITATIONS
1	Breast cancer hypoxia in relation to prognosis and benefit from radiotherapy after breast-conserving surgery in a large, randomised trial with long-term follow-up. <i>British Journal of Cancer</i> , 2022, 126, 1145-1156.	2.9	20
2	Abstract PD9-05: Prognostic and tamoxifen-predictive effect of PAM50 and ROR score in premenopausal women included in the randomised SBII:2 trial. <i>Cancer Research</i> , 2022, 82, PD9-05-PD9-05.	0.4	0
3	The Prognostic Role of Intratumoral Stromal Content in Lobular Breast Cancer. <i>Cancers</i> , 2022, 14, 941.	1.7	5
4	PAM50 subtyping and ROR score add long-term prognostic information in premenopausal breast cancer patients. <i>Npj Breast Cancer</i> , 2022, 8, 61.	2.3	5
5	Abstract PS5-09: Tumour infiltrating lymphocytes of prognostic value in different molecular breast cancer subgroups and as a suggestive predictive factor for adjuvant tamoxifen benefit in premenopausal patients after 30 years follow-up. , 2021, , .		0
6	Prognostic and predictive impact of stroma cells defined by PDGFRb expression in early breast cancer: results from the randomized SweBCG91RT trial. <i>Breast Cancer Research and Treatment</i> , 2021, 187, 45-55.	1.1	8
7	High CYP27A1 expression is a biomarker of favorable prognosis in premenopausal patients with estrogen receptor positive primary breast cancer. <i>Npj Breast Cancer</i> , 2021, 7, 127.	2.3	9
8	Disseminated tumour cells from the bone marrow of early breast cancer patients: Results from an international pooled analysis. <i>European Journal of Cancer</i> , 2021, 154, 128-137.	1.3	24
9	The Distribution of Circulating Tumor Cells Is Different in Metastatic Lobular Compared to Ductal Carcinoma of the Breast—Long-Term Prognostic Significance. <i>Cells</i> , 2020, 9, 1718.	1.8	10
10	Expression of HGF, pMet, and pAkt is related to benefit of radiotherapy after breast-conserving surgery: a long-term follow-up of the SweBCG91RT randomised trial. <i>Molecular Oncology</i> , 2020, 14, 2713-2726.	2.1	2
11	Co-localization of CD169 macrophages and cancer cells in lymph node metastases of breast cancer patients is linked to improved prognosis and PDL1 expression. <i>Oncolmmunology</i> , 2020, 9, 1848067.	2.1	9
12	Comprehensive Transcriptomic Profiling Identifies Breast Cancer Patients Who May Be Spared Adjuvant Systemic Therapy. <i>Clinical Cancer Research</i> , 2020, 26, 171-182.	3.2	14
13	Plasma membrane expression of G protein-coupled estrogen receptor (GPER)/G protein-coupled receptor 30 (GPR30) is associated with worse outcome in metachronous contralateral breast cancer. <i>PLoS ONE</i> , 2020, 15, e0231786.	1.1	15
14	Expression of epithelial-mesenchymal transition-related markers and phenotypes during breast cancer progression. <i>Breast Cancer Research and Treatment</i> , 2020, 181, 369-381.	1.1	25
15	Tumour-infiltrating lymphocytes as a prognostic and tamoxifen predictive marker in premenopausal breast cancer: data from a randomised trial with long-term follow-up. <i>Breast Cancer Research</i> , 2020, 22, 140.	2.2	25
16	Agreement between molecular subtyping and surrogate subtype classification: a contemporary population-based study of ER-positive/HER2-negative primary breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019, 178, 459-467.	1.1	23
17	Refinement of breast cancer molecular classification by miRNA expression profiles. <i>BMC Genomics</i> , 2019, 20, 503.	1.2	75
18	The estrogen receptor coactivator AIB1 is a new putative prognostic biomarker in ER-positive/HER2-negative invasive lobular carcinoma of the breast. <i>Breast Cancer Research and Treatment</i> , 2019, 175, 305-316.	1.1	8

#	ARTICLE	IF	CITATIONS
19	Assessment of early response biomarkers in relation to long-term survival in patients with HER2-negative breast cancer receiving neoadjuvant chemotherapy plus bevacizumab: Results from the Phase II PROMIX trial. <i>International Journal of Cancer</i> , 2018, 142, 618-628.	2.3	27
20	Macrophage-derived lipocalin-2 transports iron in the tumor microenvironment. <i>Oncolmmunology</i> , 2018, 7, e1408751.	2.1	64
21	Non-linear transformations of age at diagnosis, tumor size, and number of positive lymph nodes in prediction of clinical outcome in breast cancer. <i>BMC Cancer</i> , 2018, 18, 1226.	1.1	3
22	Stability of oestrogen and progesterone receptor antigenicity in formalin-fixed paraffin-embedded breast cancer tissue over time. <i>Apmis</i> , 2018, 126, 746-754.	0.9	4
23	Identification and validation of single-sample breast cancer radiosensitivity gene expression predictors. <i>Breast Cancer Research</i> , 2018, 20, 64.	2.2	40
24	Comprehensive transcriptomic profiling to identify breast cancer patients that may be spared adjuvant systemic therapy.. <i>Journal of Clinical Oncology</i> , 2018, 36, 535-535.	0.8	1
25	PAM50 Provides Prognostic Information When Applied to the Lymph Node Metastases of Advanced Breast Cancer Patients. <i>Clinical Cancer Research</i> , 2017, 23, 7225-7231.	3.2	17
26	Histological grade provides significant prognostic information in addition to breast cancer subtypes defined according to St Gallen 2013. <i>Acta Oncologica</i> , 2017, 56, 68-74.	0.8	51
27	Response to Radiotherapy After Breast-Conserving Surgery in Different Breast Cancer Subtypes in the Swedish Breast Cancer Group 91 Radiotherapy Randomized Clinical Trial. <i>Journal of Clinical Oncology</i> , 2017, 35, 3222-3229.	0.8	74
28	Tumor tissue protein signatures reflect histological grade of breast cancer. <i>PLoS ONE</i> , 2017, 12, e0179775.	1.1	8
29	Two Years of Adjuvant Tamoxifen Provides a Survival Benefit Compared With No Systemic Treatment in Premenopausal Patients With Primary Breast Cancer: Long-Term Follow-Up (> 25 years) of the Phase III SBl:2pre Trial. <i>Journal of Clinical Oncology</i> , 2016, 34, 2232-2238.	0.8	30
30	Transcriptional Profiling of Breast Cancer Metastases Identifies Liver Metastasis-Selective Genes Associated with Adverse Outcome in Luminal A Primary Breast Cancer. <i>Clinical Cancer Research</i> , 2016, 22, 146-157.	3.2	38
31	Prior Adjuvant Tamoxifen Treatment in Breast Cancer Is Linked to Increased AIB1 and HER2 Expression in Metachronous Contralateral Breast Cancer. <i>PLoS ONE</i> , 2016, 11, e0150977.	1.1	9
32	Protease Activated Receptors 1 and 2 Correlate Differently with Breast Cancer Aggressiveness Depending on Tumor ER Status. <i>PLoS ONE</i> , 2015, 10, e0134932.	1.1	7
33	Serial monitoring of circulating tumor DNA in patients with primary breast cancer for detection of occult metastatic disease. <i>EMBO Molecular Medicine</i> , 2015, 7, 1034-1047.	3.3	380
34	Stem cell biomarker ALDH1A1 in breast cancer shows an association with prognosis and clinicopathological variables that is highly cut-off dependent. <i>Journal of Clinical Pathology</i> , 2015, 68, 1012-1019.	1.0	15
35	Prognosis, stage and oestrogen receptor status of contralateral breast cancer in relation to characteristics of the first tumour, prior endocrine treatment and radiotherapy. <i>European Journal of Cancer</i> , 2015, 51, 2304-2313.	1.3	8
36	A Molecular Pathologic Framework for Risk Stratification of Stage T1 Urothelial Carcinoma. <i>European Urology</i> , 2015, 68, 824-832.	0.9	111

#	ARTICLE	IF	CITATIONS
37	Loss of TGF $\beta$ 2 Receptor Type 2 Expression Impairs Estrogen Response and Confers Tamoxifen Resistance. <i>Cancer Research</i> , 2015, 75, 1457-1469.	0.4	26
38	Contralateral breast cancer can represent a metastatic spread of the first primary tumor: determination of clonal relationship between contralateral breast cancers using next-generation whole genome sequencing. <i>Breast Cancer Research</i> , 2015, 17, 102.	2.2	30
39	Contrasting breast cancer molecular subtypes across serial tumor progression stages: biological and prognostic implications. <i>Oncotarget</i> , 2015, 6, 33306-33318.	0.8	31
40	The Three Receptor Tyrosine Kinases c-KIT, VEGFR2 and PDGFR $\beta$ , Closely Spaced at 4q12, Show Increased Protein Expression in Triple-Negative Breast Cancer. <i>PLoS ONE</i> , 2014, 9, e102176.	1.1	49
41	Invasive lobular carcinoma of the breast: long-term prognostic value of Ki67 and histological grade, alone and in combination with estrogen receptor. <i>SpringerPlus</i> , 2014, 3, 70.	1.2	14
42	Infiltration of CD3+ and CD68+ cells in bladder cancer is subtype specific and affects the outcome of patients with muscle-invasive tumors11Grant support: The Swedish Cancer Society, the Swedish research council, the Nilsson Cancer foundation, the BioCARE Strategic Cancer Research program, the Lund Medical Faculty, and FoU Landstinget Kronoberg and SÄdra RegionvÄrdnÄmnden.. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 791-797.	0.8	106
43	Claudin $\beta$ 2 is an independent negative prognostic factor in breast cancer and specifically predicts early liver recurrences. <i>Molecular Oncology</i> , 2014, 8, 119-128.	2.1	61
44	Tumor-Specific Expression of Vascular Endothelial Growth Factor Receptor 2 but Not Vascular Endothelial Growth Factor or Human Epidermal Growth Factor Receptor 2 Is Associated With Impaired Response to Adjuvant Tamoxifen in Premenopausal Breast Cancer. <i>Journal of Clinical Oncology</i> , 2005, 23, 4695-4704.	0.8	80
45	Two years of adjuvant tamoxifen in premenopausal patients with breast cancer: a randomised, controlled trial with long-term follow-up. <i>European Journal of Cancer</i> , 2005, 41, 256-264.	1.3	92
46	Correlation between estrogen receptor $\beta$ expression, collagen content and stiffness in human uterine arteries. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2003, 82, 610-615.	1.3	7
47	Estrogen and progesterone receptors in vessel walls: Biochemical and immunochemical assays. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1993, 72, 10-16.	1.3	57