

Jacqueline K Harris

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8244282/publications.pdf>

Version: 2024-02-01

13
papers

205
citations

1306789

7
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

394
citing authors

#	ARTICLE	IF	CITATIONS
1	Hippocampal tail volume as a predictive biomarker of antidepressant treatment outcomes in patients with major depressive disorder: a CAN-BIND report. <i>Neuropsychopharmacology</i> , 2020, 45, 283-291.	2.8	37
2	The Canadian Biomarker Integration Network in Depression (CAN-BIND): magnetic resonance imaging protocols. <i>Journal of Psychiatry and Neuroscience</i> , 2019, 44, 223-236.	1.4	37
3	Testing a deep convolutional neural network for automated hippocampus segmentation in a longitudinal sample of healthy participants. <i>NeuroImage</i> , 2019, 197, 589-597.	2.1	24
4	Clinical, behavioral, and neural measures of reward processing correlate with escitalopram response in depression: a Canadian Biomarker Integration Network in Depression (CAN-BIND-1) Report. <i>Neuropsychopharmacology</i> , 2020, 45, 1390-1397.	2.8	23
5	White Matter Indices of Medication Response in Major Depression: A Diffusion Tensor Imaging Study. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 913-924.	1.1	21
6	Reduced accuracy accompanied by reduced neural activity during the performance of an emotional conflict task by unmedicated patients with major depression: A CAN-BIND fMRI study. <i>Journal of Affective Disorders</i> , 2019, 257, 765-773.	2.0	20
7	Structural covariance pattern abnormalities of insula in major depressive disorder: A CAN-BIND study report. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 111, 110194.	2.5	11
8	Exploring brain connectivity changes in major depressive disorder using functional structural data fusion: A CAN-BIND study. <i>Human Brain Mapping</i> , 2021, 42, 4940-4957.	1.9	8
9	Reliability of a functional magnetic resonance imaging task of emotional conflict in healthy participants. <i>Human Brain Mapping</i> , 2020, 41, 1400-1415.	1.9	7
10	Escitalopram ameliorates differences in neural activity between healthy comparison and major depressive disorder groups on an fMRI Emotional conflict task: A CAN-BIND-1 study. <i>Journal of Affective Disorders</i> , 2020, 264, 414-424.	2.0	6
11	Cerebello-limbic functional connectivity patterns in youth at clinical high risk for psychosis. <i>Schizophrenia Research</i> , 2022, 240, 220-227.	1.1	6
12	White matter microstructure in youth at risk for serious mental illness: A comparative analysis. <i>Psychiatry Research - Neuroimaging</i> , 2021, 312, 111289.	0.9	4
13	Association between the expression of lncRNA BASP-AS1 and volume of right hippocampal tail moderated by episode duration in major depressive disorder: a CAN-BIND 1 report. <i>Translational Psychiatry</i> , 2021, 11, 469.	2.4	1